



Source details

Advances in the Theory of Nonlinear Analysis and its Applications

Open Access ⓘ

Scopus coverage years: from 2019 to Present

Publisher: Erdal Karapinar

E-ISSN: 2587-2648

Subject area: Mathematics: Analysis Mathematics: Applied Mathematics

Source type: Journal

[View all documents >](#)[Set document alert](#)[Save to source list](#) [Source Homepage](#)

CiteScore 2020

0.5 ⓘ

SJR 2020

0.278 ⓘ

SNIP 2020

0.666 ⓘ

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

Improved CiteScore methodology

CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore 2020

0.5 = 32 Citations 2017 - 2020
61 Documents 2017 - 2020

Calculated on 05 May, 2021

CiteScoreTracker 2021 ⓘ

1.0 = 73 Citations to date
74 Documents to date

Last updated on 04 July, 2021 • Updated monthly

CiteScore rank 2020 ⓘ

Category	Rank	Percentile
Mathematics		
Analysis	#141/164	14th
Mathematics		
Applied Mathematics	#482/548	12th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site >](#)

About Scopus

[What is Scopus](#)
[Content coverage](#)
[Scopus blog](#)
[Scopus API](#)
[Privacy matters](#)

Language

[日本語に切り替える](#)
[切换到简体中文](#)
[切换到繁體中文](#)
[Русский язык](#)

Customer Service

[Help](#)
[Contact us](#)

Volume: 5 - Issue: 4

👁 353 | 📄 303

Contents

📄 Articles

Strong and weak convergences in 2-probabilistic normed spaces (<https://dergipark.org.tr/en/pub/atnaa/issue/62841/921900>)
 / Pages : 454-466 PDF (/en/download/article-file/1718882)
 Harikrishnan PANACKAL, Bernardo GUILLEN, Ravi AGARWAL, Hamid MORADI

Hagen-Poiseuille Flow in Circular Cylinder when Temperature is Exponential and Sinusoidal Function of Length (<https://dergipark.org.tr/en/pub/atnaa/issue/62841/954432>) / Pages : 467-481 PDF (/en/download/article-file/1831871)
 Radhika KHANDELWAL, Surendra AGARWAL

New Faster Four Step Iterative Algorithm for Suzuki Generalized Nonexpansive Mappings With an Application (<https://dergipark.org.tr/en/pub/atnaa/issue/62841/869046>) / Pages : 482-506 PDF (/en/download/article-file/1536094)
 Austine OFEM, Donatus IGBOKWE

Inertial hybrid self-adaptive subgradient extragradient method for fixed point of quasi- ϕ -nonexpansive multivalued mappings and equilibrium problem (<https://dergipark.org.tr/en/pub/atnaa/issue/62841/822150>) / Pages : 507-522 PDF (/en/download/article-file/1383131)
 Murtala HARBAU

Existence of an initial value problem for time-fractional Oldroyd-B fluid equation using Banach fixed point theorem (<https://dergipark.org.tr/en/pub/atnaa/issue/62841/943242>) / Pages : 523-530 PDF (/en/download/article-file/1789376)
 Vo Viet TRI

Common Fixed Point Theorem for Hybrid Pair of Mappings in a Generalized (F, ξ, η) -contraction in weak Partial b -Metric Spaces with an Application (<https://dergipark.org.tr/en/pub/atnaa/issue/62841/934778>) / Pages : 531-550 PDF (/en/download/article-file/1758826)
 Lucas WANGWE, Santosh KUMAR

Note on abstract elliptic equations with nonlocal boundary in time condition (<https://dergipark.org.tr/en/pub/atnaa/issue/62841/925768>) / Pages : 551-558 PDF (/en/download/article-file/1727089)
 Kim Van HO THI

Some new fixed point results for monotone enriched nonexpansive mappings in ordered Banach spaces (<https://dergipark.org.tr/en/pub/atnaa/issue/62841/954446>) / Pages : 559-567 PDF (/en/download/article-file/1831917)
 Rahul SHUKLA, Rajendra PANT

📌 On efficient matrix-free method via quasi-Newton approach for solving system of nonlinear equations (<https://dergipark.org.tr/en/pub/atnaa/issue/62841/890281>) / Pages : 568-579 PDF (/en/download/article-file/1614839) 📌
 Muhammad ABDULLAHI, Abubakar HALILU, Aliyu AWWAL, Nuttapol PAKKARANANG

News!

We are happy to announce that our journal “Advances in the Theory of Nonlinear Analysis and its Applications (ATNAA)”

has been indexed /abstracted by

SCOPUS

zbMATH

DOAJ

The journal of “Advances in the Theory of Nonlinear Analysis and its Applications(ATNAA)” was founded on April 2017 as an electronic scientific peer-reviewed journal according to international standards.

ATNAA is an Open Access journal devoted exclusively to the publication of high-quality research papers in the fields of nonlinear analysis.

It is an open access quarterly journal without any page charge.

ISSN : 2587-2648 (Online)

Alternative links:

<https://atnaa.wordpress.com/>

<http://atnaa.org/>

Erdal KARAPINAR © Contact (<https://dergipark.org.tr/en/pub/atnaa/contacts>)
 ULAKBİM Dergi Sistemleri (/dergipark.org.tr/en/)

Erdal KARAPINAR © Contact (<https://dergipark.org.tr/en/pub/atnaa/contacts>)

ULAKBİM Dergi Sistemleri (/dergipark.org.tr/en/)



On efficient matrix-free method via quasi-Newton approach for solving system of nonlinear equations

Muhammad Abdullahi^a, Abubakar Sani Halilu^a, Aliyu Muhammed Awwal^b, Nuttapol Pakkaranang^c

^aDepartment of Mathematics and Computer Sciences, Sule Lamido University, Kafin Hausa, Nigeria.

^bDepartment of Mathematics, Faculty of Science, Gombe State University, Gombe, Nigeria.

^cDepartment of Mathematics, Faculty of Science and Technology, Phetchabun Rajabhat University, Phetchabun 67000, Thailand.

Abstract

In this paper, a matrix-free method for solving large-scale system of nonlinear equations is presented. The method is derived via quasi-Newton approach, where the approximation to the Broyden's update is done by constructing diagonal matrix using acceleration parameter. A fascinating feature of the method is that it is a matrix-free, so is suitable for solving large-scale problems. Furthermore, the convergence analysis of the new method is discussed based on some standard condition. Preliminary numerical results on some test problems show that the method is promising.

Keywords: Matrix-free, Descent direction, Global convergence, Acceleration parameter.

2010 MSC: 65H11, 65K05, 65H12, 65H18.

1. Introduction

Many of problems in sciences, engineering and economics can be expressed as optimization problems or nonlinear system of equations, which are usually solved using iterative methods. This paper focuses on the following system

$$F(x) = 0, \quad (1)$$

where $x \in \mathbb{R}^n$ and the nonlinear function $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous. Throughout this paper, the symbol $\mathbb{R}^n$ denotes the n -dimensional real space equipped with the Euclidean norm $\|\cdot\|$, $F_k = F(x_k)$ where $x_k \in \mathbb{R}^n$ is the point at certain iteration $k = 1, 2, \dots$

Email addresses: muhammad.abdullahi@slu.edu.ng (Muhammad Abdullahi), abubakarsani.halilu@slu.edu.ng (Abubakar Sani Halilu), aliyumagsu@gmail.com (Aliyu Muhammed Awwal), nuttapol.pak@pcru.ac.th (Nuttapol Pakkaranang)

Received : March 3, 2021; *Accepted:* July 8, 2021; *Online:* July 10, 2021

Moreover, the system (1) can be obtained from general unconstrained optimization problems [9]. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a merit function defined by

$$f(x) = \frac{1}{2} \|F(x)\|^2, \quad x \in \mathbb{R}^n. \quad (2)$$

Then the nonlinear equations problem (1) is equivalent to the following unconstrained optimization problem

$$\min f(x), \quad x \in \mathbb{R}^n.$$

The study of such mappings is applied in a variety of scientific areas, including economic and chemical equilibrium systems [37, 39, 38]. Some iterative methods for solving these problems include Newton method [41], the quasi-Newton methods [4, 6, 9], the Levenberg-Marquardt methods [42, 43], the double direction methods [23, 33, 29], The double step length methods [17, 18, 28], and derivative-free methods [34, 45, 46, 44]. But, the famous method used to solve (1) is Newton method that determines the search direction d_k by solving the following linear system of equations,

$$F_k + F'_k d_k = 0, \quad (3)$$

where F'_k is the Jacobian matrix of F at x_k . The Newton method is appealing because it converges quadratically from a reasonably good starting point [14]. Despite its excellent convergence property, the method has some shortcomings, which includes storing of Jacobian matrix and solving system of linear equations in every iteration. In order to overcome some of the challenges associated with Newton method, alternatives such as quasi-Newton methods have been developed [4, 6]. These Methods avoid the computation of the exact Jacobian matrix and a matrix which is an approximation the Jacobian matrix or its inverse is used instead. This matrix is there by updated in every iteration. It has been shown that most of the quasi-Newton methods have superlinear order of convergence [14]. One of the successful quasi-Newton method, known as Broyden's method, generates a sequence of iterates $\{x_k\}$ using

$$x_{k+1} = x_k - B_k^{-1} F_k, \quad k = 0, 1, 2, \dots, \quad (4)$$

where the Broyden matrix B_k is the approximation of the Jacobian matrix, such that the following quasi-Newton equation

$$B_{k+1}(x_{k+1} - x_k) = F_{k+1} - F_k, \quad (5)$$

is satisfied for all k . It is important to note that Broyden's method requires the computation and storage of $n \times n$ matrix at every iteration. Therefore, for large-scale problems, this could result to serious memory constraints. Efforts have been made by different researchers to reduce the storage problem associated with quasi-Newton methods. For instance, some modifications of the Broyden's method have been done in the literature in order to reduce its computational cost [5, 8, 11, 12]. These methods are usually referred to as limited memory Broyden methods [12, 16].

As mentioned earlier, the quasi-Newton methods has contributed in overcoming of the shortcomings of Newton's method which is computing Jacobian matrix in every iteration. However, the prize paid by the quasi-Newton method is that only superlinear rate of convergence can be achieved instead of quadratic rate. In order to improve the convergence order of quasi-Newton method, many higher order approaches have been proposed. There is a great deal of literature on the family of derivative-free methods used to solve nonlinear equations. In [16], a family of conjugate gradient methods for solving nonlinear monotone equations has been presented. The advantage of the method is that, the computation of Jacobian matrix is completely avoided throughout the iteration process. Also, a derivative-free methods for nonlinear monotone equations has been proposed in [26] and it has shown to converged Q-linearly to the solution of the monotone equations based on the assumption that the underlying function is Lipschitz continuous. Recently, some matrix-free methods have been proposed [19, 20, 21, 24, 21, 25].

Motivated by the above contributions, this paper aimed at proposing the derivative-free method for solving large-scale problem (1) that is globally convergent. The remaining part of the paper is organized as follows. In Section 2, we present the algorithms of the proposed method. Convergence analysis is presented in Section 3. Numerical results of the methods are reported in Section 4. Concluding remarks are given in Section 5.

2. Main Result

In this section, we present the proposed method for solving large scale system of nonlinear equations. The method is based on approximation of quasi-Newton's update in (4) via

$$B_k \approx \lambda_k I, \quad (6)$$

where $\lambda_k \in \mathbb{R}^n$ and I is an identity matrix.

In order to enhance good direction toward the solution, we suggest new direction d_k to be defined as

$$d_k = -\lambda_k^{-1} F_k, \quad (7)$$

where $\lambda_k \in \mathbb{R}$ is an acceleration parameter to be determined.

Furthermore, the search direction d_k is usually needed to satisfy the descent condition

$$\nabla f(x_k)^T d_k < 0.$$

Now, consider the Broyden's matrix updating formula given by

$$B_{k+1} = B_k + \frac{(y_k - B_k s_k) s_k^T}{s_k^T s_k}, \quad (8)$$

where $s_k = x_{k+1} - x_k$ and $y_k = F_{k+1} - F_k$. Despite the attractive features of this method, it is not suitable for solving the large-scale problems due the matrix storage at each iteration. Motivated by this reason, this work is aim at proposing a new matrix-free method for solving large-scale problems.

Now, from (6) and (8), it can be deduced that

$$\lambda_{k+1} I = \lambda_k I + \frac{(y_k - \lambda_k s_k) s_k^T}{s_k^T s_k}, \quad (9)$$

and by multiplying (8) by F_k , we have

$$\lambda_{k+1} F_k = \lambda_k F_k + \frac{(y_k - \lambda_k s_k) s_k^T F_k}{s_k^T s_k}. \quad (10)$$

Again, multiplying (10) by s_k^T , we have

$$\lambda_{k+1} s_k^T F_k = \lambda_k s_k^T F_k + \frac{s_k^T (y_k - \lambda_k s_k) s_k^T F_k}{s_k^T s_k}, \quad (11)$$

Dividing (11) by $s_k^T F_k$, where $s_k^T F_k \neq 0$ yields

$$\lambda_{k+1} = \lambda_k + \frac{s_k^T (y_k - \lambda_k s_k)}{s_k^T s_k}. \quad (12)$$

We finally present our iterative scheme as

$$x_{k+1} = x_k + \alpha_k d_k, \quad (13)$$

where $\alpha_k > 0$ is the step length and d_k is the search direction. Moreover, inexact line search proposed in [9] is used in this work to compute the step length α_k as follows.

Given some positive constants $\eta_1, \eta_2 > 0$ and let $h \in (0, 1)$. Suppose that $\{\omega_k\}$ is a sequence of some positive numbers for which

$$\sum_{k=0}^{\infty} \omega_k < \omega < \infty, \quad (14)$$

and

$$f(x_k + \alpha d_k) - f(x_k) \leq -\eta_1 \|\alpha F(x_k)\|^2 - \eta_2 \|\alpha d_k\|^2 + \omega_k f(x_k), \quad (15)$$

where $\alpha = h^i$ with i being the least nonnegative integer for which (15) holds. Set $\alpha_k = \alpha$.

Algorithm 1: On Efficient Matrix-Free Method Via Quasi-Newton Approach (EMQN)

Input: Given x_0 , $\lambda_0 = 0.01$, $\epsilon = 10^{-4}$, set $k = 0$.

Step 1: Compute $F(x_k)$.

Step 2: If $\|F_k\| \leq \epsilon$ then stop, else go to **Step 3**.

Step 3: Compute $d_k = -\lambda_k^{-1} F(x_k)$.

Step 4: Compute step length α_k (15).

Step 5: Set $x_{k+1} = x_k + \alpha_k d_k$.

Step 6: Compute F_{k+1} .

Step 7: Determine $\lambda_{k+1} = \lambda_k + \frac{s_k^T (y_k - \lambda_k s_k)}{s_k^T s_k}$.

Step 9: Set $k = k + 1$, and go to **Step 2**.

Remark 2.1. It can be seen that the parameter λ_{k+1} defined by (12) is a scalar for all k . In addition, the gradient of F is not needed in the implementation of Algorithm 1. With these into consideration, we can conclude that the Algorithm 1 is derivative-free as well as matrix-free. Therefore, Algorithm 1 is suitable for large-scale problems as well as nonsmooth problems. Furthermore, we show in Lemma 3.4 that the search direction generated by Algorithm 1 is sufficiently descent.

3. Convergence Result

In this section, we present the global convergence of our method (EMQN). To start, let the level set be defined as

$$\Omega = \{x \mid \|F(x)\| \leq \|F(x_0)\|\}. \quad (16)$$

Assumption 3.1. We now state the following assumptions to establish the convergence result of EMQN Algorithm .

(1) There exists a point $x^* \in \mathbb{R}^n$ such that $F(x^*) = 0$.

(2) F is continuously differentiable in some neighborhood say A of x^* containing Ω .

(3) The Jacobian of function F is positive definite bounded on A , namely, there exists some positive constants $G > g > 0$ such that

$$\|F'(x)\| \leq G, \quad \forall x \in A, \quad (17)$$

and

$$g\|d\|^2 \leq d^T F'(x)d, \quad \forall x \in A, d \in \mathbb{R}^n. \quad (18)$$

Remark 3.2. Assumption (3.1) implies that there exists a constants $G > g > 0$ such that

$$g\|d\| \leq \|F'(x)d\| \leq G\|d\|, \quad \forall x \in A, d \in \mathbb{R}^n. \quad (19)$$

$$g\|x - y\| \leq \|F(x) - F(y)\| \leq G\|x - y\|, \quad \forall x, y \in A. \quad (20)$$

Since $\lambda_k I$ approximates F'_k along direction d_k , let us state the following assumption.

Assumption 3.3. $\lambda_k I$ is a good approximation to $F'(x_k)$, i.e.,

$$\|(F'(x_k) - \lambda_k I)d_k\| \leq \epsilon \|F(x_k)\|, \quad (21)$$

where $\epsilon \in (0, 1)$ [13].

Lemma 3.4. Suppose that Assumption (3.3) holds and let $\{x_k\}$ be generated by EMQN algorithm. Then d_k is a descent direction of f at x_k i.e

$$\nabla f(x_k)^T d_k < 0. \quad (22)$$

Proof. From (7), we have

$$\begin{aligned} \nabla f(x_k)^T d_k &= F_k^T F'_k d_k \\ &= F_k^T [(F'_k - \lambda_k I) d_k - F_k] \\ &= F_k^T (F'_k - \lambda_k I) d_k - \|F_k\|^2, \end{aligned} \quad (23)$$

by Cauchy-Schwarz we have,

$$\begin{aligned} \nabla f(x_k)^T d_k &\leq \|F_k\| \|(F'_k - \lambda_k I) d_k - F_k\| \\ &\leq -(1 - \epsilon) \|F_k\|^2. \end{aligned} \quad (24)$$

Hence for $\epsilon \in (0, 1)$ we have (22).

Since the search direction satisfied the decent condition in (22), it means that the inequality $\|F_{k+1}\| \leq \|F_k\|$ holds. $\square$

Lemma 3.5. Suppose that Assumption (3.3) holds and $\{x_k\}$ be generated by EMQN algorithm. Then $\{x_k\} \subset \Omega$.

Proof. From lemma (3.4) we have $\|F_{k+1}\| \leq \|F_k\|$. In addition, for all k we have

$$\|F_{k+1}\| \leq \|F_k\| \leq \|F_{k-1}\| \leq \dots \leq \|F_0\|.$$

This shows that $\{x_k\} \subset \Omega$. $\square$

Lemma 3.6. (see[3]) Suppose that Assumption (3.1) holds and $\{x_k\}$ be generated by EMQN algorithm. Then there exists a constant $g > 0$ such that for all k

$$y_k^T s_k \geq g \|s_k\|^2. \quad (25)$$

Lemma 3.7. Suppose that Assumption (3.1) holds and $\{x_k\}$ is generated by EMQN algorithm. Then we have

$$\lim_{k \rightarrow \infty} \|\alpha_k d_k\| = \lim_{k \rightarrow \infty} \|s_k\| = 0, \quad (26)$$

and

$$\lim_{k \rightarrow \infty} \|\alpha_k F_k\| = 0. \quad (27)$$

Proof. By (15), we have for all $k > 0$,

$$\begin{aligned} \eta_2 \|\alpha_k d_k\|^2 &\leq \eta_1 \|\alpha_k F_k\|^2 + \eta_2 \|\alpha_k d_k\|^2 \\ &\leq \|F_k\|^2 - \|F_{k+1}\|^2 + \omega_k \|F_k\|^2. \end{aligned} \quad (28)$$

By summing the inequality above, we have

$$\begin{aligned} \eta_2 \sum_{i=0}^k \|\alpha_i d_i\|^2 &\leq \sum_{i=0}^k (\|F_i\|^2 - \|F_{i+1}\|^2) + \sum_{i=0}^k \omega_i \|F_i\|^2 \\ &= \|F_0\|^2 - \|F_{k+1}\|^2 + \sum_{i=0}^k \omega_i \|F_i\|^2 \\ &\leq \|F_0\|^2 + \|F_0\|^2 \sum_{i=0}^k \omega_i \\ &\leq \|F_0\|^2 + \|F_0\|^2 \sum_{i=0}^{\infty} \omega_i. \end{aligned} \quad (29)$$

So from the level set and fact that $\{\omega_k\}$ satisfies (14) then the series $\sum_{i=0}^{\infty} \|\alpha_i d_i\|^2$ is convergent. This implies (26). Following the similar arguments as above but with $\eta_1 \|\alpha_k F_k\|^2$ on the left hand side, we obtain (27). $\square$

Lemma 3.8. Suppose that Assumption (3.1) holds and let $\{x_k\}$ be generated by algorithm 1. Then there exists a constant $m_3 > 0$ such that for all $k > 0$,

$$\|d_k\| \leq m_3. \quad (30)$$

Proof. From (7) and (25) we have,

$$\begin{aligned} \|d_k\| &= \left\| -\lambda_k^{-1} F_k \right\| \\ &= \left\| -\left(\lambda_{k-1} I + \frac{(y_{k-1} - \lambda_{k-1} s_{k-1}) s_{k-1}^T}{s_{k-1}^T s_{k-1}} \right)^{-1} F_k \right\| \\ &= \left\| -\left(\lambda_{k-1} + \frac{y_{k-1}^T s_{k-1}}{s_{k-1}^T s_{k-1}} - \lambda_k \frac{s_{k-1}^T s_{k-1}}{s_{k-1}^T s_{k-1}} \right)^{-1} F_k \right\| \\ &= \left\| -\frac{\|s_{k-1}\|^2}{y_{k-1}^T s_{k-1}} F_k \right\| \\ &\leq \frac{\|s_{k-1}\|^2 \|F_k\|}{g \|s_{k-1}\|^2} \\ &\leq \frac{\|F_0\|}{g}. \end{aligned} \quad (31)$$

Taking $m_3 = \frac{\|F_0\|}{g}$, we have (30). $\square$

Theorem 3.9. Suppose that Assumption (3.1) holds and $\{x_k\}$ is generated by EMQN Algorithm. We further assume that for all $k > 0$,

$$\alpha_k \geq h \frac{|F_k^T d_k|}{\|d_k\|^2}, \quad (32)$$

where $h > 0$. Then

$$\lim_{k \rightarrow \infty} \|F_k\| = 0. \quad (33)$$

Proof. From Lemma (3.8), we have (30). Therefore by (26) and the boundedness of $\{\|d_k\|\}$, we have

$$\lim_{k \rightarrow \infty} \alpha_k \|d_k\|^2 = 0. \quad (34)$$

From (32) and (34), we have

$$\lim_{k \rightarrow \infty} |F_k^T d_k| = 0. \quad (35)$$

On the other hand from (7), we have

$$F_k^T d_k = -\lambda_k^{-1} \|F_k\|^2, \quad (36)$$

$$\begin{aligned} \|F_k\|^2 &= |-F_k^T d_k \lambda_k| \\ &\leq |F_k^T d_k| |\lambda_k|. \end{aligned} \quad (37)$$

By using (20), we obtain

$$\lambda_k = \lambda_{k-1} + \frac{y_{k-1}^T s_{k-1}}{\|s_{k-1}\|^2} - \frac{\lambda_{k-1} \|s_{k-1}\|^2}{\|s_{k-1}\|^2} = \frac{y_{k-1}^T s_{k-1}}{\|s_{k-1}\|^2} \leq \frac{\|y_{k-1}\| \|s_{k-1}\|}{\|s_{k-1}\|^2} \leq G,$$

which means, $|\lambda_k| \leq G$. So from (37), we have

$$\|F_k\|^2 \leq |F_k^T d_k| G. \quad (38)$$

Thus,

$$0 \leq \|F_k\|^2 \leq |F_k^T d_k| G \rightarrow 0. \quad (39)$$

Therefore,

$$\lim_{k \rightarrow \infty} \|F_k\| = 0. \quad (40)$$

The proof is completed. $\square$

4. Numerical results

In this section, some numerical results are presented to demonstrate the efficiency of the proposed method by comparing it with the following existing methods in the literature.

- An improved derivative-free method via double direction approach for solving systems of nonlinear equations (**IDFDD**) [3].
- Classical Broyden's method (**CBM**) for solving system of nonlinear equations.

The three algorithms were implemented using the same line search (15) in the course of the experiments and the following parameters are set: $\eta_1 = \eta_2 = 10^{-4}$, $h = 0.35$ and $\omega_k = \frac{1}{(k+1)^2}$. However, for the classical Broyden's method, we set $B_0 = I$, I is an identity matrix.

The computer codes used were written in Matlab 8.3.0.532 (R2014a) and run on a personal computer equipped with a 1.40.00 GHz CPU processor and 4 GB RAM memory. We have tried the three methods on three test problems with different initial points and dimension (n -values) between 100 to 10,000. The iteration is set to stop for all the methods if $\|F_k\| \leq 10^{-4}$. The symbol '-' represents failure due to:

- Failure to complete execution due to insufficient memory.
- Number of iterations exceed 1000 but no x_k satisfy the stopping criterion.

Table 1: Initial points

INITIAL GUESS (IP)	VALUES
x_1	$(\frac{1}{2}, \frac{1}{2}, \dots, \frac{1}{2})^T$
x_2	$(-1.5, -1.5, \dots, -1.5)^T$
x_3	$(-25, -25, \dots, -25)^T$
x_4	$(5, 5, \dots, 5)^T$
x_5	$(14, 14, \dots, 14)^T$

Problem 1: [4].

$$F_i(x) = 2x_i - \sin |x_i|, \quad i = 1, 2, \dots, n.$$

Problem 2: [10].

$$F_i(x) = \cos(x_i^2 - 1)^2 - 1, \quad i = 1, 2, \dots, n.$$

Problem 3.

$$F_1(x) = \frac{1}{3}x_1^3 + \frac{1}{2}x_2^2$$

$$F_i(x) = -\frac{1}{2}x_i^2 + \frac{i}{3}x_i^3 + \frac{1}{2}x_{i+1}^2, \quad i = 1, 2, \dots, n-1,$$

$$F_n(x) = -\frac{1}{2}x_n^2 + \frac{n}{3}x_n^3.$$

The results of the numerical experiments for the IDFDD and CBM methods as well as our proposed method

Table 2: Numerical results of EMQN, CMB and IDFDD methods for problem 1

Dimension	Initial Guess	EMQN		CBM		IDFDD	
		NIT	CPUT	NIT	CPUT	NIT	CPUT
100	x_1	28	0.142571	19	0.535173	23	0.143207
	x_2	24	0.028437	8	0.099478	25	0.112461
	x_3	9	0.055164	37	0.495608	-	-
	x_4	147	0.366485	78	0.931389	366	1.349186
	x_5	9	0.044106	109	1.283275	-	-
1000	x_1	31	0.107026	18	27.2388	26	0.246356
	x_2	25	0.119676	8	11.92841	28	0.344145
	x_3	9	0.086169	42	91.47828	-	-
	x_4	126	0.812398	140	242.139	397	3.385763
	x_5	9	0.070279	27	41.0478	-	-
10000	x_1	34	0.654757	-	-	27	1.842495
	x_2	29	0.619744	-	-	31	1.519259
	x_3	9	0.383985	-	-	-	-
	x_4	146	3.49417	-	-	428	23.4393
	x_5	9	0.384739	-	-	-	-

Table 3: Numerical results of EMQN, CMB and IDFDD methods for problem 2

Dimension	Initial Guess	EMQN		CBM		IDFDD	
		NIT	CPUT	NIT	CPUT	NIT	CPUT
100	x_1	9	0.054347	10	0.592524	4	0.091207
	x_2	6	0.026207	6	0.110625	-	-
	x_3	10	0.01284	8	0.183785	4	0.016239
	x_4	7	0.009098	10	0.196762	-	-
	x_5	8	0.010415	10	0.44328	-	-
1000	x_1	11	0.041969	11	19.17501	-	-
	x_2	7	0.022809	7	12.06882	-	-
	x_3	11	0.024054	9	15.57875	4	0.095884
	x_4	8	0.032852	11	19.02109	-	-
	x_5	9	0.051642	11	19.28136	-	-
10000	x_1	12	0.177906	-	-	-	-
	x_2	8	0.271242	-	-	-	-
	x_3	12	0.269865	-	-	4	0.566027
	x_4	10	0.171702	-	-	-	-
	x_5	13	0.305941	-	-	-	-

are reported in Tables 2-4, where NIT and CPUT are respectively stand for the number of iterations and the number of time taken for each method to successfully obtained the solution of each problem. Tables 3-4 indicated that the proposed method EMQN has minimum number of iterations and CPU time, compared to CBM and IDFDD methods, except at Table 2 with initial guesses x_1 and x_4 for the dimension 100 and x_1 and x_2 in 1000 dimension, where the number of iteration of CBM method is less than that of EMQN and IDFDD methods. Therefore, EMQN method out performed CBM and IDFDD methods. One can easily observe that our claim is fully justified from the Tables, that is, the proposed method has less CPU time and

Table 4: Numerical results of EMQN, CMB and IDFDD methods for Problem 3

Dimension	Initial Guess	EMQN		CBM		IDFDD	
		NIT	CPUT	NIT	CPUT	NIT	CPUT
100	x_1	12	0.239779	21	0.700504	27	1.230928
	x_2	11	0.226276	19	0.513159	29	1.29121
	x_3	13	0.270103	21	0.656313	23	1.156257
	x_4	14	0.244887	24	0.687868	28	0.968806
	x_5	14	0.226145	25	0.764494	26	1.047147
1000	x_1	14	0.944261	23	34.40515	35	5.13596
	x_2	11	0.892505	28	76.17269	29	5.233836
	x_3	13	0.951678	21	46.46457	31	5.45977
	x_4	14	0.947066	24	54.18002	32	6.261874
	x_5	14	1.036368	23	48.51827	40	7.275419
10000	x_1	14	88.4897	-	-	35	647.7165
	x_2	12	102.4096	-	-	29	526.0935
	x_3	15	99.16974	-	-	31	524.7104
	x_4	14	75.62304	-	-	32	480.4203




Figure 1: Performance profile of EMQN,CBM and IDFDD methods with respect to the number of iteration for the problems 1-3.

number of iterations for each of the test problems with the exception of Problem 1. Furthermore, on average, the CPU time of the proposed method is the smallest which signifies that our method is fully derivative-free and matrix-free (i.e., no computation of matrix at all).

Figures 1-2 show the summary of the numerical performance of the IDFDD and CBM methods against the proposed method in terms of iterations number and CPU time. The summary is evaluated based on the famous performance profiles developed by Dolan and Moré [4]. This means, for each method, the fraction $P(\tau)$ of the problems for which the method falls within a factor τ of the best time is plotted. The curve that stays longer on the vertical axis corresponds to the method that solved highest percentage of the test problems considered in a time that was within a factor τ of the best time.




Figure 2: Performance profile of EMQN,CBM and IDFDD methods with respect to the CPU time (in second) for the problems 1-3.

5. Conclusion

In this paper, an efficient matrix-free method via quasi-Newton update for handling nonlinear system of equations has been developed. This was achieved by approximating the Broyden's Update via acceleration parameter. The proposed method is completely matrix-free iterative method that is globally convergent under certain appropriate conditions. The efficiency as well as the performance of the proposed method have been compared with that of classical broyden method (CBM) and IDFDD method [3]. Numerical comparisons have been done using a set of large-scale test problems. Moreover, Table 2-4 and Figure 1-2, showed that the proposed method is quite efficient because it has the least number of iteration compared to IDFDD and CBM methods. Future research include using the proposed method to solve nonlinear problems as discussed in [35, 36, 40].

Acknowledgment

Nuttapol Pakkaranang would like to thank Phetchabun Rajabhat University.

References

- [1] R.P. Agarwal, B. Xu and W. Zhang, Stability of functional equations in single variable, *J. Math. Anal. Appl.*, 288 (2003), 852–869.
- [2] J.A. Baker, The stability of certain functional equations, *Proc. Amer. Math. Soc.*, 112 (1991), 729–732.
- [3] A.S. Halilu and M.Y. Waziri, An improved derivative-free method via double direction approach for solving systems of nonlinear equations, *J. of the Ramanujan Math. Soc.*, 33 (2018), 75–89.
- [4] J.E. Dennis and J.J. More, A characterization of superlinear convergence and its application to quasi-Newton methods, *Math. Comp.*, 28 (1974), 549–560.
- [5] M. Mamat, K. Muhammad and M.Y. Waziri, Trapezoidal Broyden's method for solving systems of nonlinear equations, *Appl. Math. Sci.*, 6 (2014), 251–260.
- [6] C.G. Broyden, A class of methods for solving nonlinear simultaneous equations, *Math. Comput.*, 19 (1965), 577–593.
- [7] M.Y. Waziri, Y.M. Kufena, and A.S. Halilu, Derivative-free three-term spectral conjugate gradient method for symmetric nonlinear equations, *Thai J. Math.*, 18 (2020), 1417–1431.
- [8] M. Ziani and H.F. Guyomarch, An autoadaptive limited memory Broyden's method to solve systems of nonlinear equations, *Appl. Math. Comput.*, 205 (2008), 205–215.
- [9] D. Li and M. Fukushima, A global and superlinear convergent Gauss-Newton based BFGS method for symmetric nonlinear equation, *SIAM J. Numer. Anal.*, 37 (1999), 152–172.
- [10] M.Y. Waziri and L.Y. Uba, Three-step derivative-free diagonal updating method for solving large-scale systems of nonlinear equations, *J. Numer. Math. Stoch.*, 6 (2014), 73–83.
- [11] W. Leong, M.A. Hassan and M.Y. Waziri, A matrix-free quasi-Newton method for solving large-scale nonlinear systems, *Comput. Math. Appl.*, 62 (2011), 2354–2363.
- [12] A. Ramli, M.L. Abdullah and M. Mamat, Broyden's method for solving fuzzy nonlinear equations, *Adv. Fuzzy Syst.*, (2010), Art. ID 763270, 6 pages.
- [13] A.S. Halilu and M.Y. Waziri, A transformed double step length method for solving large-scale systems of nonlinear equations, *J. Numer. Math. Stoch.*, 9 (2017), 20–32.
- [14] C.T. Kelley, *Solving nonlinear equations with Newton's method*, SIAM, Philadelphia (2003).
- [15] J.E. Dennis and R.B. Schnabel, *Numerical methods for unconstrained optimization and nonlinear equations*, SIAM, Philadelphia (1993).
- [16] W. Sun and Y.X. Yuan, *Optimization theory and methods; Nonlinear programming*, Springer, New York (2006).
- [17] A.S. Halilu and M.Y. Waziri, An enhanced matrix-free method via double step length approach for solving systems of nonlinear equations, *International Journal of Applied Mathematical Research*, 6(4) (2017), 147–156.
- [18] A.S. Halilu, A. Majumder, M.Y. Waziri, H. Abdullahi, Double direction and step length method for solving system of nonlinear equations, *Euro. J. Mol. Clinic. Med.*, 7(7) (2020), 3899–3913.
- [19] S. Aji, P. Kumam, A.M. Awwal, M.M. Yahaya and K. Sitthithakerngkiet, An efficient DY-type spectral conjugate gradient method for system of nonlinear monotone equations with application in signal recovery, *AIMS Math.*, 6 (2021): 8078–8106.
- [20] A.M. Awwal, P. Kumam and H. Mohammad, Iterative algorithm with structured diagonal Hessian approximation for solving nonlinear least squares problems, *J. Nonlinear Convex Anal.*, 22(6) (2021), 1173–1188.
- [21] A.M. Awwal, P. Kumam, K. Sitthithakerngkiet, A.M. Bakoji, A.S. Halilu and I.M. Sulaiman, Derivative-free method based on DFP updating formula for solving convex constrained nonlinear monotone equations and application, *AIMS Math.*, 6(8) (2021), 8792–8814.
- [22] S. Aji, P. Kumam, A.M. Awwal, M.M. Yahaya and W. Kumam, Two hybrid spectral methods with inertial effect for solving system of nonlinear monotone equations with application in robotics, *IEEE Access*, 9 (2021), 30918–30928.
- [23] A.S. Halilu and M.Y. Waziri, Solving systems of nonlinear equations using improved double direction method, *J. Nigerian Math. Soc.*, 39(2) (2020), 287–301.

- [24] A.M. Awwal, P. Kumam, L. Wang, S. Huang and W. Kumam, Inertial-based derivative-free method for system of monotone nonlinear equations and application, *IEEE Access*, 8 (2020) 226921–226930.
- [25] A.M. Awwal, L. Wang, P. Kumam, H. Mohammad and W. Wathayu, A projection Hestenes–Stiefel method with spectral parameter for nonlinear monotone equations and signal processing, *Math. Comput. Appl.*, 25(2) (2020), Art. ID 27–28, 29 pages.
- [26] Y.B. Zhao and D. Li, Monotonicity of fixed point and normal mapping associated with variational inequality and its application, *SIAM J. Optim.*, 11 (2001), 962–997.
- [27] M. Li, H. Liu and Z. Liu, A new family of conjugate gradient methods for unconstrained optimization, *J. Appl. Math. Comput.*, 58 (2018) 219–234.
- [28] A.S. Halilu, and M.Y. Waziri, Inexact double step length method for solving systems of nonlinear equations, *Stat. Optim. Inf. Comput.*, 8(1) (2020), 165–174.
- [29] H. Abdullahi, A.S. Halilu, and M.Y. Waziri, A modified conjugate gradient method via a double direction approach for solving large-scale symmetric nonlinear systems, *Journal of Numerical Mathematics and Stochastics*, 10(1) (2018), 32–44.
- [30] Y.H. Dai and C.X. Kou, A nonlinear conjugate gradient algorithm with an optimal property and an improved wolfe line search, *SIAM J. Optim.*, 23 (2013), 296–320.
- [31] Y.H. Dai and L.Z. Lio, New conjugacy conditions and related nonlinear conjugate gradient methods, *Appl. Math. Optim.*, 43 (2001), 87–101.
- [32] W.W. Hager and H. Zhang, A survey of nonlinear conjugate gradient methods, *Pac. J. Optim.*, 2 (2006), 35–58.
- [33] A.S. Halilu, M.Y. Waziri and I. Yusuf, Efficient matrix-free direction method with line search for solving large-scale system of nonlinear equations, *Yugosl. J. Oper. Res.*, 30(4) (2020), 399–412.
- [34] M.Y. Waziri, K. Ahmad, and A.S. Halilu, Enhanced Dai–Liao conjugate gradient methods for systems of monotone nonlinear equations, *SeMA J.*, 78(1) (2020), 15–51.
- [35] L. Li and D. Wu, The convergence of Ishikawa iteration for generalized Φ -contractive mappings, *Results in Nonlinear Analysis*, 4(1) (2021), 47–56.
- [36] P. Lo'lo' and M. Shabibi, Common best proximity points theorems for H-contractive non-self mappings, *Advances in the Theory of Nonlinear Analysis and its Application*, 5(2) (2021), 173–179.
- [37] K. Meintjes, and A.P. Morgan, A methodology for solving chemical equilibrium systems, *Appl. Math. Comput.*, 22 (1987), 333–361.
- [38] M. Sun, M.Y. Tian, and Y.J. Wang, Multi-step discrete-time Zhang neural networks with application to time-varying nonlinear optimization, *Discrete Dyn. Nat. Soc.*, Art. ID 4745759, (2019) 1–14.
- [39] A.M. Awwal, L. Wang, P. Kumam, and H. Mohammad, A two-step spectral gradient projection method for system of nonlinear monotone equations and image deblurring problems, *Symmetry*, 12(6) (2020), Art. ID 874, 20 pages.
- [40] N. Wairojjana, N. Pakkaranang, I. Uddin, P. Kumam and A.M. Awwal, Modified proximal point algorithms involving convex combination technique for solving minimization problems with convergence analysis, *Optimization*, 69(7–8) (2020), 1655–1680.
- [41] J.E. Dennis and R.B Schnabel, Numerical method for unconstrained optimization and non-linear equations, practice Hall, Englewood cliffs, NJ, USA, 1983.
- [42] D.W. Marquardt, An algorithm for least-squares estimation of nonlinear parameters, *J. Soc. Indust. Appl. Math.*, 11 (1963), 431–441.
- [43] Y.B. Musa, M.Y. Waziri, and A.S. Halilu, On computing the regularization Parameter for the Levenberg–Marquardt method via the spectral radius approach to solving systems of nonlinear equations, *J. Numer. Math. Stoch.*, 9(1) (2017), 80–94.
- [44] A.S. Halilu, M.K. Dauda, M.Y. Waziri, and M. Mamat, A derivative-free decent method via acceleration parameter for solving systems of nonlinear equations, *Open J. sci. tech.*, 2(3) (2019) 1–4.
- [45] A.S. Halilu, A. Majumder, M.Y. Waziri, K. Ahmed, Signal recovery with convex constrained nonlinear monotone equations through conjugate gradient hybrid approach, *Math. Comput. Simulation*, 187 (2021), 520–539.
- [46] M.Y. Waziri, H.U. Muhammad, A.S. Halilu, and K. Ahmad, Modified matrix-free methods for solving system of nonlinear equations, (2020) DOI: 10.1080/02331934.2020.1778689.