

Source details

Nonlinear Functional Analysis and Applications

Scopus coverage years: from 2017 to 2021

Publisher: Kyungnam University Press

ISSN: 1229-1595 E-ISSN: 2466-0973

Subject area:

Mathematics: Analysis

Mathematics: Numerical Analysis

Mathematics: Applied Mathematics

Mathematics: Control and Optimization

Source type: Journal

CiteScore 2020 1.1 ⓘ

SJR 2020 0.353 ⓘ

SNIP 2020 0.803 ⓘ

View all documents >

Set document alert

 Save to source list

Source Homepage

CiteScore

CiteScore rank & trend

Scopus content coverage

i Improved CiteScore methodology ⓘ

CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore 2020 ▾

1.1 = $\frac{244 \text{ Citations 2017 - 2020}}{230 \text{ Documents 2017 - 2020}}$

Calculated on 05 May, 2021

CiteScoreTracker 2021 ⓘ

1.2 = $\frac{276 \text{ Citations to date}}{224 \text{ Documents to date}}$

Last updated on 06 March, 2022 • Updated monthly

CiteScore rank 2020 ⓘ

Category	Rank	Percentile
Mathematics		
Analysis	#108/164	34th
Mathematics		
Numerical Analysis	#46/66	31st

View CiteScore methodology >

CiteScore FAQ >

Add CiteScore to your site 

About Scopus

- What is Scopus
- Content coverage
- Scopus blog
- Scopus API
- Privacy matters

Language

- 日本語に切り替える
- 切换到简体中文
- 切换到繁體中文
- Русский язык

Customer Service

- Help
- Tutorials
- Contact us

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © Elsevier B.V ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.
We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.



Nonlinear Functional Analysis and Applications

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
South Korea  Universities and research institutions in South Korea	Mathematics Analysis Applied Mathematics Control and Optimization Numerical Analysis	Kyungnam University Press	6
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	12291595, 24660973	2017-2020	Homepage How to publish in this journal jongkyuk@kyungnam.ac.kr

SCOPE

The international mathematical journal NFAA will publish carefully selected original research papers on nonlinear functional analysis and applications, that is, ordinary differential equations, all kinds of partial differential equations, functional differential equations, integrodifferential equations, control theory, approximation theory, optimal control, optimization theory, numerical analysis, variational inequality, asymptotic behavior, fixed point theory, dynamic systems and complementarity problems. Papers for publication will be communicated and recommended by the members of the Editorial Board.


 Join the conversation about this journal

 Quartiles



FIND SIMILAR JOURNALS


1 Thai Journal of Mathematics THA 81% similarity	2 Panamerican Mathematical Journal USA 74% similarity	3 Fixed Point Theory and Applications GBR 72% similarity	4 Carpathian Journal of Mathematics ROU 71% similarity	5 Journ and A CHE
---	--	---	---	--



SCImago Graphica

Explore, visually communicate and make sense of data with our new free tool.

Get it



Metrics based on Scopus® data as of April 2021

Gh. Mostafa 2 years ago

necessary conditions



Nonlinear Functional Analysis and Applications

Volume 27

Number 1

2022

Editor-in-Chief

Jong Kyu Kim, Kyungnam University, Korea

Print ISSN 1229-1595 / Online ISSN 2466-0973

Published Online: 7 March 2022



Nonlinear Functional Analysis and Applications (NFAA)

HOME	ABOUT	LOG IN	REGISTER	SEARCH	CURRENT	ARCHIVES
AIM AND SCOPE	EDITORIAL BOARD					

Home > Archives > **Volume 27, Number 1, March 2022**

Volume 27, Number 1, March 2022

Table of Contents

Articles

<u>A NEW EXPLICIT EXTRAGRADIENT METHOD FOR SOLVING EQUILIBRIUM PROBLEMS WITH CONVEX CONSTRAINTS</u>	PDF
<i>Kanikar Muangchoo</i>	1-22
<u>GENERAL BICONVEX FUNCTIONS AND BIVARIATIONAL-LIKE INEQUALITIES</u>	PDF
<i>Muhammad Aslam Noor</i>	23-44
<u>STRONG AND Δ-CONVERGENCE THEOREMS FOR A COUNTABLE FAMILY OF MULTI-VALUED DEMICONTRACTIVE MAPS IN HADAMARD SPACES</u>	PDF
<i>Ma'aruf Shehu Minjibir, Sani Salisu</i>	45-58
<u>IMPROVED GENERALIZED M-ITERATION FOR QUASI-NONEXPANSIVE MULTIVALUED MAPPINGS WITH APPLICATION IN REAL HILBERT SPACES</u>	PDF
<i>Francis Akutsah, Ojen Kumar Narain, Jong Kyu Kim</i>	59-82
<u>SETVALUED MIXED QUASI-EQUILIBRIUM PROBLEMS WITH OPERATOR SOLUTIONS</u>	PDF
<i>Tirth Ram, Anu Kumari Khanna, Ravdeep Kour</i>	83-97
<u>ON SUBCLASSES OF ANALYTIC FUNCTIONS ASSOCIATED WITH STRUVE FUNCTIONS</u>	PDF
<i>B.A. Frasin, Tariq Al-Hawary, Feras Yousef, I. Aldawish</i>	99-110
<u>DISKCYCLIC C_0-SEMICROUPS AND DISKCYCLICITY CRITERIA</u>	PDF
<i>Mansoorh Moosapoor</i>	111-119
<u>HALPERN TSENG'S EXTRAGRADIENT METHODS FOR SOLVING VARIATIONAL INEQUALITIES INVOLVING SEMISTRICTLY QUASIMONOTONE OPERATOR</u>	PDF
<i>Nopparat Wairojjana, Nuttapol Pakkaranang</i>	121-140
<u>SOME INEQUALITIES ON POLAR DERIVATIVE OF A POLYNOMIAL</u>	PDF
<i>Khangembam Babina Devi, Kshetrimayum Krishnadas, Barchand Chanam</i>	141-148
<u>ITERATIVE METHOD FOR SOLVING FINITE FAMILIES OF VARIATIONAL INEQUALITY AND FIXED POINT PROBLEMS OF CERTAIN MULTI-VALUED MAPPINGS</u>	PDF
<i>Musa Adewale Olona, Ojen Kumar Narain</i>	149-167
<u>A POWER SERIES ASSOCIATED WITH THE GENERALIZED HYPERGEOMETRIC FUNCTIONS WITH THE UNIT ARGUMENT WHICH ARE INVOLVED IN BELL POLYNOMIALS</u>	PDF
<i>Junesang Choi, Mohd Idris Qureshi, Javid Majid, Jahan Ara</i>	169-187
<u>NUMERICAL SOLUTIONS OF NONLINEAR VOLTERRA-FREDHOLM INTEGRO-DIFFERENTIAL EQUATIONS BY USING MADM AND VIM</u>	PDF
<i>Ayoob M. Abed, Muhammed F. Younis, Ahmed A. Hamoud</i>	189-201
<u>HYBRID INERTIAL CONTRACTION PROJECTION METHODS EXTENDED TO VARIATIONAL INEQUALITY PROBLEMS</u>	PDF
<i>N. D. Truong, J. K. Kim, T. H. H. Anh</i>	203-221

ISSN: 1229-1595 (Print), 2466-0973 (Online)

(51767) 7 Kyungnamdaehak-ro, Masanhappo-gu, Changwon-si, Gyeongsangnam-do, Republic of Korea

USER

Username

Password

☐ Remember me

NOTIFICATIONS

- [View](#)
- [Subscribe / Unsubscribe](#)

JOURNAL CONTENT

Search

All

Browse

- [By Issue](#)
- [By Author](#)
- [By Title](#)

FONT SIZE

INFORMATION

- [For Readers](#)
- [For Authors](#)
- [For Librarians](#)



HALPERN TSENG'S EXTRAGRADIENT METHODS FOR SOLVING VARIATIONAL INEQUALITIES INVOLVING SEMISTRICTLY QUASIMONOTONE OPERATOR

Nopparat Wairojjana¹ and Nuttapol Pakkaranang²

¹Applied Mathematics Program, Faculty of Science and Technology
Valaya Alongkorn Rajabhat University under the Royal Patronage (VRU)
1 Moo 20 Phaholyothin Road, Klong Neung
Klong Luang, Pathumthani, 13180, Thailand
e-mail: nopparat@vru.ac.th

²Mathematics and Computing Science Program, Faculty of Science and Technology
Phetchabun Rajabhat University, Phetchabun 67000, Thailand
e-mail: nuttapol.pak@prcu.ac.th

Abstract. In this paper, we study the strong convergence of new methods for solving classical variational inequalities problems involving semistrictly quasimonotone and Lipschitz-continuous operators in a real Hilbert space. Three proposed methods are based on Tseng's extragradient method and use a simple self-adaptive step size rule that is independent of the Lipschitz constant. The step size rule is built around two techniques: the monotone and the non-monotone step size rule. We establish strong convergence theorems for the proposed methods that do not require any additional projections or knowledge of an involved operator's Lipschitz constant. Finally, we present some numerical experiments that demonstrate the efficiency and advantages of the proposed methods.

1. INTRODUCTION

Let $\mathcal{H}$ be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and the induced norm $\| \cdot \|$. The strong converge of the sequence $\{x_n\}$ to an element x is denoted by

⁰Received May 27, 2021. Revised September 23, 2021. Accepted November 6, 2021.

⁰2020 Mathematics Subject Classification: 47J20, 90C25, 90C30, 90C52, 65Y05.

⁰Keywords: Variational Inequalities, Halpern Tseng's extragradient method, strong convergence theorems, semistrictly quasimonotone operator, Lipschitz continuous.

⁰Corresponding author: Nuttapol Pakkaranang(nuttapol.pak@prcu.ac.th).

$x_n \rightarrow x$. For a given closed and convex subset $\mathcal{C} \subset \mathcal{H}$, the variational inequality problem denoted by $VI(\mathcal{C}, \mathcal{G})$ is to find $x^* \in \mathcal{C}$ such that

$$\langle \mathcal{G}(x^*), y - x^* \rangle \geq 0, \quad \forall y \in \mathcal{C}, \quad (\text{VIP})$$

where $\mathcal{G} : \mathcal{H} \rightarrow \mathcal{H}$ is an operator. It is well known (see for details [7]) that the original problem is closely related to find a point $x^* \in \mathcal{C}$ such that

$$\langle \mathcal{G}(y), y - x^* \rangle \geq 0, \quad \forall y \in \mathcal{C}. \quad (\text{DVIP})$$

We will refer to problem (DVIP) as the dual variational inequality problem $DVI(\mathcal{C}, \mathcal{G})$ of (VIP) based on the paper [7]. For a closed and convex $\mathcal{C} \subset \mathcal{H}$, the metric projection $P_{\mathcal{C}} : \mathcal{H} \rightarrow \mathcal{C}$ is described for all $x \in \mathcal{H}$ such that

$$P_{\mathcal{C}}(x) = \arg \min \{\|x - y\| : y \in \mathcal{C}\}.$$

Moreover, $\mathbb{R}, \mathbb{N}$ are the set of real numbers and natural numbers, respectively. It is useful to note that the problem (VIP) is equivalent to solve the following problem:

$$\text{Find } x^* \in \mathcal{C} \text{ such that } x^* = P_{\mathcal{C}}[x^* - \lambda \mathcal{G}(x^*)],$$

where λ is any positive real number.

The theory of variational inequalities has been used extensively in the investigation of a collection of topics, consisting of physics, engineering, economics, and optimization theory. This problem was first introduced by Stampacchia [33] in 1964, and it is also well established that the problem (VIP) is a pivotal problem in nonlinear analysis. It is a significant mathematical problem that incorporates several important topics of applied mathematics, such as network equilibrium problems, the necessary optimality conditions, complementarity problems, and systems of nonlinear equations (for more details [8, 13, 14, 15, 16, 17, 28, 29, 30, 32]). On the other hand, the projection methods are effective iterative methods to solve variational inequalities. Many iterative methods for solving variational inequalities have been designed and investigated previously (see for more details [1, 4, 5, 6, 11, 12, 18, 20, 21, 22, 23, 24, 25, 26, 27, 31, 34, 35, 37]).

Then, Korpelevich [18] and Antipin [2] were the first to propose the extragradient method. The method is of the following design:

$$\begin{cases} x_0 \in \mathcal{C}, \\ y_n = P_{\mathcal{C}}[x_n - \lambda \mathcal{G}(x_n)], \\ x_{n+1} = P_{\mathcal{C}}[x_n - \lambda \mathcal{G}(y_n)], \end{cases} \quad (1.1)$$

where $0 < \lambda < \frac{1}{L}$ is the Lipschitz constant of an operator $\mathcal{G}$ and L is the Lipschitz constant of an operator $\mathcal{G}$. In view of the above method, we use two projections on the underlying set $\mathcal{C}$ over each iteration. This, of course, can affect the computational effectiveness of the used method if the feasible set

$\mathcal{C}$ has a complicated structure. Here, we restrict in our interest to presenting some methods which can address this drawback. The first is the following subgradient extragradient method due to Censor et al. [4]. This method takes the form of

$$\begin{cases} x_0 \in \mathcal{C}, \\ y_n = P_{\mathcal{C}}[x_n - \lambda \mathcal{G}(x_n)], \\ x_{n+1} = P_{\mathcal{H}_n}[x_n - \lambda \mathcal{G}(y_n)], \end{cases} \quad (1.2)$$

where $0 < \lambda < \frac{1}{L}$ and

$$\mathcal{H}_n = \{z \in \mathcal{H} : \langle x_n - \lambda \mathcal{G}(x_n) - y_n, z - y_n \rangle \leq 0\}.$$

In this article, we focus on the Tseng's extragradient method [34] that need to calculate only one projection for each iteration:

$$\begin{cases} x_0 \in \mathcal{C}, \\ y_n = P_{\mathcal{C}}[x_n - \lambda \mathcal{G}(x_n)], \\ x_{n+1} = y_n + \lambda [\mathcal{G}(x_n) - \mathcal{G}(y_n)], \end{cases} \quad (1.3)$$

where $0 < \lambda < \frac{1}{L}$. The above-mentioned methods have two significant disadvantages. The first is the constant step size, which necessarily involves the knowledge or estimation of the Lipschitz constant of the involved operator and only converges weakly in Hilbert spaces. In certain cases, the Lipschitz constants are unknown since they are difficult to compute for an operator. Estimating the Lipschitz constant a priori may be difficult from a computational point of view, which may affect the method's convergence rate and applicability.

The primary objective of this paper is to investigate semistrictly quasimonotone variational inequalities in infinite-dimensional Hilbert spaces. To prove that the iterative sequence generated by Tseng's extragradient method for the solution of semistrictly quasimonotone variational inequalities converges strongly to some solutions. The proposed methods are inspired by the projection method [34] and the Halpern method [10]. The proposed methods only require solving one projection onto the feasible set per iteration. If suitable conditions are imposed on control parameters, the iterative sequences generated by our methods strongly converge to some solution to the problem. We also provide examples to demonstrate the computational performance of the new methods.

The paper is organized in the following manner. In Sect. 2, some preliminary results were presented. Sect. 3 provides new algorithms and their convergence study. Finally, Sect. 4 presents some numerical results to point out the practical efficiency of the proposed methods.

2. PRELIMINARIES

Let $\mathcal{H}$ be a real Hilbert space. Given $x, y \in \mathcal{H}$, we interpret the closed line segment

$$[x, y] = \{tx + (1 - t)y : 0 \leq t \leq 1\}.$$

The segments $(x, y]$, $[x, y)$, and (x, y) are defined similarly. A *metric projection* $P_{\mathcal{C}}(x)$ of $x \in \mathcal{H}$ is defined by

$$P_{\mathcal{C}}(x) = \arg \min\{\|x - y\| : y \in \mathcal{C}\}.$$

For all $x, y \in \mathcal{H}$, we have

$$\|x + y\|^2 = \|x\|^2 + 2\langle x, y \rangle + \|y\|^2.$$

Lemma 2.1. *Assume $\mathcal{C}$ be a nonempty, closed and convex subset of a real Hilbert space $\mathcal{H}$ and $P_{\mathcal{C}} : \mathcal{H} \rightarrow \mathcal{C}$ be a metric projection from $\mathcal{H}$ onto $\mathcal{C}$.*

(i) *Let $x \in \mathcal{C}$ and $y \in \mathcal{H}$, we have*

$$\|x - P_{\mathcal{C}}(y)\|^2 + \|P_{\mathcal{C}}(y) - y\|^2 \leq \|x - y\|^2.$$

(ii) *$z = P_{\mathcal{C}}(x)$ if and only if*

$$\langle x - z, y - z \rangle \leq 0, \quad \forall y \in \mathcal{C}.$$

(iii) *For $y \in \mathcal{C}$ and $x \in \mathcal{H}$*

$$\|x - P_{\mathcal{C}}(x)\| \leq \|x - y\|.$$

Definition 2.2. Suppose that $\mathcal{C}$ is a nonempty, closed and convex subset of a real Hilbert space $\mathcal{H}$, and let $\mathcal{G} : \mathcal{C} \rightarrow \mathcal{H}$ be a mapping. The mapping $\mathcal{G}$ is said to be:

(a) *strongly monotone* on $\mathcal{C}$ with constant $\gamma > 0$ if for each pair of points $x, y \in \mathcal{C}$, we have

$$\langle \mathcal{G}(x) - \mathcal{G}(y), x - y \rangle \geq \gamma \|x - y\|^2;$$

(b) *strictly monotone* on $\mathcal{C}$ if for all distinct $x, y \in \mathcal{C}$, we have

$$\langle \mathcal{G}(x) - \mathcal{G}(y), x - y \rangle > 0;$$

(c) *monotone* on $\mathcal{C}$ if for all distinct $x, y \in \mathcal{C}$, we have

$$\langle \mathcal{G}(x) - \mathcal{G}(y), x - y \rangle \geq 0;$$

(d) *pseudomonotone* on $\mathcal{C}$ if for all distinct $x, y \in \mathcal{C}$, $\langle \mathcal{G}(y), x - y \rangle \geq 0$, then

$$\langle \mathcal{G}(x), x - y \rangle \geq 0;$$

(e) *quasimonotone* on $\mathcal{C}$ if for all distinct $x, y \in \mathcal{C}$, $\langle \mathcal{G}(y), x - y \rangle > 0$, then

$$\langle \mathcal{G}(x), x - y \rangle \geq 0;$$

- (f) *semistrictly quasimonotone* [9] on $\mathcal{C}$ if $\mathcal{G}$ is quasimonotone on $\mathcal{C}$ and for all distinct of points $x, y \in \mathcal{C}$, $\langle \mathcal{G}(y), x - y \rangle > 0$, then

$$\langle \mathcal{G}(z), x - y \rangle \geq 0, \text{ for some } z \in \left(\frac{x+y}{2}, x \right).$$

Remark 2.3. The implications are as follows:

$$(a) \implies (b) \implies (c) \implies (d) \implies (e) \quad \text{and} \quad (f) \implies (e).$$

However, in general, the inverse assertions are false.

Lemma 2.4. ([3]) *For any $x, y \in \mathcal{H}$ and $\ell \in \mathbb{R}$, we have*

- (i) $\|\ell x + (1 - \ell)y\|^2 = \ell\|x\|^2 + (1 - \ell)\|y\|^2 - \ell(1 - \ell)\|x - y\|^2$.
- (ii) $\|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y \rangle$.

Definition 2.5. A mapping $\mathcal{G} : \mathcal{H} \rightarrow \mathcal{H}$ is said to be

- (i) *weakly hemicontinuous* if $\mathcal{G}$ is upper semicontinuous from line segments in $\mathcal{H}$ to the weak topology of $\mathcal{H}$;
- (ii) *weakly sequentially continuous* if $\{\mathcal{G}(x_n)\}$ converges weakly to $\mathcal{G}(x)$ for every sequence $\{x_n\}$ converges weakly to x .

Remark 2.6. It is easy to prove that if $\mathcal{G} : \mathcal{H} \rightarrow \mathcal{H}$ is weakly sequentially continuous, then $\mathcal{G}$ must be weakly hemicontinuous.

It is known that the following conclusion is true.

Lemma 2.7. ([7]) *A solution of problem (DVIP) is always a solution of problem (VIP) provided that the operator $\mathcal{G}$ is, say, weakly hemicontinuous.*

Lemma 2.8. ([36]) *Let $\{p_n\} \subset [0, +\infty)$ be a sequence such that*

$$p_{n+1} \leq (1 - q_n)p_n + q_nr_n, \quad \forall n \in \mathbb{N}.$$

Moreover, two sequence $\{q_n\} \subset (0, 1)$ and $\{r_n\} \subset \mathbb{R}$ such that

$$\lim_{n \rightarrow +\infty} q_n = 0, \quad \sum_{n=1}^{+\infty} q_n = +\infty \quad \text{and} \quad \limsup_{n \rightarrow +\infty} r_n \leq 0.$$

Then, $\lim_{n \rightarrow +\infty} p_n = 0$.

Lemma 2.9. ([19]) *Let a real number sequence $\{p_n\}$ and there exists a subsequence $\{n_i\}$ of $\{n\}$ such that*

$$p_{n_i} < p_{n_{i+1}}, \quad \forall i \in \mathbb{N}.$$

Then, there exist a nondecreasing sequence $\{m_k\} \subset \mathbb{N}$ such that $m_k \rightarrow +\infty$ as $k \rightarrow +\infty$, and satisfying the following inequality for $k \in \mathbb{N}$:

$$p_{m_k} \leq p_{m_{k+1}} \quad \text{and} \quad p_k \leq p_{m_{k+1}}.$$

Indeed, $m_k = \max\{j \leq k : p_j \leq p_{j+1}\}$.

3. MAIN RESULTS

In this section, we introduce different iterative methods for variational inequalities involving semistrictly quasimonotone based on Tseng's extragradient method and the Halpern method that does not involve knowledge of the operator's Lipschitz constant or additional projection.

Algorithm 1.

Step 0: Choose $x_1 \in \mathcal{C}$, $0 < \lambda < \frac{1}{L}$, $\mu \in (0, 1)$ and $\{\alpha_n\} \subset (0, 1)$ satisfies the following conditions:

$$\lim_{n \rightarrow +\infty} \alpha_n = 0 \quad \text{and} \quad \sum_{n=1}^{+\infty} \alpha_n = +\infty.$$

Step 1: Compute

$$y_n = P_{\mathcal{C}}(x_n - \lambda \mathcal{G}(x_n)).$$

If $x_n = y_n$, then STOP and y_n is a solution. Otherwise, go to **Step 2**.

Step 2: Compute

$$z_n = y_n + \lambda [\mathcal{G}(x_n) - \mathcal{G}(y_n)].$$

Step 3: Compute

$$x_{n+1} = \alpha_n x_0 + (1 - \alpha_n) z_n.$$

Set $n = n + 1$ and go back to **Step 1**.

To prove the convergence analysis, it is assumed that the following conditions have been met:

- (G1) The solution set of problem (VIP) is denoted by Ω is nonempty;
- (G2) An operator $\mathcal{G} : \mathcal{H} \rightarrow \mathcal{H}$ is semistrictly quasimonotone, that is, if $\mathcal{G}$ is quasimonotone on $\mathcal{C}$ and $\langle \mathcal{G}(y), x - y \rangle > 0$, then

$$\langle \mathcal{G}(z), x - y \rangle \geq 0, \quad \text{for some } z \in \left(\frac{x + y}{2}, x \right); \quad (\text{SQM})$$

- (G3) An operator $\mathcal{G} : \mathcal{H} \rightarrow \mathcal{H}$ is *Lipschitz continuous* with constant $L > 0$, that is,

$$\|\mathcal{G}(x) - \mathcal{G}(y)\| \leq L\|x - y\|, \quad \forall x, y \in \mathcal{C}; \quad (\text{LC})$$

- (G4) An operator $\mathcal{G} : \mathcal{H} \rightarrow \mathcal{H}$ is *sequentially weakly continuous*, that is, if $\{\mathcal{G}(x_n)\}$ converges weakly to $\mathcal{G}(x)$ for every sequence $\{x_n\}$ converges weakly to x .

Lemma 3.1. *Suppose that $\mathcal{G} : \mathcal{H} \rightarrow \mathcal{H}$ satisfies the conditions (G1)-(G4) and sequence $\{x_n\}$ generated by Algorithm 1. Then, we have*

$$\|z_n - x^*\|^2 \leq \|x_n - x^*\|^2 - (1 - \lambda^2 L^2) \|x_n - y_n\|^2.$$

Proof. Since $x^* \in \Omega$, we have

$$\begin{aligned} & \|z_n - x^*\|^2 \\ &= \|y_n + \lambda[\mathcal{G}(x_n) - \mathcal{G}(y_n)] - x^*\|^2 \\ &= \|y_n - x^*\|^2 + \lambda^2 \|\mathcal{G}(x_n) - \mathcal{G}(y_n)\|^2 + 2\lambda \langle y_n - x^*, \mathcal{G}(x_n) - \mathcal{G}(y_n) \rangle \\ &= \|y_n + x_n - x_n - x^*\|^2 + \lambda^2 \|\mathcal{G}(x_n) - \mathcal{G}(y_n)\|^2 + 2\lambda \langle y_n - x^*, \mathcal{G}(x_n) - \mathcal{G}(y_n) \rangle \\ &= \|y_n - x_n\|^2 + \|x_n - x^*\|^2 + 2\langle y_n - x_n, x_n - x^* \rangle \\ &\quad + \lambda^2 \|\mathcal{G}(x_n) - \mathcal{G}(y_n)\|^2 + 2\lambda \langle y_n - x^*, \mathcal{G}(x_n) - \mathcal{G}(y_n) \rangle \\ &= \|x_n - x^*\|^2 + \|y_n - x_n\|^2 + 2\langle y_n - x_n, y_n - x^* \rangle + 2\langle y_n - x_n, x_n - y_n \rangle \\ &\quad + \lambda^2 \|\mathcal{G}(x_n) - \mathcal{G}(y_n)\|^2 + 2\lambda \langle y_n - x^*, \mathcal{G}(x_n) - \mathcal{G}(y_n) \rangle. \end{aligned} \quad (3.1)$$

It is given that

$$y_n = P_{\mathcal{C}}[x_n - \lambda \mathcal{G}(x_n)]$$

and it gives that

$$\langle x_n - \lambda \mathcal{G}(x_n) - y_n, y - y_n \rangle \leq 0, \quad \forall y \in \mathcal{C}. \quad (3.2)$$

Thus, we have

$$\langle x_n - y_n, x^* - y_n \rangle \leq \lambda \langle \mathcal{G}(x_n), x^* - y_n \rangle. \quad (3.3)$$

Combining expressions (3.1) and (3.3), we have

$$\begin{aligned} & \|z_n - x^*\|^2 \\ &\leq \|x_n - x^*\|^2 + \|y_n - x_n\|^2 + 2\lambda \langle \mathcal{G}(x_n), x^* - y_n \rangle - 2\langle x_n - y_n, x_n - y_n \rangle \\ &\quad + \lambda^2 \|\mathcal{G}(x_n) - \mathcal{G}(y_n)\|^2 - 2\lambda \langle \mathcal{G}(x_n) - \mathcal{G}(y_n), x^* - y_n \rangle \\ &= \|x_n - x^*\|^2 - \|x_n - y_n\|^2 + \lambda^2 \|\mathcal{G}(x_n) - \mathcal{G}(y_n)\|^2 - 2\lambda \langle \mathcal{G}(y_n), y_n - x^* \rangle. \end{aligned} \quad (3.4)$$

It is given that x^* is the solution of the problem (VIP) implies that

$$\langle \mathcal{G}(x^*), y - x^* \rangle \geq 0, \quad \forall y \in \mathcal{C}.$$

It implies that

$$\langle \mathcal{G}(y), y - x^* \rangle \geq 0, \quad \forall y \in \mathcal{C}.$$

Substituting $y = y_n \in \mathcal{C}$, we have

$$\langle \mathcal{G}(y_n), y_n - x^* \rangle \geq 0. \quad (3.5)$$

From expressions (3.4) and (3.5), we obtain

$$\begin{aligned} \|z_n - x^*\|^2 &\leq \|x_n - x^*\|^2 - \|x_n - y_n\|^2 + \lambda^2 L^2 \|x_n - y_n\|^2 \\ &= \|x_n - x^*\|^2 - (1 - \lambda^2 L^2) \|x_n - y_n\|^2. \end{aligned} \quad (3.6)$$

□

Lemma 3.2. *Let $\mathcal{G} : \mathcal{H} \rightarrow \mathcal{H}$ satisfying (G1)–(G4). If there exists a weakly convergent subsequence $\{x_{n_k}\}$ of $\{x_n\}$ to $\hat{x}$ and $\lim_{k \rightarrow +\infty} \|x_{n_k} - y_{n_k}\| = 0$. Then, $\hat{x}$ is the solution of (VIP).*

Proof. Indeed, we have

$$y_{n_k} = P_{\mathcal{C}}[x_{n_k} - \lambda_{n_k} \mathcal{G}(x_{n_k})],$$

that is equivalent to

$$\langle x_{n_k} - \lambda_{n_k} \mathcal{G}(x_{n_k}) - y_{n_k}, y - y_{n_k} \rangle \leq 0, \quad \forall y \in \mathcal{C}. \quad (3.7)$$

The inequality mentioned above implies that

$$\langle x_{n_k} - y_{n_k}, y - y_{n_k} \rangle \leq \lambda_{n_k} \langle \mathcal{G}(x_{n_k}), y - y_{n_k} \rangle, \quad \forall y \in \mathcal{C}. \quad (3.8)$$

Thus, we obtain

$$\frac{1}{\lambda_{n_k}} \langle x_{n_k} - y_{n_k}, y - y_{n_k} \rangle + \langle \mathcal{G}(x_{n_k}), y_{n_k} - x_{n_k} \rangle \leq \langle \mathcal{G}(x_{n_k}), y - x_{n_k} \rangle, \quad \forall y \in \mathcal{C}. \quad (3.9)$$

Since $\min\{\frac{\mu}{L}, \lambda_1\} \leq \lambda \leq \lambda_1$ and $\{x_{n_k}\}$ is a bounded sequence, from the condition $\lim_{k \rightarrow +\infty} \|x_{n_k} - y_{n_k}\| = 0$ and $k \rightarrow +\infty$ in (3.9), we obtain

$$\liminf_{k \rightarrow +\infty} \langle \mathcal{G}(x_{n_k}), y - x_{n_k} \rangle \geq 0, \quad \forall y \in \mathcal{C}. \quad (3.10)$$

Moreover, we have

$$\begin{aligned} &\langle \mathcal{G}(y_{n_k}), y - y_{n_k} \rangle \\ &= \langle \mathcal{G}(y_{n_k}) - \mathcal{G}(x_{n_k}), y - x_{n_k} \rangle + \langle \mathcal{G}(x_{n_k}), y - x_{n_k} \rangle + \langle \mathcal{G}(y_{n_k}), x_{n_k} - y_{n_k} \rangle. \end{aligned} \quad (3.11)$$

Since $\lim_{k \rightarrow +\infty} \|x_{n_k} - y_{n_k}\| = 0$ and $\mathcal{G}$ is L -Lipschitz continuity on $\mathcal{H}$ implies that

$$\lim_{k \rightarrow +\infty} \|\mathcal{G}(x_{n_k}) - \mathcal{G}(y_{n_k})\| = 0, \quad (3.12)$$

which together with (3.11) and (3.12), we obtain

$$\liminf_{k \rightarrow +\infty} \langle \mathcal{G}(y_{n_k}), y - y_{n_k} \rangle \geq 0, \quad \forall y \in \mathcal{C}. \quad (3.13)$$

To prove further, let us take a positive sequence $\{\epsilon_k\}$ that is convergent to zero and decreasing. For each $\{\epsilon_k\}$, we denote by m_k the smallest positive integer such that

$$\langle \mathcal{G}(x_{n_i}), y - x_{n_i} \rangle + \epsilon_k > 0, \quad \forall i \geq m_k, \quad (3.14)$$

where the existence of m_k follows from (3.13). Since $\{\epsilon_k\}$ is decreasing, it is easy to see that the sequence $\{m_k\}$ is increasing.

Case I: If there is a subsequence $\{x_{n_{m_{k_j}}}\}$ of $\{x_{n_{m_k}}\}$ such that $\mathcal{G}(x_{n_{m_{k_j}}}) = 0$, for all j . Let $j \rightarrow +\infty$, we obtain

$$\langle \mathcal{G}(\hat{x}), y - \hat{x} \rangle = \lim_{j \rightarrow +\infty} \langle \mathcal{G}(x_{n_{m_{k_j}}}), y - \hat{x} \rangle = 0. \quad (3.15)$$

Thus, $\hat{x} \in \mathcal{C}$ and imply that $\hat{x} \in VI(\mathcal{C}, \mathcal{G})$.

Case II: If there exists $N_0 \in \mathbb{N}$ such that for all $n_{m_k} \geq N_0$, $\mathcal{G}(x_{n_{m_k}}) \neq 0$. Consider that

$$F_{n_{m_k}} = \frac{\mathcal{G}(x_{n_{m_k}})}{\|\mathcal{G}(x_{n_{m_k}})\|^2}, \quad \forall n_{m_k} \geq N_0. \quad (3.16)$$

Due to the above definition, we obtain

$$\langle \mathcal{G}(x_{n_{m_k}}), F_{n_{m_k}} \rangle = 1, \quad \forall n_{m_k} \geq N_0. \quad (3.17)$$

Moreover, expressions (3.14) and (3.17), for all $n_{m_k} \geq N_0$, we have

$$\langle \mathcal{G}(x_{n_{m_k}}), y + \epsilon_k F_{n_{m_k}} - x_{n_{m_k}} \rangle > 0. \quad (3.18)$$

Since $\mathcal{G}$ is quasimonotone, then

$$\langle \mathcal{G}(y + \epsilon_k F_{n_{m_k}}), y + \epsilon_k F_{n_{m_k}} - x_{n_{m_k}} \rangle > 0. \quad (3.19)$$

For all $n_{m_k} \geq N_0$, we have

$$\langle \mathcal{G}(y), y - x_{n_{m_k}} \rangle \geq \langle \mathcal{G}(y) - \mathcal{G}(y + \epsilon_k F_{n_{m_k}}), y + \epsilon_k F_{n_{m_k}} - x_{n_{m_k}} \rangle - \epsilon_k \langle \mathcal{G}(y), F_{n_{m_k}} \rangle. \quad (3.20)$$

Due to $\{x_{n_k}\}$ weakly converges to $\hat{x} \in \mathcal{C}$ through $\mathcal{G}$ is weakly sequentially continuous on the set $\mathcal{C}$, we get $\{\mathcal{G}(x_{n_k})\}$ weakly converges to $\mathcal{G}(\hat{x})$. Suppose that $\mathcal{G}(\hat{x}) \neq 0$, we have

$$\|\mathcal{G}(\hat{x})\| \leq \liminf_{k \rightarrow +\infty} \|\mathcal{G}(x_{n_k})\|. \quad (3.21)$$

Since $\{x_{n_{m_k}}\} \subset \{x_{n_k}\}$ and $\lim_{k \rightarrow +\infty} \epsilon_k = 0$, we have

$$0 \leq \lim_{k \rightarrow +\infty} \|\epsilon_k F_{n_{m_k}}\| = \lim_{k \rightarrow +\infty} \frac{\epsilon_k}{\|\mathcal{G}(x_{n_{m_k}})\|} \leq \frac{0}{\|\mathcal{G}(\hat{x})\|} = 0. \quad (3.22)$$

Next, consider $k \rightarrow +\infty$ in (3.20), we obtain

$$\langle \mathcal{G}(y), y - \hat{x} \rangle \geq 0, \quad \forall y \in \mathcal{C}. \quad (3.23)$$

Thus, we infer that $\hat{x} \in VI(\mathcal{C}, \mathcal{G})$. $\square$

Theorem 3.3. *Assume that a mapping $\mathcal{G} : \mathcal{H} \rightarrow \mathcal{H}$ satisfies the conditions (G1)–(G4). Then, the sequence $\{x_n\}$ generated by the Algorithm 1 converges strongly to $x^* = P_\Omega(x_0)$.*

Proof. By using Lemma 3.1, we have

$$\|z_n - x^*\|^2 \leq \|x_n - x^*\|^2 - (1 - \lambda^2 L^2) \|x_n - y_n\|^2. \quad (3.24)$$

Since $0 < \lambda < \frac{1}{L}$, we obtain

$$\|z_n - x^*\|^2 \leq \|x_n - x^*\|^2. \quad (3.25)$$

By the use of definition of $\{x_{n+1}\}$, we obtain

$$\begin{aligned} \|x_{n+1} - x^*\| &= \|\alpha_n x_0 + (1 - \alpha_n) z_n - x^*\| \\ &= \|\alpha_n [x_0 - x^*] + (1 - \alpha_n) [z_n - x^*]\| \\ &\leq \alpha_n \|x_0 - x^*\| + (1 - \alpha_n) \|z_n - x^*\|. \end{aligned} \quad (3.26)$$

Combining expressions (3.25) and (3.26), we obtain

$$\begin{aligned} \|x_{n+1} - x^*\| &\leq \alpha_n \|x_0 - x^*\| + (1 - \alpha_n) \|x_n - x^*\| \\ &\leq \max \left\{ \|x_0 - x^*\|, \|x_n - x^*\| \right\}. \end{aligned} \quad (3.27)$$

By induction, we have

$$\|x_n - x^*\| \leq \|x_0 - x^*\|. \quad (3.28)$$

Thus, we conclude that $\{x_n\}$ is bounded sequence. By using Lemma 2.4 (i), we obtain

$$\begin{aligned} &\|x_{n+1} - x^*\|^2 \\ &= \|\alpha_n x_0 + (1 - \alpha_n) z_n - x^*\|^2 \\ &= \|\alpha_n [x_0 - x^*] + (1 - \alpha_n) [z_n - x^*]\|^2 \\ &= \alpha_n \|x_0 - x^*\|^2 + (1 - \alpha_n) \|z_n - x^*\|^2 - \alpha_n (1 - \alpha_n) \|x_0 - z_n\|^2 \\ &\leq \alpha_n \|x_0 - x^*\|^2 + (1 - \alpha_n) \left[\|x_n - x^*\|^2 - (1 - \lambda^2 L^2) \|x_n - y_n\|^2 \right] \\ &\quad - \alpha_n (1 - \alpha_n) \|x_0 - z_n\|^2 \\ &\leq \alpha_n \|x_0 - x^*\|^2 + \|x_n - x^*\|^2 - (1 - \alpha_n) (1 - \lambda^2 L^2) \|x_n - y_n\|^2. \end{aligned} \quad (3.29)$$

The above relationship implies that

$$(1 - \alpha_n) (1 - \lambda^2 L^2) \|x_n - y_n\|^2 \leq \alpha_n \|x_0 - x^*\|^2 + \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2. \quad (3.30)$$

From Lemma 2.4, we have

$$\begin{aligned}
& \|x_{n+1} - x^*\|^2 \\
&= \|\alpha_n x_0 + (1 - \alpha_n)z_n - x^*\|^2 \\
&= \|\alpha_n[x_0 - x^*] + (1 - \alpha_n)[z_n - x^*]\|^2 \\
&\leq (1 - \alpha_n)^2 \|z_n - x^*\|^2 + 2\alpha_n \langle x_0 - x^*, (1 - \alpha_n)[z_n - x^*] + \alpha_n[x_0 - x^*] \rangle \\
&= (1 - \alpha_n)^2 \|z_n - x^*\|^2 + 2\alpha_n \langle x_0 - x^*, x_{n+1} - x^* \rangle \\
&\leq (1 - \alpha_n) \|x_n - x^*\|^2 + 2\alpha_n \langle x_0 - x^*, x_{n+1} - x^* \rangle.
\end{aligned} \tag{3.31}$$

Case 1: Assume that there exists a fixed number $n_1 \in \mathbb{N}$ such that

$$\|x_{n+1} - x^*\| \leq \|x_n - x^*\|, \quad \forall n \geq n_1. \tag{3.32}$$

Thus, above implies that $\lim_{n \rightarrow +\infty} \|x_n - x^*\|$ exists and let $\lim_{n \rightarrow +\infty} \|x_n - x^*\| = l$. From (3.30), we obtain

$$\begin{aligned}
& (1 - \alpha_n)(1 - \lambda^2 L^2) \|x_n - y_n\|^2 \\
&\leq \alpha_n \|x_0 - x^*\|^2 + \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2.
\end{aligned} \tag{3.33}$$

The existence of $\lim_{n \rightarrow +\infty} \|x_n - x^*\| = l$ and $\alpha_n \rightarrow 0$, we can deduce that

$$\lim_{n \rightarrow +\infty} \|x_n - y_n\| = 0. \tag{3.34}$$

Thus, we have

$$\lim_{n \rightarrow +\infty} \|y_n - x^*\| = l. \tag{3.35}$$

It follows that

$$\|z_n - y_n\| = \|y_n + \lambda[\mathcal{G}(x_n) - \mathcal{G}(y_n)] - y_n\| \leq \lambda L \|x_n - y_n\|.$$

The above expression implies that

$$\lim_{n \rightarrow +\infty} \|z_n - y_n\| = 0. \tag{3.36}$$

It follows that

$$\lim_{n \rightarrow +\infty} \|x_n - z_n\| \leq \lim_{n \rightarrow +\infty} \|x_n - y_n\| + \lim_{n \rightarrow +\infty} \|y_n - z_n\| = 0. \tag{3.37}$$

Furthermore, we obtain

$$\begin{aligned}
\|x_{n+1} - x_n\| &= \|\alpha_n x_0 + (1 - \alpha_n)z_n - x_n\| \\
&= \|\alpha_n[x_0 - x_n] + (1 - \alpha_n)[z_n - x_n]\| \\
&\leq \alpha_n \|x_0 - x_n\| + (1 - \alpha_n) \|z_n - x_n\|.
\end{aligned} \tag{3.38}$$

It follows that

$$\lim_{n \rightarrow +\infty} \|x_{n+1} - x_n\| = 0. \tag{3.39}$$

It is given that $x^* = P_\Omega(x_0)$, we have

$$\langle x_0 - x^*, y - x^* \rangle \leq 0, \quad \forall y \in \Omega. \quad (3.40)$$

Indeed, since $\{x_n\}$ is bounded, we assume that there exists a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $x_{n_k} \rightharpoonup \hat{x} \in \mathcal{H}$. By Lemma 3.2, we have

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \langle x_0 - x^*, x_n - x^* \rangle &= \limsup_{k \rightarrow +\infty} \langle x_0 - x^*, x_{n_k} - x^* \rangle \\ &= \langle x_0 - x^*, \hat{x} - x^* \rangle \\ &\leq 0. \end{aligned} \quad (3.41)$$

By the use of $\lim_{n \rightarrow +\infty} \|x_{n+1} - x_n\| = 0$. We can deduce that

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \langle x_0 - x^*, x_{n+1} - x^* \rangle &\leq \limsup_{n \rightarrow +\infty} \langle x_0 - x^*, x_{n+1} - x_n \rangle \\ &\quad + \limsup_{n \rightarrow +\infty} \langle x_0 - x^*, x_n - x^* \rangle \\ &\leq 0. \end{aligned} \quad (3.42)$$

By the use of expressions (3.31), (3.42) and Lemma 2.8, we can derive that

$$\|x_n - x^*\| \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

Case 2: Assume there exists a subsequence $\{n_i\}$ of $\{n\}$ such that

$$\|x_{n_i} - x^*\| \leq \|x_{n_{i+1}} - x^*\|, \quad \forall i \in \mathbb{N}.$$

Thus, by Lemma 2.9, there exists a sequence $\{m_k\} \subset \mathbb{N}$ as $\{m_k\} \rightarrow +\infty$, such that

$$\|x_{m_k} - x^*\| \leq \|x_{m_{k+1}} - x^*\| \quad \text{and} \quad \|x_k - x^*\| \leq \|x_{m_{k+1}} - x^*\|, \quad \text{for all } k \in \mathbb{N}. \quad (3.43)$$

As similar to Case 1, expression (3.30) provides that

$$\begin{aligned} &(1 - \alpha_{m_k})(1 - \lambda^2 L^2) \|x_{m_k} - y_{m_k}\|^2 \\ &\leq \alpha_{m_k} \|x_0 - x^*\|^2 + \|x_{m_k} - x^*\|^2 - \|x_{m_{k+1}} - x^*\|^2. \end{aligned} \quad (3.44)$$

Due to $\alpha_{m_k} \rightarrow 0$, we deduce the following:

$$\lim_{k \rightarrow +\infty} \|x_{m_k} - y_{m_k}\| = 0. \quad (3.45)$$

It follows that

$$\|z_{m_k} - y_{m_k}\| = \|y_{m_k} + \lambda[\mathcal{G}(x_{m_k}) - \mathcal{G}(y_{m_k})] - y_{m_k}\| \leq \lambda L \|x_{m_k} - y_{m_k}\|.$$

The above expression implies that

$$\lim_{k \rightarrow +\infty} \|z_{m_k} - y_{m_k}\| = 0. \quad (3.46)$$

It follows that

$$\lim_{k \rightarrow +\infty} \|x_{m_k} - z_{m_k}\| \leq \lim_{k \rightarrow +\infty} \|x_{m_k} - y_{m_k}\| + \lim_{k \rightarrow +\infty} \|y_{m_k} - z_{m_k}\| = 0. \quad (3.47)$$

Furthermore, we obtain

$$\begin{aligned} \|x_{m_{k+1}} - x_{m_k}\| &= \|\alpha_{m_k}x_0 + (1 - \alpha_{m_k})z_{m_k} - x_{m_k}\| \\ &= \|\alpha_{m_k}[x_0 - x_{m_k}] + (1 - \alpha_{m_k})[z_{m_k} - x_{m_k}]\| \\ &\leq \alpha_{m_k}\|x_0 - x_{m_k}\| + (1 - \alpha_{m_k})\|z_{m_k} - x_{m_k}\|. \end{aligned} \quad (3.48)$$

It follows that

$$\lim_{k \rightarrow +\infty} \|x_{m_{k+1}} - x_{m_k}\| = 0. \quad (3.49)$$

We use the same argument as in Case 1, which is as follows:

$$\limsup_{k \rightarrow +\infty} \langle x_0 - x^*, x_{m_{k+1}} - x^* \rangle \leq 0. \quad (3.50)$$

Now, using expressions (3.31), we have

$$\begin{aligned} \|x_{m_{k+1}} - x^*\|^2 &\leq (1 - \alpha_{m_k})\|x_{m_k} - x^*\|^2 + 2\alpha_{m_k}\langle x_0 - x^*, x_{m_{k+1}} - x^* \rangle \\ &\leq (1 - \alpha_{m_k})\|x_{m_{k+1}} - x^*\|^2 + 2\alpha_{m_k}\langle x_0 - x^*, x_{m_{k+1}} - x^* \rangle. \end{aligned} \quad (3.51)$$

It continues from that

$$\|x_{m_{k+1}} - x^*\|^2 \leq 2\langle x_0 - x^*, x_{m_{k+1}} - x^* \rangle. \quad (3.52)$$

Thus, (3.42) and (3.52) we obtain

$$\|x_{m_{k+1}} - x^*\|^2 \rightarrow 0, \text{ as } k \rightarrow +\infty. \quad (3.53)$$

It implies that

$$\lim_{k \rightarrow +\infty} \|x_k - x^*\|^2 \leq \lim_{k \rightarrow +\infty} \|x_{m_{k+1}} - x^*\|^2 \leq 0. \quad (3.54)$$

Consequently, $x_n \rightarrow x^*$. This completes the proof. $\square$

Next, we introduce the first variant of Algorithm 1 in which the constant step size λ is chosen adaptively and thus produced a sequence $\{\lambda_n\}$ that does not require the knowledge of the Lipschitz-type constant L .

Algorithm 2.

Step 0: Choose $x_1 \in \mathcal{C}$, $\lambda_1 > 0$, $\mu \in (0, 1)$ and $\{\alpha_n\} \subset (0, 1)$ satisfies the following conditions:

$$\lim_{n \rightarrow +\infty} \alpha_n = 0 \text{ and } \sum_{n=1}^{+\infty} \alpha_n = +\infty.$$

Step 1: Compute

$$y_n = P_{\mathcal{C}}(x_n - \lambda \mathcal{G}(x_n)).$$

If $x_n = y_n$, then STOP and y_n is a solution. Otherwise, go to **Step 2**.

Step 2: Compute

$$z_n = y_n + \lambda [\mathcal{G}(x_n) - \mathcal{G}(y_n)].$$

Step 3: Compute

$$x_{n+1} = \alpha_n x_0 + (1 - \alpha_n) z_n.$$

Step 4: Compute

$$\lambda_{n+1} = \begin{cases} \min \left\{ \lambda_n, \frac{\mu \|x_n - y_n\|}{\|\mathcal{G}(x_n) - \mathcal{G}(y_n)\|} \right\} & \text{if } \mathcal{G}(x_n) - \mathcal{G}(y_n) \neq 0, \\ \lambda_n & \text{otherwise.} \end{cases} \quad (3.55)$$

Set $n = n + 1$ and go back to **Step 1**.

Lemma 3.4. *The sequence $\{\lambda_n\}$ generated by (3.55) is decreasing monotonically and converges to $\lambda > 0$.*

Proof. It is given that $\mathcal{G}$ is Lipschitz-continuous with constant $L > 0$. Let $\mathcal{G}(x_n) \neq \mathcal{G}(y_n)$ such that

$$\begin{aligned} \frac{\mu \|x_n - y_n\|}{\|\mathcal{G}(x_n) - \mathcal{G}(y_n)\|} &\geq \frac{\mu \|x_n - y_n\|}{L \|x_n - y_n\|} \\ &\geq \frac{\mu}{L}. \end{aligned} \quad (3.56)$$

The above expression implies that $\lim_{n \rightarrow +\infty} \lambda_n = \lambda$. $\square$

Lemma 3.5. *Suppose that $\mathcal{G} : \mathcal{H} \rightarrow \mathcal{H}$ satisfies conditions (G1)-(G4) and sequence $\{x_n\}$ generated by Algorithm 2. Then, we have*

$$\|x_{n+1} - x^*\|^2 \leq \|x_n - x^*\|^2 - \left(1 - \mu^2 \frac{\lambda_n^2}{\lambda_{n+1}^2}\right) \|x_n - y_n\|^2.$$

Next, we introduce the second variant of Algorithm 1 in which the constant step size λ is chosen adaptively and thus produced a sequence $\{\lambda_n\}$ that does not require the knowledge of the Lipschitz-like constants L . In this case, the step size sequence $\{\lambda_n\}$ is not monotone.

Algorithm 3.

Step 0: Choose $x_1 \in \mathcal{C}$, $\lambda_1 > 0$, $\mu \in (0, 1)$ and select a nonnegative real sequence $\{\varphi_n\}$ such that $\sum_{n=1}^{+\infty} \varphi_n < +\infty$. Moreover, choose $\{\alpha_n\} \subset (0, 1)$ satisfies the following conditions:

$$\lim_{n \rightarrow +\infty} \alpha_n = 0 \quad \text{and} \quad \sum_{n=1}^{+\infty} \alpha_n = +\infty.$$

Step 1: Compute

$$y_n = P_C(x_n - \lambda \mathcal{G}(x_n)).$$

If $x_n = y_n$, then STOP and y_n is a solution. Otherwise, go to **Step 2**.

Step 2: Compute

$$z_n = y_n + \lambda [\mathcal{G}(x_n) - \mathcal{G}(y_n)].$$

Step 3: Compute

$$x_{n+1} = \alpha_n x_0 + (1 - \alpha_n) z_n.$$

Step 4: Compute

$$\lambda_{n+1} = \begin{cases} \min \left\{ \lambda_n + \varphi_n, \frac{\mu \|x_n - y_n\|}{\|\mathcal{G}(x_n) - \mathcal{G}(y_n)\|} \right\} & \text{if } \mathcal{G}(x_n) - \mathcal{G}(y_n) \neq 0, \\ \lambda_n + \varphi_n & \text{otherwise.} \end{cases} \quad (3.57)$$

Set $n = n + 1$ and go back to **Step 1**.

Lemma 3.6. *The sequence $\{\lambda_n\}$ generated by (3.57) is convergent to λ and also satisfy the following inequality*

$$\min \left\{ \frac{\mu}{L}, \lambda_0 \right\} \leq \lambda \leq \lambda_0 + P, \quad \text{where } P = \sum_{n=1}^{+\infty} \varphi_n.$$

Proof. Due to the Lipschitz continuity of a mapping $\mathcal{G}$ there exists a fixed number $L > 0$. Consider that $\mathcal{G}(x_n) - \mathcal{G}(y_n) \neq 0$ such that

$$\begin{aligned} \frac{\mu \|x_n - y_n\|}{\|\mathcal{G}(x_n) - \mathcal{G}(y_n)\|} &\geq \frac{\mu \|x_n - y_n\|}{L \|x_n - y_n\|} \\ &\geq \frac{\mu}{L}. \end{aligned} \quad (3.58)$$

By using mathematical induction on the definition of λ_{n+1} , we have

$$\min \left\{ \frac{\mu}{L}, \lambda_0 \right\} \leq \lambda_n \leq \lambda_0 + P.$$

Let

$$[\lambda_{n+1} - \lambda_n]^+ = \max \{0, \lambda_{n+1} - \lambda_n\}$$

and

$$[\lambda_{n+1} - \lambda_n]^- = \max \{0, -(\lambda_{n+1} - \lambda_n)\}.$$

From the definition of $\{\lambda_n\}$, we have

$$\sum_{n=1}^{+\infty} [\lambda_{n+1} - \lambda_n]^+ = \sum_{n=1}^{+\infty} \max \{0, \lambda_{n+1} - \lambda_n\} \leq P < +\infty. \quad (3.59)$$

That is, the series $\sum_{n=1}^{+\infty} [\lambda_{n+1} - \lambda_n]^+$ is convergent.

Next we need to prove the convergence of $\sum_{n=1}^{+\infty} (\lambda_{n+1} - \lambda_n)^-$.

Let $\sum_{n=1}^{+\infty} [\lambda_{n+1} - \lambda_n]^- = +\infty$. Due to the reason that

$$\lambda_{n+1} - \lambda_n = (\lambda_{n+1} - \lambda_n)^+ - (\lambda_{n+1} - \lambda_n)^-,$$

we have

$$\lambda_{k+1} - \lambda_0 = \sum_{n=0}^k (\lambda_{n+1} - \lambda_n) = \sum_{n=0}^k [\lambda_{n+1} - \lambda_n]^+ - \sum_{n=0}^k [\lambda_{n+1} - \lambda_n]^-. \quad (3.60)$$

By allowing $k \rightarrow +\infty$ in (3.60), we have $\lambda_k \rightarrow -\infty$ as $k \rightarrow +\infty$. This is a contradiction. Due to the convergence of the series $\sum_{n=0}^k [\lambda_{n+1} - \lambda_n]^+$ and

$\sum_{n=0}^k [\lambda_{n+1} - \lambda_n]^-$ taking $k \rightarrow +\infty$ in (3.60), we obtain $\lim_{n \rightarrow +\infty} \lambda_n = \lambda$. This completes the proof. $\square$

Remark 3.7. (i) Three extragradient-type methods are established to find an approximate solution of variational inequalities involving semistrictly quasimonotone and Lipschitz-continuous operators in a real Hilbert space.

(ii) A strongly convergent result, corresponding to the proposed algorithms have been proved.

(iii) It is important to note that these methods have been used fixed, monotonic and non-monotonic step size rules that use operator value rather than the Lipschitz constant of an operator.

4. NUMERICAL ILLUSTRATIONS

The computational results of the proposed schemes are described in this section, in contrast to some related work in the literature and also in the analysis of how variations in control parameters affect the numerical effectiveness of the proposed algorithms. All computations are done in MATLAB R2018b and run on HP i- 5 Core(TM)i5-6200 8.00 GB (7.78 GB usable) RAM laptop.

Example 4.1. Consider that $\mathcal{H} = l_2$ is a real Hilbert space with sequences of real numbers satisfying the following condition

$$\|x_1\|^2 + \|x_2\|^2 + \cdots + \|x_n\|^2 + \cdots < +\infty. \quad (4.1)$$

Assume that $\mathcal{G} : \mathcal{C} \rightarrow \mathcal{C}$ is defined by

$$G(x) = (5 - \|x\|)x, \quad \forall x \in \mathcal{H},$$

where $C = \{x \in \mathcal{H} : \|x\| \leq 3\}$. It is easy to see that $\mathcal{G}$ is weakly sequentially continuous on $\mathcal{H}$ and $VI(\mathcal{C}, \mathcal{G}) = \{0\}$. For any $x, y \in \mathcal{H}$, we have

$$\begin{aligned} \|\mathcal{G}(x) - \mathcal{G}(y)\| &= \|(5 - \|x\|)x - (5 - \|y\|)y\| \\ &= \|5(x - y) - \|x\|(x - y) - (\|x\| - \|y\|)y\| \\ &\leq 5\|x - y\| + \|x\|\|x - y\| + \|\|x\| - \|y\|\|\|y\| \\ &\leq 5\|x - y\| + 3\|x - y\| + 3\|x - y\| \\ &\leq 11\|x - y\|. \end{aligned} \tag{4.2}$$

Hence $\mathcal{G}$ is L -Lipschitz continuous with $L = 11$. For any $x, y \in \mathcal{H}$, let $\langle \mathcal{G}(x), y - x \rangle > 0$ such that

$$(5 - \|x\|)\langle x, y - x \rangle > 0.$$

Since $\|x\| \leq 3$ implies that

$$\langle x, y - x \rangle > 0.$$

Thus, we have

$$\begin{aligned} \langle \mathcal{G}(y), y - x \rangle &= (5 - \|y\|)\langle y, y - x \rangle \\ &\geq (5 - \|y\|)\langle y, y - x \rangle - (5 - \|y\|)\langle x, y - x \rangle \\ &\geq 2\|x - y\|^2 \geq 0. \end{aligned} \tag{4.3}$$

Thus, we shown that $\mathcal{G}$ is quasimonotone on $\mathcal{C}$. Let $x = (\frac{5}{2}, 0, 0, \dots, 0, \dots)$ and $y = (3, 0, 0, \dots, 0, \dots)$ such that

$$\langle \mathcal{G}(x) - \mathcal{G}(y), x - y \rangle = (2.5 - 3)^2 < 0.$$

A projection on the set C is computed explicitly as follows:

$$P_C(x) = \begin{cases} x, & \text{if } \|x\| \leq 3, \\ \frac{3x}{\|x\|}, & \text{otherwise.} \end{cases}$$

The control conditions have been taken as follows:

$$\lambda = \frac{1}{2L}, \quad \lambda_0 = \frac{5}{11}, \quad \mu = 0.44, \quad \varphi_n = \frac{100}{(n+3)^2}.$$

TABLE 1. Numerical results values for Example 4.1.

x_0	Number of Iterations		
	Algorithm 3	Algorithm 3	Algorithm 3
$(3, 3, \dots, 3_{10000}, 0, 0, \dots)$	102	91	89
$(1, 2, \dots, 10000, 0, 0, \dots)$	112	94	99
$(7, 7, \dots, 7_{100000}, 0, 0, \dots)$	117	101	110
$(20, 20, \dots, 20_{100000}, 0, 0, \dots)$	129	109	115

TABLE 2. Numerical results values for Example 4.1.

x_0	Elapsed time in seconds		
	Algorithm 3	Algorithm 3	Algorithm 3
$(3, 3, \dots, 3_{10000}, 0, 0, \dots)$	15.658362	13.575637	14.0057768
$(1, 2, \dots, 10000, 0, 0, \dots)$	16.274758	14.019125	15.0084828
$(7, 7, \dots, 7_{100000}, 0, 0, \dots)$	18.564383	17.476847	17.5384939
$(20, 20, \dots, 20_{100000}, 0, 0, \dots)$	22.657362	19.475593	21.5701981

Remark 4.2. (i) We observe from numerical results of Example 4.1 that our proposed Algorithm 3 is efficient and easy to implement for both finite and infinite dimensional spaces, see Tables 1 and 2.

(ii) The performance is better both in CPU time and the number of iterations for Example 4.1 (see Tables 1 and 2).

(iii) We also observe that different choice of initial points x_0 has significant effect of the CPU (time) and number of iterations.

CONCLUSION

In this study, we considered three strong convergence results for variational inequalities problem involving semistrictly quasimonotone and Lipschitz continuous monotone operator, but the Lipschitz constant is unknown. We modify the extragradient method with a natural step size rule. The strong convergence result is demonstrated without any provision of additional projections.

Acknowledgement: Nuttapol Pakkaranang was partially financial supported by Phetchabun Rajabhat University.

REFERENCES

- [1] J. Abubakar, P. Kumam, A.H. Ibrahim, and A. Padcharoen, *Relaxed inertial Tseng's type method for solving the inclusion problem with application to image restoration*, Mathematics, **8**(5) (2020), Article ID 818.
- [2] A.S. Antipin, *On a method for convex programs using a symmetrical modification of the Lagrange function*, Ekonomika i Matem. Metody., **12** (1976), 1164–1173.
- [3] H.H. Bauschke and P.L. Combettes, *Convex analysis and monotone operator theory in Hilbert spaces*, Springer, New York, 2011.
- [4] Y. Censor, A. Gibali and S. Reich, *The subgradient extragradient method for solving variational inequalities in Hilbert space*, J. Optim. Theory Appl., **148**(2) (2010), 318–335.
- [5] Y. Censor, A. Gibali and S. Reich, *Strong convergence of subgradient extragradient methods for the variational inequality problem in Hilbert space*, Optim. Methods Softw., **26**(4-5) (2011), 827–845.
- [6] P. Cholamjiak, D.V. Hieu and Y.J. Cho, *Relaxed Forward-Backward Splitting Methods for Solving Variational Inclusions and Applications*, J. Sci. Comput., **88** (2021). <https://doi.org/10.1007/s10915-021-01608-7>.

- [7] A. Daniilidis and N. Hadjisavvas, *Characterization of nonsmooth semistrictly quasiconvex and strictly quasiconvex functions*, J. Optim. Theory Appl., **102**(3) (1999), 525–536.
- [8] C.M. Elliott, *Variational and quasivariational inequalities: applications to free-boundary problems (Claudio Baiocchi and antónio Capelo)*, SIAM Rev., **29**(2) (1987), 314–315.
- [9] N. Hadjisavvas and S. Schaible, *On strong pseudomonotonicity and (semi)strict quasimonotonicity*, J. Optim. Theory Appl., **79**(1) (1993), 139–155.
- [10] B. Halpern, *Fixed points of nonexpanding maps*, Bull. Amer. Math. Soc., **73** (1967), 957–961.
- [11] D.V. Hieu and P. Chalamjiak, *Modified extragradient method with Bregman distance for variational inequalities*, Appl. Anal., (2020), Accepted. <https://doi.org/10.1080/00036811.2020.1757078>.
- [12] A.N. Iusem and B.F. Svaiter, *A variant of Korpelevich's method for variational inequalities with a new search strategy*, Optimization, **42**(4) (1997), 309–321.
- [13] G. Kassay, J. Kolumbán and Z. Páles, *On Nash stationary points*, Publ. Math. Debrecen, **54**(3-4) (1999), 267–279.
- [14] G. Kassay, J. Kolumbán and Z. Páles, *Factorization of minty and stampanchia variational inequality systems*, European J. Oper. Res., **143**(2) (2002), 377–389.
- [15] J.K. Kim and Salahuddin, *Extragradient methods for generalized mixed equilibrium problems and fixed point problem in Hilbert spaces*, Nonlinear Funct. Anal. Appl., **22**(4) (2017), 693–709.
- [16] D. Kinderlehrer and G. Stampacchia, *An introduction to variational inequalities and their applications*, Classic in Applied Mathematics, Society for Industrial and Applied Mathematics, (SIAM), Philadelphia, PA, 2000.
- [17] I. Konnov, *Equilibrium models and variational inequalities*, **210**, Elsevier, 2007.
- [18] G. Korpelevich, *The extragradient method for finding saddle points and other problems*, Ékonom. i Mat. Metody, **12**(4) (1976), 747–756.
- [19] P.-E. Maingé, *Strong convergence of projected subgradient methods for nonsmooth and nonstrictly convex minimization*, Set-Valued Anal., **16**(7-8) (2008), 899–912.
- [20] Y.V. Malitsky and V.V. Semenov, *An extragradient algorithm for monotone variational inequalities*, Cybernet. Systems Anal., **50**(2) (2014), 271–277.
- [21] A. Moudafi, *Viscosity approximation methods for fixed-points problems*, J. Math. Anal. Appl., **241**(1) (2000), 46–55.
- [22] K. Muangchoo, *A Viscosity type projection method for solving pseudomonotone variational inequalities*, Nonlinear Funct. Anal. Appl., **26**(2) (2021), 347–371.
- [23] M.A. Noor, *Some iterative methods for nonconvex variational inequalities*, Comput. Math. Model., **21**(1) (2010), 97–108.
- [24] A. Padcharoen, D. Kitkuan, W. Kumam and P. Kumam, *Tseng methods with inertial for solving inclusion problems and application to image deblurring and image recovery problems*, Comput. Math. Meth., **3** (2020), e1088. <https://doi.org/10.1002/cmm4.1088>.
- [25] N. Pakkaranang, H. ur Rehman, and W. Kumam, *Two strongly convergent self-adaptive iterative schemes for solving pseudo-monotone equilibrium problems with applications*, Demonstr. Math., **54**(1) (2021), 280–298.
- [26] S. Reich, D.V. Thong, P. Chalamjiak and L.V. Long, *Inertial projection-type methods for solving pseudomonotone variational inequality problems in Hilbert space*, Numer. Algorithms, **88**(2) (2021), 813–835.
- [27] H. ur Rehman, N. Pakkaranang, A. Hussain and N. Wairojjana, *A modied extra-gradient method for a family of strongly pseudomonotone equilibrium problems in real Hilbert spaces*, J. Math. Comput. Sci., **22**(1) (2021), 38–48.

- [28] H. ur Rehman, P. Kumam, I.K. Argyros, W. Deebani and W. Kumam, *Inertial extra-gradient method for solving a family of strongly pseudomonotone equilibrium problems in real hilbert spaces with application in variational inequality problem*, Symmetry, **12**(4) (2020), Article ID 503. <https://doi.org/10.3390/sym12040503>.
- [29] H. ur Rehman, P. Kumam, Y.J. Cho, Y.I. Suleiman and W. Kumam, *Modified Popov's explicit iterative algorithms for solving pseudomonotone equilibrium problems*, Optim. Methods Softw., **36**(1) (2021), 82–113.
- [30] H. ur Rehman, P. Kumam, Y.J. Cho and P. Yordsorn, *Weak convergence of explicit extragradient algorithms for solving equilibrium problems*, J. Inequal. Appl., **2019** (2019), Article ID 282. <https://doi.org/10.1186/s13660-019-2233-1>.
- [31] H. ur Rehman, P. Kumam, A. Gibali, and W. Kumam, *Convergence analysis of a general inertial projection-type method for solving pseudomonotone equilibrium problems with applications*, J. Inequal. Appl., **2021** (2021), Article ID 63. <https://doi.org/10.1186/s13660-021-02591-1>.
- [32] H. ur Rehman, P. Kumam, W. Kumam, M. Shutaywi and W. Jirakitpuwapat, *The inertial sub-gradient extra-gradient method for a class of pseudo-monotone equilibrium problems*, Symmetry, **12**(3) (2020), Article ID 463. <https://doi.org/10.3390/sym12030463>.
- [33] G. Stampacchia, *Formes bilinéaires coercitives sur les ensembles convexes*, C. R. Acad. Sci. Paris, **258** (1964), 4413–4416.
- [34] P. Tseng, *A modified forward-backward splitting method for maximal monotone mappings*, SIAM J. Control Optim., **38**(2) (2000), 431–446.
- [35] N. Wairojjana, H. ur Rehman, M.D. la Sen and N. Pakkaranang, *A general inertial projection-type algorithm for solving equilibrium problem in Hilbert spaces with applications in fixed-point problems*, Axioms **9**(3) (2020), Article ID 101. <https://doi.org/10.3390/axioms9030101>.
- [36] H.-K. Xu, *Another control condition in an iterative method for nonexpansive mappings*, Bull. Austral. Math. Soc., **65**(1) (2002), 109–113.
- [37] L. Zhang, C. Fang and S. Chen, *An inertial subgradient-type method for solving single-valued variational inequalities and fixed point problems*, Numer. Algo., **79** (2018), 941–956.