



รายละเอียดของวารสาร**ชื่อวารสาร:****Journal Name:** Progress in Applied Science and Technology**ชื่อบรนการ:** Arranee Chotiko**ชื่อย่อของวารสาร:****Abbreviation Name:****ISSN:** 2730-3012**E-ISSN:** 2730-3020**ที่อยู่สำหรับการติดต่อ:** Progress in Applied Science and Technology, Faculty of Science and Technology, Rajamangala University of Technology Thanyaburi 39 Moo 1, Klong 6, Khlong Luang Pathum Thani 12110 Thailand**เจ้าของ:** คณะวิทยาศาสตร์และเทคโนโลยี มหาวิทยาลัยเทคโนโลยีราชมงคลรัตนบุรี / Faculty of Science and Technology, Rajamangala University of Technology Thanyaburi**จำนวนฉบับต่อปี:** 3**Email:** scitechrmutt@rmutt.ac.th**Website:** <https://ph02.tci-thaijo.org/index.php/past>**TCI กลุ่มที่:** 1**สาขาวิชาของวารสาร:** Physical Sciences**สาขาวิชของวารสาร:** Chemistry / Computer Science / Mathematics

**หมายเหตุ:**

- เริ่มใช้ ISSN และชื่อวารสารใหม่ ตั้งแต่ปีที่ 10 ฉบับที่ 2 กรกฎาคม 2563 เป็นต้นไป
- ISSN และชื่อวารสาร (เดิม) คือ ISSN : 2229-1547, วารสารวิทยาศาสตร์และเทคโนโลยี มก. รัตนบุรี | Science and Technology RMUTT Journal

**กลุ่มของวารสารในฐานข้อมูล TCI****Publication 10 Years**



## แบบตอบรับการตีพิมพ์บทความวิจัยในวารสารวิชาการ

วารสาร Progress in Applied Science and Technology

คณะวิทยาศาสตร์และเทคโนโลยี  
มหาวิทยาลัยเทคโนโลยีราชมงคลล้านนา  
39 หมู่ที่ 1 ถนนรังสิต-นครนายก ตำบลคลองหก  
อำเภอคลองหลวง จังหวัดปทุมธานี 12110

วันที่ 6 มิถุนายน 2566

เรื่อง การตอบรับบทความวิจัยเพื่อลงตีพิมพ์ในวารสารวิชาการ  
เรียน ผู้ช่วยศาสตราจารย์ ดร.สัจจารักษ์ ลัดสุขเนิน

ตามที่ท่านมีความประสงค์ส่งบทความวิจัยเรื่อง “Property A Characterization of  $S_\beta$ -continuous Fixed Point” ลงพิมพ์เผยแพร่ในวารสาร Progress in Applied Science and Technology ซึ่งเป็นวารสารวิชาการของคณะวิทยาศาสตร์และเทคโนโลยี มหาวิทยาลัยเทคโนโลยีราชมงคลล้านนา กองบรรณาธิการได้พิจารณาแล้วว่า บทความนี้มีคุณภาพดีและมีประโยชน์ต่อวิชาการ จึงอนุมัติให้ตีพิมพ์เผยแพร่ในวารสาร Progress in Applied Science and Technology Vol. 13 No. 2 (May – August 2023) เมื่อวันที่ 2 มิถุนายน 2566

จึงเรียนมาเพื่อโปรดทราบ

*On*

(ผู้ช่วยศาสตราจารย์ ดร.อรณี โชคติโก)

บรรณาธิการ

วารสาร Progress in Applied Science and Technology

กองบรรณาธิการ

วารสาร Progress in Applied Science and Technology

โทรศัพท์ 02-549-4176

โทรสาร 02-549-4119

Home (<https://ph02.tci-thaijo.org/index.php/past/index>)  
 / Archives (<https://ph02.tci-thaijo.org/index.php/past/issue/archive>)  
 / Vol. 13 No. 2 (2023): May - August 2023 (<https://ph02.tci-thaijo.org/index.php/past/issue/view/17027>)  
 / Mathematics and Applied Statistics

## A Characterization of S\_beta-continuous Fixed Point Property



(<https://ph02.tci-thaijo.org/index.php/past/issue/view/17027>)

PDF (<https://ph02.tci-thaijo.org/index.php/past/article/view/247828/169255>)

Published: Jul 20, 2023

DOI: <https://doi.org/10.14456/past.2023.12> (<https://doi.org/10.14456/past.2023.12>)

Keywords: <https://www.nstda.or.th/home/nstda-privacy-policy/>  
 S\_beta-open sets S\_beta-continuous function fixed point property S\_beta-continuous  
 fixed point property

Sajjarak Ladsungnern

Mathematics Program, Faculty of Education, Chaiyaphum Rajabhat University

Ardoon Jongrak

Mathematics Program, Faculty of Science and Technology, Phetchabun Rajabhat University

### Abstract

In this paper, we define and investigate the  $S_\beta$ -continuous retraction and the  $S_\beta$ -continuous fixed point property which apply the  $S_\beta$ -continuity in (6). The study shown that, for the regular and locally indiscrete topological space  $(X, \tau)$  with the  $S_\beta$ -continuous fixed point property, if a topology  $\sigma$  for  $X$  is stronger than  $\tau$  and for every open subset  $G$  in  $\sigma$  with the closure of  $G$  in  $\sigma$  and the closure of  $G$  in  $\tau$  are equal, then  $(X, \sigma)$  has the fixed point property.

#### How to Cite

Ladsungnern S, Jongrak A. A Characterization of S\_beta-continuous Fixed Point Property. Prog Appl Sci Tech. [Internet]. 2023 Jul. 20 [cited 2023 Oct. 10];13(2):9-16. Available from: <https://ph02.tci-thaijo.org/index.php/past/article/view/247828>

More Citation Formats ▾

Download Citation ▾

Submit  
Your  
Article



(<https://ph02.tci-thaijo.org/index.php/past/submit>)

Publication  
Ethics

Publication  
Ethics  
(<https://ph02.tci-thaijo.org/index.php/past/Ethics>)

Home  
ThaiJo

THAIJO  
(<https://www.tci-thaijo.org/>)

Journal  
Information

Info  
Tier  
(/public/api/infoTier.php)

Inde  
thai:  
◀ ▶

Editor  
:  
Arranee  
Chotiko

Google  
Scholar

Google Scholar  
([https://scholar.google.co.th/scholar?as\\_q=&as\\_epq=progress+in+applied+science+and+technol](https://scholar.google.co.th/scholar?as_q=&as_epq=progress+in+applied+science+and+technol))

Information

For  
Readers  
(<https://ph02.tci-thaijo.org/index.php/past/information/readers>)

For  
Author  
(<https://ph02.tci-thaijo.org/index.php/past/Author>)

Visitors  
Flag Counter  
(<https://info.flagcounter.com/EXWO>)

Issue
Vol. 13 No. 2 (2023): May - August 2023 ( <a href="https://ph02.tci-thaijo.org/index.php/past/issue/view/17027">https://ph02.tci-thaijo.org/index.php/past/issue/view/17027</a> )

Counter  
installed  
: 04 May  
2019

Section
Mathematics and Applied Statistics



(<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Current Issue

<https://ph02.tci-thaijo.org/index.php/past/gateway/plugin/WebFeedGatew>

<https://ph02.tci-thaijo.org/index.php/past/gateway/plugin/WebFeedGatew>

<https://ph02.tci-thaijo.org/index.php/past/gateway/plugin/WebFeedGatew>

## References

Adams CC, Franzosa RD. Introduction to topology pure and applied. New Jersey: Upper saddle river; 2008.

Cammarolo F, Noiri T. On the delta-continuous fixed point property. Int. J. Math. Math. Sci. 1990;13(1):45-50.

Connell EH. Property of fixed point spaces. Proc. Am. Math. Soc. 1959;10(3):974-979.

Dugundji J. Topology. Boston: Allyn and Bacon; 1966.

Jongrak A. Some properties of S\_beta-open Mappings. The proceedings of annual meeting in mathematics. 2017;22:TP0031-035.

Khalaf AB, Ahmed NK. S\_beta-open sets and S\_beta-continuity in topological spaces. Thai J. Math. 2013;11(2):319-335.

Levine N. Semi-open sets and semi-continuity in topological spaces. Am. Math. Mon. 1963;29(70):36-4.

Monsef ME, Deeb SN, Mahmoud RA. beta-open sets and beta-continuous mappings. Bullentin of the Faculty of Science, Assiut University. 1983;12(1):77-90.

Puturong N. On strongly theta-semi- continuous functions. Thai J. Math. 2007; 5(3):11-23.

Progress in Applied Science and Technology  
Faculty of Science and Technology (<http://www.sci.rmutt.ac.th/>)  
Rajamangala University of Technology Thanyaburi (<http://www.sci.rmutt.ac.th/>)  
39 Moo 1, Klong 6, Khlong Luang Pathum Thani 12110 Thailand  
Tel : +66(0)2 549 4150, 4164, 4176  
Fax : +66(0)2 549 4158, 4119  
E-mail : [scitech@rmutt.ac.th](mailto:scitech@rmutt.ac.th) (<mailto:scitech@rmutt.ac.th>)

<https://ph02.tci-thaijo.org/index.php/past/about/aboutThisPublishingSystem>

นโยบายการคุ้มครองข้อมูลส่วนบุคคล

(<https://www.nstda.or.th/home/nstda-privacy-policy/>)

Privacy policy (<https://www.nstda.or.th/en/nstda-privacy-policy.html>)

## Research Article

Received: December 06, 2022

Revised: April 12, 2023

Accepted: June 02, 2023

DOI: 10.14456/past.2023.12

## A Characterization of $S_\beta$ -continuous Fixed Point Property

Sajjarak Ladsungnern<sup>1</sup> and Ardoon Jongrak<sup>2\*</sup>

<sup>1</sup> Mathematics Program, Faculty of Education,  
Chaiyaphum Rajabhat University, Chaiyaphum 36000, Thailand

<sup>2</sup> Mathematics Program, Faculty of Science and Technology,  
Phetchabun Rajabhat University, Phetchabun 67000, Thailand

\*E-mail: ardoon.jon@pcru.ac.th

### Abstract

In this paper, we define and investigate the  $S_\beta$ -continuous retraction and the  $S_\beta$ -continuous fixed point property which apply the  $S_\beta$ -continuity in (6). The study shown that, for the regular and locally indiscrete topological space  $(X, \tau)$  with the  $S_\beta$ -continuous fixed point property, if a topology  $\sigma$  for  $X$  is stronger than  $\tau$  and for every open subset  $G$  in  $\sigma$  with the closure of  $G$  in  $\sigma$  and the closure of  $G$  in  $\tau$  are equal, then  $(X, \sigma)$  has the fixed point property.

**Keywords:**  $S_\beta$ -open sets,  $S_\beta$ -continuous function, fixed point property,  $S_\beta$ -continuous fixed point property

### 1. Introduction

For the two topological spaces  $(X, \tau)$  and  $(X, \sigma)$ , in 1959 Conell EH. (3) has introduced the concept of the fixed point property. If  $(X, \tau)$  is a regular space with the fixed point property,  $\sigma$  is stronger than  $\tau$ ,  $G$  is an open subset in  $\sigma$ , its closure is the same in both topologies, then  $(X, \sigma)$  has the fixed point property. In 1990, Cammaroto F. and Noiri T. (2) has introduced the concept of the  $\delta$ -continuous fixed point property. If  $(X, \tau)$  is an almost-regular space with the  $\delta$ -continuous fixed point property,  $\sigma$  is stronger than  $\tau$ ,  $G$  is an open subset in  $\sigma$ , its closure is the same in both topologies, then  $(X, \sigma)$  has the fixed point property. In 2007, Puturong N. (9) has introduced the concept of the strongly  $\theta$ -semi-continuous fixed point property. If  $(X, \tau)$  is a

regular space with the strongly  $\theta$ -semi-continuous fixed point property,  $\sigma$  is stronger than  $\tau$ ,  $G$  is an open subset in  $\sigma$ , its closure are the same in both topologies, then  $(X, \sigma)$  has the fixed point property. In 2013, Khalaf BA. and Ahmed KN. (6) has introduced  $S_\beta$ -continuous function  $f: (X, \tau) \rightarrow (Y, \sigma)$ , if for each  $x$  in  $X$  and for each open subset  $H$  in  $Y$  containing  $f(x)$ , there exists an  $S_\beta$ -open subset  $G$  in  $X$  containing  $x$  such that  $f(G) \subseteq H$ . The purpose of this paper is to apply  $S_\beta$ -continuity to the retraction and fixed point property. We shall offer some statements,  $(X, \tau)$  is a locally indiscrete space with  $S_\beta$ -continuous fixed point property and the same above conditions implies that  $(X, \sigma)$  has the fixed point property.

## 2. Preliminaries

Throughout the present note spaces always mean topological spaces on which no separation axioms are assumed unless explicitly stated. Recall the following definitions and characterizations. In a topological space  $(X, \tau)$  (or simply  $X$ ) and let  $A$  be a subset of  $X$ . The intersection of all closed sets containing  $A$  is called the closure of  $A$  and is denoted by  $cl(A)$ . The interior of  $A$  is defined by the union of all open sets contained in  $A$  and is denoted by  $int(A)$ . We shall define semi-open set, semi-closed set,  $\beta$ -closed set,  $S_\beta$ -open set, and  $S_\beta$ -closed set as follows.

**Definition 2.1.** (6-8) Let  $(X, \tau)$  be a topological space. A subset  $A$  of  $X$  is called

(i) *semi-open* if  $A \subseteq cl(int(A))$ , the complement of a semi-open set is called semi-closed set.

(ii)  *$\beta$ - open* if  $A \subseteq cl(int(cl(A)))$ , the complement of a  $\beta$ - open set is called  $\beta$ -closed set.

(iii)  $S_\beta$ -open if  $A$  is a semi-open set and for each  $x \in A$  there exists a  $\beta$ -closed set  $F$  such that  $x \in F \subseteq A$ , and  $B \subseteq X$  is called  $S_\beta$ -closed set if the complement of  $B$  is  $S_\beta$ -open set.

The set of all semi-open (resp.  $\beta$ -open,  $S_\beta$ -open) sets is denoted by  $SO(X, \tau)$  (resp.  $\beta O(X, \tau)$ ,  $S_\beta O(X, \tau)$ ). The set of all semi-closed (resp.  $\beta$ -closed,  $S_\beta$ -closed) sets is denoted by  $SC(X, \tau)$  (resp.  $\beta C(X, \tau)$ ,  $S_\beta C(X, \tau)$ ). The semi closure, semi interior,  $\beta$ -closure,  $\beta$ -interior,  $S_\beta$ -closure, and  $S_\beta$ -interior of a set  $A$  are defined in similar way of closure and interior of a set  $A$ . The closure of  $G$  in  $(X, \tau)$  is denoted by  $cl(G)^\tau$ .

**Definition 2.2.** (4) A topological space  $(X, \tau)$  is said to be:

(i) *locally indiscrete* if every open subset of  $X$  is closed. (A set  $A \subseteq X$  is called clopen, if  $A$  are both open and closed.)

(ii) *regular* if given any point  $x \in X$  and any closed subset  $F \subseteq X - \{x\}$  implies that there exist open sets  $U$  and  $V$  such that  $x \in U$ ,  $F \subseteq V$  and  $U \cap V = \emptyset$ .

**Definition 2.3.** (6) Let  $(X, \tau)$  be a topological space. A subset  $N$  of  $X$  is called

$S_\beta$ -neighbourhood of subset  $A$  in  $X$  if, there exists a  $S_\beta$ -open set  $U$  such that  $A \subseteq U \subseteq N$ . We say that  $N$  is a  $S_\beta$ -neighbourhood of an element  $x \in X$  if  $A = \{x\}$ , the set of all  $S_\beta$ -neighbourhood of an element  $x$  denoted by  $S_\beta \eta(x)$ . The  $S_\beta$ -interior and  $S_\beta$ -closure of a set  $A$  denoted by  $S_\beta int(A) = \{x \in X \mid A \in S_\beta \eta(x)\}$  and  $S_\beta cl(A) = \{x \in X \mid A \cap W \neq \emptyset, \forall W \in S_\beta \eta(x)\}$  respectively.

**Example 2.1** (6) Let  $X = \{p, q, r, s\}$ ,

$\tau = \{\emptyset, \{p\}, \{q\}, \{p, q\}, \{p, q, r\}, X\}$  be topology on  $X$ . We get that  $S_\beta O(X) = \{\emptyset, \{q\}, \{p, r, s\}, X\}$ .

Consider  $A = \{r, s\}$ . Then  $S_\beta$ -neighbourhood of subset  $A$  are  $\{p, r, s\}$  and  $X$ ,  $S_\beta \eta(q) = \{\{q\}, X\}$ ,  $S_\beta int(A) = \emptyset$ , and  $S_\beta cl(A) = \{p, r, s\}$ .

Some properties of  $S_\beta$ -interior and  $S_\beta$ -closure sets are mentioned in the following result. For the routine proofs are omitted.

**Theorem 2.1.** (6) For any subset  $A$  and  $B$  of a topological space  $X$ . The following statements are holds:

(i) The  $S_\beta$ -interior of  $A$  is the union of all  $S_\beta$ -open sets contained in  $A$ .

(ii)  $S_\beta int(A)$  is the largest  $S_\beta$ -open set contained in  $A$ .

(iii) A set  $A$  is  $S_\beta$ -open set if and only if  $A = S_\beta int(A)$ .

(iv)  $S_\beta int(\emptyset) = \emptyset$  and  $S_\beta int(X) = X$ .

(v)  $S_\beta int(A) \subseteq A$ .

(vi) If  $A \subseteq B$ , then  $S_\beta int(A) \subseteq S_\beta int(B)$ .

(vii)  $S_\beta \text{int}(A) \cup S_\beta \text{int}(B)$   
 $\subseteq S_\beta \text{int}(A \cup B)$ .

(viii)  $S_\beta \text{int}(A \cap B)$   
 $\subseteq S_\beta \text{int}(A) \cap S_\beta \text{int}(B)$ .

(ix)  $S_\beta \text{int}(A \setminus B)$   
 $\subseteq S_\beta \text{int}(A) \setminus S_\beta \text{int}(B)$ .

Here we give some properties of  $S_\beta$ -closure of a set similar forms of closure of a set.

**Theorem 2.2.** (6) For any subset  $F$  and  $E$  of a topological space  $X$ . The following statements are true:

- (i) The  $S_\beta$ -closure of  $F$  is the intersection of all  $S_\beta$ -closed sets in  $X$  containing  $F$ .
- (ii)  $S_\beta cl(F)$  is the smallest  $S_\beta$ -closed set containing  $F$ .
- (iii)  $F$  is  $S_\beta$ -closed set if and only if  $F = S_\beta cl(F)$ .
- (iv)  $S_\beta cl(\emptyset) = \emptyset$  and  $S_\beta cl(X) = X$ .
- (v)  $F \subseteq S_\beta cl(F)$ .
- (vi) If  $F \subseteq E$ , then  $S_\beta cl(F) \subseteq S_\beta cl(E)$ .
- (vii)  $S_\beta cl(F) \cup S_\beta cl(E)$   
 $\subseteq S_\beta cl(F \cup E)$ .
- (viii)  $S_\beta cl(F \cap E)$   
 $\subseteq S_\beta cl(F) \cap S_\beta cl(E)$ .
- (ix)  $X \setminus S_\beta cl(F) = S_\beta \text{int}(X \setminus F)$ .
- (x)  $X \setminus S_\beta \text{int}(F) = S_\beta cl(X \setminus F)$ .
- (xi)  $S_\beta \text{int}(F) = X \setminus S_\beta cl(X \setminus F)$ .

**Definition 2.4.** (4, 6) Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a function,

- (i)  $f$  is *continuous* at  $x \in X$  if for each open subset  $V$  of  $f(x)$ , there exists an open set  $U$  of  $x$  such that  $f(U) \subseteq V$ .
- (ii)  $f$  is  $S_\beta$ -continuous at  $x \in X$  if for each open subset  $V$  of  $f(x)$ , there exists an  $S_\beta$ -open set  $U$  of  $x$  such that  $f(U) \subseteq V$ . If  $f$  is  $S_\beta$ -continuous at  $x$  for all  $x \in X$ , we say that  $f$  is  $S_\beta$ -continuous on  $X$ .

(iii)  $f$  is *open* if for each open subset  $U$  of  $X$ ,  $f(U) \in \sigma$ .

**Definition 2.5.** (1) Let  $(X, \tau)$  be a topological space.

(i) A topological space  $(X, \tau)$  is said to be *regular* if for any point  $x \in X$  and any closed subset  $F \subseteq X \setminus \{x\}$ , implies that there exist open sets  $U$  and  $V$ , such that  $x \in U$ , and  $U \cap V = \emptyset$ .

(ii) A space  $X$  has the *fixed point property* if, every continuous function  $f : X \rightarrow X$ , there exists an  $x \in X$  such that  $f(x) = x$ , briefly denoted by *FPP*.

**Theorem 2.3.** (6) Let  $(X, \tau)$  be a topological space and  $G \subseteq X$ . The following statements hold:

- (i) If  $G \in \tau$  then  $G \in SO(X, \tau)$ .
- (ii)  $G \in S_\beta O(X, \tau)$  if and only if  $G \in SO(X, \tau)$  and  $G = \cup \{B \subseteq G \mid B \in \beta O(X, \tau)\}$ .
- (iii) If  $X$  is a locally indiscrete space and  $G \in SO(X, \tau)$  ( $G \in SC(X, \tau)$ ) then  $G$  is closed (open) in  $X$ .
- (iv) If  $X$  is a locally indiscrete space and  $G \in SO(X, \tau)$  then  $G \in S_\beta O(X, \tau)$ .


**Theorem 2.4.** (6) Let  $(X, \tau)$  and  $(Y, \sigma)$  be a topological spaces, and  $f : X \rightarrow Y$ , following are equivalent:

- (i)  $f$  is  $S_\beta$ -continuous.
- (ii) For each open set  $B$  in  $Y$   $f^{-1}(B) \in S_\beta O(X, \tau)$ .
- (iii) For each closed set  $B$  in  $Y$ ,  $f^{-1}(B) \in S_\beta C(X, \tau)$ .
- (iv) For each subset  $A$  in  $X$ ,  $f(S_\beta cl(A)) \subseteq cl(f(A))$ .
- (v) For each subset  $A$  in  $X$ ,  $\text{int}(f(A)) \subseteq f(S_\beta \text{int}(A))$ .
- (vi) For each subset  $B$  in  $Y$ ,  $S_\beta cl(f^{-1}(B)) \subseteq f^{-1}(cl(B))$ .
- (vii) For each subset  $B$  in  $Y$ ,  $f^{-1}(\text{int}(B)) \subseteq S_\beta \text{int}(f^{-1}(cl(B)))$ .

**Theorem 2.5.** (6) Let  $(X, \tau)$  and  $(Y, \sigma)$  be topological spaces, and  $f: X \rightarrow Y, A \subseteq X$ . If  $A$  is a clopen set then the restriction of a function  $f$  is  $S_\beta$ -continuous, as  $f|_A: A \rightarrow Y$  is  $S_\beta$ -continuous.

The following theorem indicates that compositions of  $S_\beta$ -continuous functions are  $S_\beta$ -continuous, when a domain of the second function is locally indiscrete space.

**Theorem 2.6.** Let  $(X, \tau)$ ,  $(Y, \sigma)$  and  $(Z, \nu)$  be topological spaces, and  $Y$  is a locally indiscrete space, if  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  be the two  $S_\beta$ -continuous then the composite function  $g \circ f$  (see in Figure 1) is  $S_\beta$ -continuous.



**Figure 1** The composite function  $g \circ f$ .

**Proof.** Let  $G$  be an open set in  $Z$ . Since  $g$  is  $S_\beta$ -continuous, by theorem 2.4. (ii), so that  $g^{-1}(G) \in S_\beta O(Y, \sigma)$ . Moreover, since  $Y$  is a locally indiscrete space, by Theorem 2.3. (i, iv),  $g^{-1}(G)$  is open set in  $Y$ . For assumption  $f$  is  $S_\beta$ -continuous, we obtain  $f^{-1}(g^{-1}(G)) \in S_\beta O(X, \tau)$ .

But  $(g \circ f)^{-1}(G) = f^{-1}(g^{-1}(G))$ , therefore and hence  $g \circ f$  is  $S_\beta$ -continuous.  $\square$

### 3. Main Results

The purpose of this section is to explore properties and characterizations of  $S_\beta$ -continuous retract and  $S_\beta$ -continuous fixed point property. Cammarolo and Nori (2) defined a subset  $A$  of a space  $X$  to be a  $\delta$ -continuous

retract of  $X$ , defined a space  $X$  has the  $\delta$ -continuous fixed point property. We shall similarly defined a  $S_\beta$ -continuous retract and  $S_\beta$ -continuous fixed point property as follows.

**Definition 3.1.** Let  $(X, \tau)$  be a topological spaces.

(i) A subset  $A$  of space  $X$  is called a  $S_\beta$ -continuous retract of  $X$  if there exists a  $S_\beta$ -continuous  $f: X \rightarrow A$  such that  $f|_A$  is the identity on  $A$ , that is  $f(x) = x$  for every  $x \in A$ , and  $f$  is called a  $S_\beta$ -continuous retraction.

(ii) A space  $X$  is said to have the  $S_\beta$ -continuous fixed point property, briefly denoted by  $S_\beta cFPP$ , if for every  $S_\beta$ -continuous function  $f: X \rightarrow X$ , there exists an  $x \in X$  such that  $f(x) = x$ .

**Example 3.1.** Let  $X = \{p, q, r, s\}$  and let  $\tau = \{\emptyset, \{r\}, \{p, q\}, \{p, q, r\}, \{p, q, s\}, X\}$ ,  $\tau_A = \{\emptyset, \{p, q\}, A\}$ ,  $A = \{p, q, s\}$ , and  $f: (X, \tau) \rightarrow (A, \tau_A)$  defined that  $f(p) = p, f(q) = q, f(r) = r, f(s) = s$ . Since  $S_\beta O(X, \tau) = \{\emptyset, \{r\}, \{p, q\}, \{p, q, r\}, \{p, q, s\}, X\}$  and  $f$  is  $S_\beta$ -continuous. Hence  $A$  is a  $S_\beta$ -continuous retract of  $X$ .

**Example 3.2.** Let  $X = \{p, q, r\}$  and  $\tau = \{\emptyset, \{p\}, \{q, r\}, X\}$  be a topology for  $X$ . We show that  $X$  has the  $S_\beta$ -continuous fixed point property.

**Solution.** Now, we prove by contrapositive. Assume that, there exist a  $S_\beta$ -continuous function  $f: X \rightarrow X$  such that  $f(x) \neq x, \forall x \in X$ . Suppose that for  $p, q, r \in X$  such that  $f(p) = q, f(q) = r$ , and  $f(r) = q$ .

**Step1.** To show that,  $f$  is  $S_\beta$ -continuous. Because  $\emptyset, \{p\}, \{q, r\}$ , and  $X$  are open sets in  $X$ . By definition of  $f$ , we have,  $f^{-1}(\emptyset) = \emptyset, f^{-1}(p) = \emptyset$ , and  $f^{-1}(\{q, r\}) = X$ , such that  $\emptyset, X \in S_\beta O(X)$ . By Theorem 2.4. (ii), Therefore  $f$  is a  $S_\beta$ -continuous.

**Step2.** To show that,  $f(x) \neq x, \forall x \in X$ . By defined function of  $f$   $f(p)=q, f(q)=r$  and  $f(r)=q$ , such that  $p \neq q$  and  $q \neq r$ . It follows that,  $f(x) \neq x, \forall x \in X$ . Form the both steps, it shows that,  $X$  has the  $S_\beta$ -continuous fixed point property.

Next, we investigate some properties of  $S_\beta$ -continuous retract of a space  $X$ .

**Lemma 3.1.** Let  $(X, \tau)$  be a topological space,  $A \subseteq X$  with a locally indiscrete subspace  $(A, \tau_A)$ . Then  $A$  is a  $S_\beta$ -continuous retract of  $X$  if and only if for every topological space  $(Y, \sigma)$  and for each  $S_\beta$ -continuous function  $f: A \rightarrow Y$  can be extend to a  $S_\beta$ -continuous over  $X$  relative to  $Y$ .

**Proof.** ( $\Rightarrow$ ) Suppose that  $(X, \tau)$  is a topological space and  $A \subseteq X$  is locally indiscrete subspace  $(A, \tau_A)$ . Let  $f: A \rightarrow Y$  be a  $S_\beta$ -continuous function. Since  $A$  is a  $S_\beta$ -continuous retract of  $X$ . By Definition 3.1. (i), we have  $g: X \rightarrow A$  is a  $S_\beta$ -continuous function.


By Theorem 2.6. and property of locally indiscrete subspace  $(A, \tau_A)$ , therefore  $f \circ g: X \rightarrow Y$  is a  $S_\beta$ -continuous function and implies  $f \circ g(x) = f(g(x)) = f(x)$ , for every  $x \in A$ . Hence  $f \circ g$  is an extension of  $f$ .

( $\Leftarrow$ ) Suppose that  $i_A: A \rightarrow A$  is an identity function on  $A$ . Note that  $i_A$  is also a  $S_\beta$ -continuity. By assumption, we have a  $S_\beta$ -continuous function  $g: X \rightarrow A$  where  $g|_A = i_A$ . Therefore  $A$  is a  $S_\beta$ -continuous retract of  $X$ .  $\square$

A  $S_\beta$ -continuous retract of a space  $X$  with a clopen set is necessity for the  $S_\beta$ -continuous fixed point property, as following.

**Theorem 3.1.** Let  $(X, \tau)$  be a topological space,  $A$  be a clopen subset in  $X$ . If  $X$  has the  $S_\beta$ -continuous fixed point property and  $A$  is a  $S_\beta$ -

continuous retract of  $X$ , then  $A$  has the  $S_\beta$ -continuous fixed point property ( $S_\beta$ -cFPP).



**Figure 2** To show functions  $f$ ,  $F$ ,  $j$ , and  $j \circ F$ .

**Proof.** Suppose that  $f: A \rightarrow A$  is a  $S_\beta$ -continuous function, and  $A$  is a clopen (both open and closed) set in  $X$ . Then  $(A, \tau_A)$  is a locally indiscrete subspace. Since  $A$  is a  $S_\beta$ -continuous retract of  $X$ , by Lemma 3.1., we have a  $S_\beta$ -continuous function  $F: X \rightarrow A$  which it is an extension of  $f$ .


Let  $j: A \rightarrow X$  be an identity function, as  $f(x) = x$ , for all  $x \in A$ . To show that  $j \circ F$  is a  $S_\beta$ -continuous function (see in Figure 2). Let  $U$  be an open in  $X$ , hence  $j^{-1}(U) = U \cap A$  is an open in  $(A, \tau_A)$ . Since  $F: X \rightarrow A$  is a  $S_\beta$ -continuous function, implies that

$F^{-1}(j^{-1}(U)) = (j \circ F)^{-1}(U) \in S_\beta O(X)$ , and therefore  $(j \circ F): X \rightarrow X$  is a  $S_\beta$ -continuous function. Consequently to show that  $A$  has the  $S_\beta$ -continuous fixed point property. Since  $X$  has the  $S_\beta$ -continuous fixed point property. There exists an  $x \in A \subseteq X$  such that  $x = (j \circ F)(x) = j(F(x)) = j(f(x)) = f(x)$ . So, it completes the proof.  $\square$

The following theorem is a slight modification of theorem 1 in (3), Theorem 3.2. in (2), and Theorem 4.11. in (9).

**Theorem 3.2** Let  $(X, \tau)$  and  $(X, \sigma)$  be topological spaces where  $(X, \tau)$  is a locally indiscrete space, and it has the  $S_\beta$ -continuous fixed point property. If  $\sigma$  stronger than  $\tau$

(as  $\sigma \supseteq \tau$ ) and  $cl(G)^{(\tau)} = cl(G)^{(\sigma)}$ , for any  $G \in \sigma$  then  $(X, \sigma)$  has the continuous fixed point property.



**Figure 3** Diagram of functions  $f$ ,  $g$ ,  $h$ , and  $i$ .

**Proof.** We must show that,  $(X, \sigma)$  has the continuous fixed point property. Let  $f : (X, \sigma) \rightarrow (X, \sigma)$  be a continuous function. We must demonstrate  $f(x) = x, \forall x \in X$ . Defined the three functions,  $g, h$ , and  $i$  which  $g : (X, \sigma) \rightarrow (X, \tau)$ ,  $h : (X, \tau) \rightarrow (X, \tau)$  are defined by setting  $g(x) = h(x) = f(x)$ , for all  $x \in X$ , and  $i : (X, \tau) \rightarrow (X, \sigma)$  is an identity function. For every  $x \in X$ , we have  $f(x) = g(x) = i(g(x)) = i \circ g(x)$ , (see in Figure 3) so that  $f = i \circ g$  is a continuous function. To show that  $g$  is a continuous function. Let  $G \in \tau$ .

By identity function property,  $i$  is an open, one to one and onto. Therefore  $i(G) \in \sigma$ . Since  $i \circ g$  is continuity, we have  $(i \circ g)^{-1}(i(G)) \in \sigma$ . By the property of one to one, onto, and inverse function. We have

$$\begin{aligned} (i \circ g)^{-1}(i(G)) &= (g^{-1} \circ i^{-1})(i(G)) \\ &= g^{-1}(i^{-1}(i(G))) \\ &= g^{-1}(G). \end{aligned}$$

So that  $g^{-1}(G) \in \sigma$ , therefore  $g$  is a continuous function. To show that  $h$  is a  $S_\beta$ -continuous function. Let  $x \in X$  and  $h(x) \in V$ , for all  $V \in \tau$ . Since  $(X, \tau)$  is a regular spaces, there exists an  $A \in \tau$  such that  $h(x) \in A \subseteq cl(A)^{(\tau)} \subseteq V$ . Since  $g$  is a continuous function,  $h(x) \in A \in \tau$ , and the setting  $h, g$ , and  $f$ , we have  $g^{-1}(A) \in \sigma$ , and  $h^{-1}(A) = f^{-1}(A) = g^{-1}(A)$ . So that  $h^{-1}(A) = f^{-1}(A) \in \sigma$ . By closure properties and our assumption  $cl(h^{-1}(A))^{(\tau)} = cl(h^{-1}(A))^{(\sigma)}$ ,

$$\begin{aligned} x \in h^{-1}(A) &\subseteq cl(h^{-1}(A))^{(\tau)} \\ &= cl(h^{-1}(A))^{(\sigma)} = cl(f^{-1}(A))^{(\sigma)}. \end{aligned} \quad (3.1)$$

Since  $f$  is a continuous function and

$$cl(A)^{(\tau)} \subseteq V, \text{ so that}$$

$$\begin{aligned} cl(f^{-1}(A))^{(\sigma)} &\subseteq f^{-1}(cl(A)^{(\sigma)}) = f^{-1}(cl(A)^{(\tau)}), \\ \text{and } f^{-1}(cl(A)^{(\sigma)}) &\subseteq f^{-1}(V). \end{aligned} \quad (3.2)$$

By (3.1) and (3.2), we obtain  $cl(h^{-1}(A))^{(\tau)} \subseteq f^{-1}(V)$ . From defined  $h$ ,  $h^{-1}(V) = f^{-1}(V)$ , implies that  $cl(h^{-1}(A))^{(\tau)} \subseteq h^{-1}(V)$ .  $\square$

Since  $i$  is an open function with  $h^{-1}(A) \in \sigma$ , and  $\tau \subseteq \sigma$ , we have  $h^{-1}(A)$  is an open in the space  $(X, \tau)$ . Let  $U = h^{-1}(A)$ . Since  $(X, \tau)$  is a locally indiscrete space, implies that  $U$  be the both open and closed, therefore  $U$  is a semi open set in the space  $(X, \tau)$ . By Theorem 2.3. (iv),  $U$  is a  $S_\beta$ -open set in the space  $(X, \tau)$ . which  $x \in U$ . By (3.1) and (3.3), we have  $U \subseteq h^{-1}(V)$ , so that  $h(U) \subseteq h(h^{-1}(V)) \subseteq V$ . Hence  $h$  is a  $S_\beta$ -continuous function, by Definition 2.5. Finality to show that  $(X, \sigma)$  has the continuous fixed point property. By the assumption,  $(X, \tau)$  has the  $S_\beta$ -continuous fixed point property. So there exists an  $x \in X$  such that  $h(x) = x$ . But we have  $f(x) = h(x)$ , for all  $x \in X$ , implies that  $f(x) = x$ , so we have the result.  $\square$

Next, Lemma is to explore properties of an one to one, onto, open functions and the composition which it is  $S_\beta$ -continuous function.

**Lemma 3.2.** Let  $(X, \tau), (Y, \sigma)$ , and  $(Z, \nu)$  be topological spaces, and  $g : (Y, \sigma) \rightarrow (Z, \nu)$  be an one to one, onto, and open. If  $g \circ f : (X, \tau) \rightarrow (Z, \nu)$  is a  $S_\beta$ -continuous function then  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a  $S_\beta$ -continuous function.

**Proof.** Let  $V$  be an open set in space  $(Y, \sigma)$ . Since  $g$  is an open function, implies that  $g(V)$  is an open set in space  $(Z, \nu)$ . by our assumption,  $g \circ f$  is a  $S_\beta$ -continuous function, so that  $(g \circ f)^{-1}(g(V)) \in S_\beta(X, \tau)$ .

By property of inverse function, one to one, and onto function, we have

$$\begin{aligned}(g \circ f)^{-1}(g(V)) &= (f^{-1} \circ g^{-1})(g(V)) \\ &= f^{-1}(g^{-1}(g(V))) \\ &= f^{-1}(V) \in S_\beta(X, \tau).\end{aligned}$$

And by Theorem 2.4. (ii), hence  $f$  is a  $S_\beta$ -continuous function.  $\square$

In the following example, we show that the converse of the Lemma 3.2 is not true. Recall a topological space  $(X, \tau)$  of Example 3.2 in (5).

**Example 3.3.** Let  $(X, \tau), (Y, \sigma)$ , and  $(Z, \nu)$  be topological spaces, such that  $X = \{a, b, c, d\}$ ,  $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$ ,  $Y = \{p, q, r\}$ ,  $\sigma = \{\emptyset, Y\}$ ,  $Z = \{l, m, n\}$ ,  $\nu = \{\emptyset, Z, \{m\}, \{n\}, \{m, n\}\}$ . Function  $f: (X, \tau) \rightarrow (Y, \sigma)$  defined that  $f(a) = p, f(b) = q, f(c) = r = f(d)$ , and function  $g: (Y, \sigma) \rightarrow (Z, \nu)$  defined that  $g(p) = l, g(q) = m$ , and  $g(r) = n$ . Then  $g$  is a one to one, onto, and open. Since  $S_\beta O(X, \tau) = \{\emptyset, \{b\}, \{a, c, d\}, X\}$ , and

$$\begin{aligned}f^{-1}(\emptyset) &= \emptyset \in S_\beta O(X, \tau), \text{ and} \\ f^{-1}(Y) &= X \in S_\beta O(X, \tau).\end{aligned}$$

So that  $f$  is  $S_\beta$ -continuous. Because

$$\begin{aligned}g \circ f(a) &= l, g \circ f(b) = m, g \circ f(c) = n, \\ g \circ f(d) &= n, \text{ and}\end{aligned}$$

$$(g \circ f)^{-1}(\{m, n\}) = \{b, c, d\} \notin S_\beta O(X, \tau).$$

Therefore,  $g \circ f$  is not  $S_\beta$ -continuous.

In Theorem 3.2., we replace  $cl(G)^{(\tau)} = cl(G)^{(\sigma)}$  with  $S_\beta cl(G)^{(\tau)} = S_\beta cl(G)^{(\sigma)}$ . We have the following Theorem.

**Theorem 3.3.** Let  $(X, \tau)$  and  $(X, \sigma)$  be topological spaces with regular space  $(X, \tau)$  has the continuous fixed point property. If  $\sigma$  stronger than  $\tau$  and  $S_\beta cl(G)^{(\tau)} = S_\beta cl(G)^{(\sigma)}$  where  $G \in \sigma$ , for all  $G \in S_\beta O(X, \sigma)$  then  $(X, \sigma)$  has the  $S_\beta$ -continuous fixed point property.

**Proof.** Suppose that  $f: (X, \sigma) \rightarrow (X, \sigma)$  is a  $S_\beta$ -continuous, we need to establish that

$f(x) = x$ ,  $\forall x \in X$ . The three functions were defined,  $g: (X, \sigma) \rightarrow (X, \tau)$ ,  $h: (X, \tau) \rightarrow (X, \tau)$ , and  $i: (X, \tau) \rightarrow (X, \sigma)$ , as  $g(x) = h(x) = f(x)$ , for all  $x \in X$ , and  $i$  be an identity. It follows that  $f = i \circ g$  is a  $S_\beta$ -continuous function.

To show that  $g$  is a  $S_\beta$  continuous function. Let  $G \in \tau$ . The identity function imply open and bijection, we get  $i(G) \in \sigma$ . Since  $i \circ g$  is a  $S_\beta$ -continuous, by Lemma 3.2, we have  $g$  is a  $S_\beta$ -continuous function. To show that  $h$  is a continuous function. Let  $x \in X$  and  $h(x) \in V$ , for all open set  $V$  in space  $(X, \tau)$ . Since  $(X, \tau)$  is a regular space, there exists an open subset  $A$  in space  $(X, \tau)$  such that  $h(x) \in A \subseteq cl(A)^{(\tau)} \subseteq V$ . By the  $S_\beta$ -continuous function  $g$ ,  $h(x) \in A$ , and the setting  $h, g$ , and  $f$ , we have  $g^{-1}(A) \in S_\beta O(X, \sigma)$ , and  $h^{-1}(A) = f^{-1}(A) = g^{-1}(A)$ , implies that  $h^{-1}(A) = f^{-1}(A) \in S_\beta O(X, \sigma)$ . Utilize  $S_\beta$ -closure,  $S_\beta$ -interior properties, and our assumption, we have  $S_\beta cl(h^{-1}(A))^{(\tau)} = S_\beta cl(h^{-1}(A))^{(\sigma)}$ , implies that

$$\begin{aligned}x \in h^{-1}(A) &= S_\beta int(h^{-1}(A))^{(\sigma)} \\ &\subseteq S_\beta int(S_\beta cl(h^{-1}(A))^{(\sigma)})^{(\sigma)} \\ &\subseteq S_\beta cl(h^{-1}(A))^{(\sigma)} \\ &= S_\beta cl(h^{-1}(A))^{(\tau)} \\ &= S_\beta cl(f^{-1}(A))^{(\sigma)} = S_\beta cl(f^{-1}(A))^{(\tau)}. \quad (3.4)\end{aligned}$$

Since  $\tau \subseteq \sigma$ , we have  $cl(A)^{(\sigma)} \subseteq cl(A)^{(\tau)}$ . Since  $f$  is a  $S_\beta$ -continuous function and  $cl(A)^{(\tau)} \subseteq V$ , we obtain that

$$\begin{aligned}S_\beta cl(f^{-1}(A))^{(\sigma)} &\subseteq f^{-1}(cl(A)^{(\sigma)}) \\ &\subseteq f^{-1}(cl(A)^{(\sigma)}) \\ &\subseteq f^{-1}(V).\end{aligned} \quad (3.5)$$

By the (3.4) and (3.5),  $h^{-1}(A) \subseteq f^{-1}(V)$ .

Since  $h(x) = f(x)$ , for every  $x \in X$ .

So that  $h^{-1}(V) = f^{-1}(V)$ , we get  $h^{-1}(A) \subseteq h^{-1}(V)$ .

Since  $h^{-1}(A) = g^{-1}(A) \in \sigma$ ,  $\tau \subseteq \sigma$ , and using identity function, we have  $i(h^{-1}(A)) = h^{-1}(A)$ , so that  $h^{-1}(A)$  is an open set in space  $(X, \tau)$  and  $x \in h^{-1}(A)$ . Let  $U = h^{-1}(A)$ , implies that  $h(U) = h(h^{-1}(A)) \subseteq h(h^{-1}(V)) \subseteq V$ . Hence  $h$  is a continuous function on  $(X, \tau)$ . Since  $(X, \tau)$  has the continuous fixed point property, so there

exists an  $x \in X$  with  $h(x) = x$ , but  $h(x) = f(x)$ , for every  $x \in X$ , there exists an  $x \in X$  with  $f(x) = x$ , hence  $(X, \sigma)$  has the  $S_\beta$ -continuous fixed point property.  $\square$

#### 4. Conclusion

From the above study it is clear that for regular and locally indiscrete space  $(X, \tau)$ . If  $(X, \tau)$  has the  $S_\beta$ -continuous fixed point property (the continuous fixed point property),  $\sigma$  is a topology for  $X$  stronger than  $\tau$  and  $cl(G)^{(\tau)} = cl(G)^{(\sigma)}$  ( $S_\beta cl(G)^{(\tau)} = S_\beta cl(G)^{(\sigma)}$ ) for every  $G \in \sigma$ , then  $(X, \sigma)$  has the fixed point property (the  $S_\beta$ -continuous fixed point property). The results from this paper are based for further studies on the concept of other types of weak the continuous fixed point property. In the future we will study on minimal  $S_\beta$ -open sets and the continuous fixed point property in generalized topological spaces.

#### Acknowledgements

The author would like to thank the member of Mathematics program in Chaiyaphum Rajabhat University for his/her valuable comments and would like to thank Assoc. Prof. Dr. Narongrit Puturong for his useful suggestions and the reviewers for their valuable comments and suggestions. The author would like to thank Phetchabun Rajabhat University for the full financial support in this research.

#### Declaration of conflicting interests

The authors declared that they have no conflicts of interest in the research, authorship, and this article's publication.

#### References

1. Adams CC, Franzosa RD. Introduction to topology pure and applied. New Jersey: Upper saddle river; 2008.
2. Cammarolo F, Noiri T. On the  $\delta$ -continuous fixed point property. Int. J. Math. Math. Sci. 1990;13(1):45-50.
3. Connell EH. Property of fixed point spaces. Proc. Am. Math. Soc. 1959;10(3):974-979.
4. Dugundji J. Topology. Boston: Allyn and Bacon; 1966.
5. Jongrak A. Some properties of  $S_\beta$ -open Mappings. The proceedings of annual meeting in mathematics. 2017;22:TPO031-035.
6. Khalaf AB, Ahmed NK.  $S_\beta$ -open sets and  $S_\beta$ -continuity in topological spaces. Thai J. Math. 2013;11(2):319-335.
7. Levine N. Semi-open sets and semi-continuity in topological spaces. Am. Math. Mon. 1963;29(70):36-4.
8. Monsef ME, Deeb SN, Mahmood RA.  $\beta$ -open sets and  $\beta$ -continuous mappings. Bullentin of the Faculty of Science, Assiut University. 1983;12(1):77-90.
9. Puturong N. On strongly  $\theta$ -semi-continuous functions. Thai J. Math. 2007; 5(3):11-23.