

Source details

Caryologia

Open Access ⓘ

Scopus coverage years: from 1949 to 2023

Publisher: Firenze University Press

ISSN: 0008-7114 E-ISSN: 2165-5391

Subject area:

Agricultural and Biological Sciences: General Agricultural and Biological Sciences

Biochemistry, Genetics and Molecular Biology: Genetics

Source type: Journal

View all documents >

Set document alert

 Save to source list

CiteScore 2022

1.6 ⓘ

SJR 2022

0.293 ⓘ

SNIP 2022

0.425 ⓘ

CiteScore CiteScore rank & trend Scopus content coverage

i

Improved CiteScore methodology

×

CiteScore 2022 counts the citations received in 2019-2022 to articles, reviews, conference papers, book chapters and data papers published in 2019-2022, and divides this by the number of publications published in 2019-2022. [Learn more >](#)

CiteScore 2022

1.6 = $\frac{323 \text{ Citations 2019 - 2022}}{201 \text{ Documents 2019 - 2022}}$

Calculated on 05 May, 2023

CiteScoreTracker 2023 ⓘ

1.6 = $\frac{334 \text{ Citations to date}}{210 \text{ Documents to date}}$

Last updated on 05 April, 2024 • Updated monthly

CiteScore rank 2022 ⓘ

Category	Rank	Percentile
Agricultural and Biological Sciences	#104/213	51st
General Agricultural and Biological Sciences		
Biochemistry, Genetics and Molecular Biology	#283/328	13th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

- What is Scopus
- Content coverage
- Scopus blog
- Scopus API
- Privacy matters

Language

- 日本語版を表示する
- 查看简体中文版本
- 查看繁體中文版本
- Просмотр версии на русском языке

Customer Service

- Help
- Tutorials
- Contact us

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

All content on this site: Copyright © 2024 Elsevier B.V. ↗, its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the Creative Commons licensing terms apply. We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.



Caryologia








COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
United Kingdom Universities and research institutions in United Kingdom Media Ranking in United Kingdom	Agricultural and Biological Sciences Agricultural and Biological Sciences (miscellaneous) Biochemistry, Genetics and Molecular Biology Genetics	Firenze University Press Universita degli Studi di Firenze in Scimago Institutions Rankings	33
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	00087114, 21655391	1949-2023	Homepage How to publish in this journal caryo@unifi.it
SCOPE			
Caryologia is devoted to the publication of original papers, and occasionally of reviews, about plant, animal and human karyological, cytological, cytogenetic, embryological and ultrastructural studies. Articles about the structure, the organization and the biological events relating to DNA and chromatin organization in eukaryotic cells are considered. Caryologia has a strong tradition in plant and animal cytosystematics and in cytotoxicology. Bioinformatics articles may be considered, but only if they have an emphasis on the relationship between the nucleus and cytoplasm and/or the structural organization of the eukaryotic cell.			

Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS

1 Plant Systematics and Evolution AUT 32% similarity	2 Turkish Journal of Botany TUR 32% similarity	3 Acta Botanica Croatica HRV 29% similarity	4 Korean Journal of Plant Taxonomy KOR 29% similarity
---	---	--	--



Caryologia

Q3 Agricultural and Biological Sciences (miscellaneous) best quartile

SJR 2023 0.18

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scima
```

SCImago Graphica


Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of March 2024

[HOME](#) / [ARCHIVES](#) / Vol. 76 No. 4 (2023)

Vol. 76 No. 4 (2023)

**PUBLISHED:** 2024-03-15

FULL ISSUE

 **PDF**

ARTICLES

First cytogenetic study of the Somphong's rasbora (*Trigonostigma somphongsi*) (Perciformes, Cyprinidae), a critically endangered species in Thailand

Surachest Aiumsumang, Chavalit Vidthayanon, Sitthi Kulabtong, Alongklod Tanomtong, Sumalee Phimphan
3-8

 **PDF**

Evaluation of the evolutionary process within *Populus caspica* species from Hyrcanian forests by karyotype analysis

Fereshteh Asadi-Corom, Farhad Asadi, Hossein Mirzaie-Nodoushan

9-14

**Chromosomal and genome size variations in Opium poppy (*Papaver somniferum* L.) from Afghanistan**

Sayed Zia Rasekh, Ghasem Karimzadeh

15-22

**Karyological analyses in several Algerian populations of six species of the genus *Vicia* L. (Fabaceae)**

Zahia Sebkhi, Rachida Issolah, Nabila Melzi, Hassina Benmouhoub, Mohamed Mefti

23-37

**Cytogenotoxic and antimicrobial effects of *Nezara viridula* (L.) (Hemiptera: Heteroptera: Pentatomidae) alcoholic extracts**

Nicoleta Anca Șuțan, Mircea Bărbuceanu, Daniela Barbuceanu, Ionica Deliu

39-49

**Apogamous *Isoetes coromandelina* L.f. (Isoetaceae) with asynaptic meiosis**

V. Irudayaraj, A. Benniamin, S. Arokia Raj

51-57

**Evaluation of the antigenotoxic potential of fresh bovine whey in onion meristematic roots exposed to Quizalofop-P-tefuryl**

Florica Colă, Elena Bonciu, Mugurel Colă, Nicoleta Anca Șuțan

59-66



[MAKE A SUBMISSION](#)

INFORMATION

[For Readers](#)[For Authors](#)[For Librarians](#)

Caryologia: International Journal of Cytology, Cytosystematics and Cytogenetics | E-ISSN: 2165-5391 | contact: alessio.papini@unifi.it

Published by [Firenze University Press](#) | [University of Florence](#) | Via Cittadella, 7 - 50144 Firenze (FI) - Italy | contact: info@fupress.com |

[Privacy policy](#) | [Terms of use](#) | [Legal notice](#) | [Accessibility Statement](#) | [Login/Registration](#) | [Announcements](#) | [Fupress Journals](#) | [Open Access Policy of the University of Florence](#) |

The *Caryologia* site and its metadata are licensed under a [Creative Commons Attribution 4.0 International License \(CC-BY- 4.0\)](#).



Platform &
workflow by
OJS / PKP

HOME / Editorial Team

Editorial Team

Editor in Chief

Alessio Papini - *Department of Biology*
University of Florence
Via La Pira, 4 – 0121 Firenze, Italy

Associate Editors

Alfonso Carabez-Trejo - *Mexico City, Mexico*
Katsuhiko Kondo - *Hagishi-Hiroshima, Japan*
Canio G. Vosa - *Pisa, Italy*

Subject Editors

Mycology

Renato Benesperi - *University of Florence, Italy*

Human and Animal Cytogenetics

Michael Schmid - *University of Würzburg, Germany*

Plant Cytogenetics

Lorenzo Peruzzi – *University of Pisa, Italy*

Cytotoxicity

Cleiton Pereira de Souza - *São Paulo, Brazil*

Plant Karyology and Phylogeny

Andrea Coppi – *University of Florence, Italy*

Histology and Cell Biology

Alessio Papini – *University of Florence, Italy*

Nanoparticles

Sara Falsini - *University of Florence, Italy*

Zoology

Mauro Mandrioli - *University of Modena and Reggio Emilia, Italy*

Genotoxicity

Nicoleta Anca Sutan - *University of Pitesti, Romania*

Editorial Assistant

[Sara Falsini](#) - *University of Florence, Italy*

Editorial Advisory Board

Andrew Bajer - *University of Oregon, Eugene, USA*
 Erika Bellini - *University of Pisa, Pisa, Italy*
 Andrea Bennici - *University of Florence, Florence, Italy*
 Graziella Berta - *Amedeo Avogadro University of Eastern Piedmont, Alessandria, Italy*
 Elisabetta Bianchi - *University of Florence, Florence, Italy*
 Davide Bizzaro - *Polytechnic University of Marche, Ancona, Italy*
 Elena Bonciu - *University of Craiova, Romania*
 Antonio Brito Da Cunha - *University of Sao Paulo, Sao Paulo, Brazil*
 Ernesto Capanna - *University "La Sapienza", Rome, Italy*
 Duccio Cavalieri - *Edmund Mach Fondation, San Michele all'Adige, Italy*
 Roberto Cremonini - *University of Pisa, Pisa, Italy*
 Mauro Cresti - *University of Siena, Siena, Italy*
 Giovanni Cristofolini - *University of Bologna, Bologna, Italy*
 Paolo Crosti - *University of Milano-Bicocca, Milan, Italy*
 Giovanni Delfino - *University of Florence, Firenze, Italy*
 Saverio D'Emérico - *University of Bari, Bari, Italy*
 Fabio Garbari - *University of Pisa, Pisa, Italy*
 Antonio Giovino - *Council for Agricultural Research and Economics, Bagheria, Italy*
 Claudia Giuliani - *University of studies of Milan, Milan, Italy*
 Marcelo Guerra - *Federal University of Pernambuco, Recife, Brazil*
 Waheeb K. Heneen - *The Swedish University of Agricultural Sciences, Svalöf, Sweden*
 Alessio Iannucci - *University of Florence, Florence, Italy*
 Leopoldo Iannuzzi - *CNR-ISPAA, Napoli, Italy*
 Janusz Limon - *Medical University of Gdansk, Gdansk, Poland*
 Jiangtao Liu - *University of Lanzhou, Lanzhou, China*
 Nils Mandahl - *Lund University, Lund, Sweden*
 Mauro Mandrioli - *University of Modena and Reggio Emilia, Modena, Italy*
 Gian Carlo Manicardi - *University of Modena and Reggio Emilia, Modena, Italy*
 Palmer Marchi - *University "La Sapienza", Rome, Italy*
 Fábio Porto-Foresti - *Universidade Estadual Paulista, São Paulo, Brazil*
 Monica Ruffini Castiglione - *University of Pisa, Pisa, Italy*
 Anca Sutan - *University of Pitesti, Romania*
 Luigi Sanità di Toppi - *University of Pisa, Pisa, Italy*
 Claus Steinlein - *University of Würzburg, Würzburg, Germany*
 Joan Vallès - *University of Barcelona, Barcelona, Catalonia, Spain*
 Qiner Yang - *Beijing Institute of Botany, Beijing, China*

MAKE A SUBMISSION

INFORMATION

For Readers

For Authors

For Librarians

Caryologia: International Journal of Cytology, Cytosystematics and Cytogenetics | E-ISSN: **2165-5391** | contact: alessio.papini@unifi.it

Published by [Firenze University Press](#) | [University of Florence](#) | Via Cittadella, 7 - 50144 Firenze (FI) - Italy | contact: info@fupress.com |

[Privacy policy](#) | [Terms of use](#) | [Legal notice](#) | [Accessibility Statement](#) | [Login/Registration](#) | [Announcements](#) | [Fupress Journals](#) | [Open Access Policy of the University of Florence](#) |

The *Caryologia* site and its metadata are licensed under a [Creative Commons Attribution 4.0 International License \(CC-BY- 4.0\)](#)



Platform &
workflow by
OJS / PKP

Journal Impact (/)

Q

Enter journal title, issn or abbr in this box to search

Caryologia

Journal Abbreviation: CARYOLOGIA
Journal ISSN: 0008-7114

About Caryologia

CARYOLOGIA is published by the Department of Plant Biology of the University of Florence; it is a periodical devoted to the publication of original researches and, occasionally, of reviews in plant and animai cytology, cytosystematics, cytogenetics, embriology and ultra-structure. Usually one volume is issued yearly and it consists of four numbers.

Year	Impact Factor (IF)	Total Articles	Total Cites
2022 (2023 update)	2.1	-	1189
2021	0.690	-	1180
2020	0.918	31	953
2019	0.621	39	740
2018	1.174	58	1007
2017	0.608	45	896
2016	0.516	47	808
2015	0.494	51	797
2014	0.740	42	754
2013	0.849	47	810
2012	0.632	49	888

2011	0.533	29	541
2010	0.398	36	592

You may also be interested in the following journals

- ▶ PLoS One (/journal/PLOS-ONE) ▶ Irbm (/journal/IRBM) ▶ Acta Cytologica (/journal/ACTA-CYTOL)
- ▶ Cytojournal (/journal/CYTOJOURNAL) ▶ Pharmaceutical Biology (/journal/PHARM-BIOL)
- ▶ Nature Cell Biology (/journal/NAT-CELL-BIOL) ▶ Crop Science (/journal/CROP-SCI)
- ▶ Cytogenetic and Genome Research (/journal/CYTOGENET-GENOME-RES)
- ▶ Analytical and Quantitative Cytology and Histology (/journal/ANAL-QUANT-CYTOL)
- ▶ Zoology in The Middle East (/journal/ZOOL-MIDDLE-EAST)

<< Carpathian Journal of Mathematics (/journal/CARPATHIAN-J-MATH)

Castanea (/journal/CASTANEA) >>

Top Journals in biology (/subject/biology)

- Nature Reviews Genetics (/journal/NAT-REV-GENET)
- Nature Reviews Microbiology (/journal/NAT-REV-MICROBIOL)
- Cell (/journal/CELL)
- Nature Genetics (/journal/NAT-GENET)
- Nature Methods (/journal/NAT-METHODS)
- Cell Stem Cell (/journal/CELL-STEM-CELL)
- Cell Metabolism (/journal/CELL-METAB)
- Annual Review of Biochemistry (/journal/ANNU-REV-BIOCHEM)

Nature Cell Biology (/journal/NAT-CELL-BIOL)
Annual Review of Plant Biology (/journal/ANNU-REV-PLANT-BIOL)
Trends in Cell Biology (/journal/TRENDS-CELL-BIOL)
Cell Host & Microbe (/journal/CELL-HOST-MICROBE)

Journal Impact (/)



Citation: Aiumsumang, S., Vidthayanon, C., Kulabong, S., Tanomtong, A., & Phimphan, S. (2023). First cytogenetic study of the Somphong's rasbora (*Trigonostigma somphongsi*) (Perciformes, Cyprinidae), a critically endangered species in Thailand. *Caryologia* 76(4):3-8. doi:10.36253/caryologia-2439

Received: December 30, 2023

Accepted: February 25, 2024

Published: March 14, 2024

Copyright: ©2023 Aiumsumang, S., Vidthayanon, C., Kulabong, S., Tanomtong, A., & Phimphan, S. This is an open access, peer-reviewed article published by Firenze University Press (<http://www.fupress.com/caryologia>) and distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Competing Interests: The Author(s) declare(s) no conflict of interest.

First cytogenetic study of the Somphong's rasbora (*Trigonostigma somphongsi*) (Perciformes, Cyprinidae), a critically endangered species in Thailand

SURACHEST AIUMSUMANG¹, CHAVALIT VIDTHAYANON², SITTHI KULABONG³,
ALONGKLOD TANOMTONG⁴, SUMALEE PHIMPHAN^{1,*}

¹ Biology Program, Faculty of Science and Technology, Phetchabun Rajabhat University, Phetchabun 67000, Thailand

² Scientific Committee. SuebNakasathien Foundation, 140 Tiwanon Road, Bang Kraso Subdistrict, Mueang Nonthaburi District, Nonthaburi Province 11000, Thailand

³ Faculty of Agro-Industrial Technology, Rajamangala University of Technology Tawan-ok Chantaburi Campus, Chantaburi Province 22210, Thailand

⁴ Program of Biology, Faculty of Science, Khon Kaen University, Muang, Khon Kaen 40002, Thailand

*Corresponding author. E-mail: sumalee.phim@pcru.ac.th; joodoof@gmail.com

Abstract. The first chromosome study of *Trigonostigma somphongsi* (Meinken, 1958) from Thailand. Specimens were collected from Pak Phli District, Nakhornnayok Province, central Thailand. The mitotic chromosomes were directly prepared from kidney tissues of two males and two females. Conventional staining and Ag-NOR banding techniques were applied to stain the chromosomes. The results shown that diploid chromosome number of *T. somphongsi* was $2n=50$, the fundamental numbers (NF) were 92 in both male and female. The types chromosomes consisted of 10 metacentric, 20 submetacentric, 12 acrocentric and 8 telocentric chromosomes. The result also exhibited that the interstitial nucleolar organizer regions (NORs) were clearly observed at the short arm of chromosome pair 2. The karyotype formula could be deduced as: $2n$ (diploid) $50=10m+20sm+12a+8t$.

Keywords: karyotype, chromosome, *Trigonostigma somphongsi*.

INTRODUCTION

The *Trigonostigma somphongsi* is a critically endangered species in natural habitat of Thailand. It has been discovered in a deep water rice field, floodplain of Bangpakong Basin, Nakhornnayok Province, central Thailand. (Petsut et al. 2014). The *Trigonostigma* is a genus of small cyprinid fish found in Southeast Asia. There are currently five recognized species in this genus including *T. heteromorpha*, *T. hengeli*, *T. somphongsi*, *T. espei* and *T. truncate* (Tan, 2020). The *T. somphongsi* was found in the rice field and in a



Figure 1. The characteristic of *T. somphongsi*. Scale bar indicate 0.2 cm.

densely vegetated ditch, which was flooded by a nearby river. It differs from the rest of the genus in its dark pattern which occurs as a strait horizontal line that extends from the base of the caudal fin and ends just after it passed the posterior part of the dorsal fin, instead of showing the wedge-like marking of other species in the genus (Petsut et al, 2014) (Fig. 1). Although the Somphong's rasbora is important for fish biodiversity of Thailand, there were quite scarce data about cytogenetics in these fishes especially banding analysis in fish chromosomes.

Up to date, information about karyotypes in genus *Trigonostigma* are rare and usually based on conventional staining technique. Only *T. heteromorpha* has been published, and the result showed that the diploid chromosome number ($2n$) was 50 (Khuda 1979). The cytogenetic studies using conventional staining technique provided valuable information on the great karyotype diversity shown by these animals. Analyses of cytogenetic markers, included the number and karyotype formula, number and location of nucleolar organizer regions (NORs). The present study is the first report on the chromosomal characteristics of *T. somphongsi* determined using conventional staining and Ag-NOR banding techniques. *T. somphongsi* is a critically endangered species in natural habitats of Thailand. The results enhance the level of cytogenetic information available and enable future comprehensive studies to be conducted on taxonomy and evolutionary relationships. Moreover, the data provide useful basic information for conservation and on breeding practices as well as an analysis of the chromosomal evolution of this species of *Trigonostigma*.

MATERIALS AND METHODS

Sample collection

Two males and two female of *T. somphongsi* were collected from agricultural land, Pak Phli District, Nakhornnayok Province, central Thailand, and were grown

starting from May 2023 (permission from an ethical committee ID U1-04498-2559). All specimens were maintained in aerated, flowing seawater aquaria until the analysis.

Chromosome preparation

Chromosomes preparations were obtained from kidney by cell suspension technique. Briefly, the kidney was cut into small pieces and then mixed with 0.075 M potassium chloride (KCl). After discarding all large piece tissues, cell sediments were transferred to a centrifuge tube and incubated for 30 minutes, then centrifuged at 1,500 rpm for 5 minutes. The KCl was discarded from the supernatant after centrifugation at 1,500 rpm for 5 minutes. Cells were fixed in fresh cool fixative (3 methanol:1 glacial acetic acid) and gradually made up to 8 ml before centrifuging again at 1,500 rpm for 5 minutes, whereupon the supernatant was discarded. Fixation was repeated until the supernatant was clear. The mixture was dropped onto a clean and cold slide by micropipette followed by air-drying technique.

Chromosome staining

Conventional staining of the chromosomes in the air-dried slides was done using 10% Giemsa solution for 10 minute (Rooney 2001). Ag-NOR banding was carried out following method of Howell and Black (1980) by adding 4 drops of 50% silver nitrate and 2% gelatin on slides. The slides were then sealed with cover glasses and incubated at 60°C for 5 minutes. After that the slides were soaked in distilled water until the cover glasses were separated.

Chromosome checking

Twenty clearly observable cells with well spread chromosomes of each male and female were selected and photographed under Olympus Bx63 microscope. Metaphase figures were analyzed according to the chromosome classification of Turpin and Lejeune (1965). The length of the short arm chromosome (Ls) and the long arm chromosome (Ll) were measured and the length of the total arm chromosome (LT, $LT = Ls + Ll$) was calculated. The relative length (RL), the centromeric index (CI), and standard deviation (SD) of RL and CI were estimated. The CI ($q/p + q$) between 0.50-0.59, 0.60-0.69, 0.70-0.89, and 0.90-0.99 were described as metacentric, submetacentric, acrocentric and telocentric chromo-

somes, respectively. The fundamental number (NF, number of chromosome arms) was obtained by assigning a value of 2 to metacentric, submetacentric and acrocentric chromosomes and 1 to telocentric chromosome. All parameters were used in karyotyping and idiogram.

RESULTS AND DISCUSSION

This is the first report on cytogenetic characterization using conventional staining and Ag-NOR banding techniques for *T. somphongsi*. The results indicated diploid chromosome number ($2n$) were found 50 in all studies samples as show in Fig. 2. This result is coincident with *T. heteromorpha* reports by Khuda 1979, and similar to another species in *Rasbora* (Post 1965; Manna and Khuda-Bukhsh 1977; Khuda-Bukhsh et al. 1979; Donsakul and Magtoon 1995; Donsakul and Magtoon 2002; Seetapan and Moeikum 2004; Donsakul et al. 2005; Donsakul et al. 2009; Yeesaem et al. 2019; Aiumsumang et al. 2021; 2022). From the previous report, most of cyprinid species have $2n=50$, chromosome consisting of both mono- and bi-arm chromosomes. There is no observation of strange size chromosomes related to sex, which is in accordance to the author of this genus (Khuda 1979). The types chromosomes of *T. somphongsi* were 10 metacentric, 20 submetacentric, 12 acrocentric and 8 telocentric chromosomes. The mean values calculated from twenty mitotic metaphases showed the centromeric index of chromosome complements ranging from 0.569 ± 0.001 to 1.000 ± 0.000 (Table 1).

To our results, NORs could be observed in one pair of chromosomes in both male and female of *T. som-*

phongsi. The result demonstrated that the chromosome marker shows in the chromosome pair 2, which is metacentric chromosome (Fig. 3). An important characteristic of Nucleolar Organizer Regions (NORs) in fish is related to that it has inter- and intra-species polymorphism. NORs characters can be a cytogenetic marker for cytotaxonomic studies and also have been used for studying of phylogenetic relationships among the Cyprinid fishes (Amemyia and Gold 1988; Galetti Jr 1998; Almeida-Toledo et al. 2000). The important karyotype feature of *T. somphongsi* is the symmetrical karyotype, which were found in four types of chromosomes (metacentric, submetacentric, acrocentric, and telocentric chromosomes). Figure 4 show the idiograms from conventional staining and Ag-NOR banding techniques. The karyotype formula could be deduced as: $2n$ (diploid) $50=10m+20sm+12a+8t$. The study on fish chromosomes is the basic knowledge which can be applied for the several fields such as classification, evolution, heredity, systematic (Gold et al. 1990; Ueda et al. 2001; Barat et al. 2002; Barat and Sahoo 2007), breeding, rapid production of inbred lines and cytotaxonomy (Kirpichnikov 1981). Furthermore, cytogenetic studies on fish have also been used as biological indicator to determine the ecological toxicology (Klinkhardt, 1993) and cytogenetic techniques have been widely applied to improve farmed stocks in many aquaculture species in the World (Beardmore et al. 2001; Desprez et al. 2003).

Here, we have that the karyotype of *T. somphongsi* is $2n=50$ and might represent a derived character, probably also shared by all members of the *Trigonostigma* clade. Besides, our study is the first cytogenetic karyotype data to describe in detail the karyotypic features of the Som-




Figure 2. Metaphase chromosome plates and karyotypes of the *T. somphongsi* by conventional staining. Species share the karyotype composed of 50 chromosomes. Scale bar indicate 5 μ m.

Table 1. Karyomorphological details of *T. somphongsi* from 20 metaphases chromosome, $2n$ (diploid)=50.

Chro. pair	Ls	Ll	LT	RL $\pm$ SD	CI $\pm$ SD	Chro. size	Chro. type
1	2.651	2.965	5.616	0.047 $\pm$ 0.002	0.569 $\pm$ 0.001	Large	metacentric
2	2.547	2.882	5.429	0.045 $\pm$ 0.002	0.574 $\pm$ 0.001	Large	metacentric
3	2.433	2.716	5.149	0.043 $\pm$ 0.003	0.571 $\pm$ 0.002	Large	metacentric
4	2.337	2.709	5.046	0.040 $\pm$ 0.001	0.583 $\pm$ 0.002	Medium	metacentric
5	2.927	2.609	5.536	0.039 $\pm$ 0.001	0.583 $\pm$ 0.002	Medium	metacentric
6	2.800	3.982	6.782	0.045 $\pm$ 0.002	0.688 $\pm$ 0.004	Large	submetacentric
7	2.692	3.237	5.929	0.044 $\pm$ 0.001	0.680 $\pm$ 0.001	Large	submetacentric
8	2.764	3.753	6.517	0.042 $\pm$ 0.004	0.682 $\pm$ 0.002	Large	submetacentric
9	2.985	3.706	6.691	0.040 $\pm$ 0.004	0.674 $\pm$ 0.002	Large	submetacentric
10	2.597	3.646	6.243	0.037 $\pm$ 0.005	0.694 $\pm$ 0.003	Medium	submetacentric
11	2.564	3.480	6.044	0.035 $\pm$ 0.003	0.692 $\pm$ 0.005	Medium	submetacentric
12	2.488	2.842	5.330	0.030 $\pm$ 0.003	0.653 $\pm$ 0.001	Medium	submetacentric
13	2.488	2.842	5.330	0.030 $\pm$ 0.003	0.653 $\pm$ 0.001	Medium	submetacentric
14	2.293	4.470	6.763	0.029 $\pm$ 0.008	0.669 $\pm$ 0.002	Medium	submetacentric
15	2.121	3.942	6.063	0.025 $\pm$ 0.003	0.664 $\pm$ 0.003	Medium	submetacentric
16	0.549	3.998	4.547	0.041 $\pm$ 0.001	0.898 $\pm$ 0.004	Large	acrocentric
17	0.404	3.565	3.969	0.038 $\pm$ 0.001	0.894 $\pm$ 0.004	Large	acrocentric
18	0.354	3.445	3.799	0.038 $\pm$ 0.001	0.871 $\pm$ 0.004	Large	acrocentric
19	0.344	3.014	3.358	0.037 $\pm$ 0.001	0.874 $\pm$ 0.004	Medium	acrocentric
20	0.175	2.985	3.160	0.035 $\pm$ 0.001	0.872 $\pm$ 0.004	Medium	acrocentric
21	0.168	2.014	2.182	0.031 $\pm$ 0.001	0.890 $\pm$ 0.004	Medium	acrocentric
22	0.000	4.624	4.624	0.029 $\pm$ 0.001	1.000 $\pm$ 0.000	Large	telocentric
23	0.000	3.615	3.615	0.027 $\pm$ 0.001	1.000 $\pm$ 0.000	Large	telocentric
24	0.000	3.550	3.550	0.026 $\pm$ 0.001	1.000 $\pm$ 0.000	Medium	telocentric
25	0.000	2.989	2.989	0.026 $\pm$ 0.001	1.000 $\pm$ 0.000	Medium	telocentric

Remarks: Ls=short arm chromosome, Ll=length of long arm chromosome, LT=length of total chromosomes, RL=relative length, CI=centromeric index, SD=standard deviation.




Figure 3. Metaphase chromosome plates and karyotypes of the *T. somphongsi*. The arrows indicate NOR banding by Ag-NOR staining technique. Scale bar indicate 5 μ m.




Figure 4. Idiogram showing lengths and shape of chromosomes of the *T. somphongsi*, *n* (haploid)=50, by conventional staining (A) and Ag-NOR staining technique (B). The arrows indicate NOR banding.

phong's rasbora species. The data can be a support for the investigation of chromosomal cytotaxonomy evolutionary history of relationships, conservation and on breeding practices within *Trigonostigma*.

ACKNOWLEDGEMENTS

This work was financially supported by Critical Ecosystem Partnership Fund (CEPF). We would like to thank Phetchabun Rajabhat University for available help.

REFERENCES

- Aiumsumang S, Phimphan S, Suwannapoom C, Chaiyasan P, Supiwong W, Tanomtong A. 2021. A comparative chromosome study on five Minnow fishes (Cyprinidae, Cypriniformes) in Thailand. *Caryologia* 74(1): 89–196. <https://doi.org/10.36253/caryologia-1017>.
- Aiumsumang S, Phimphan S, Tanomtong A, Supiwong W. 2022. Chromosome study of *Rasbora trilineata* and *Rasbora borapetensis* (Cyprinidae, Cypriniformes): Reveal by conventional staining technique. *Science Technology and Engineering Journal* 8(2): 29–39.
- Almeida-Toledo, LF, Foresti F, Toledo-Filho SA. 1984. Complex sex chromosome system in *Eigenmannia* sp. (Pisces, Gymnotiformes). *Genetica* 64: 165–169. <https://doi.org/10.1007/BF00115340>.
- Amemiya CT, Gold JR. 1988. Chromosomal NORs as taxonomic and systematic characters in North American cyprinid fishes. *Genetica* 76: 81–90. <https://doi.org/10.1007/BF00058806>.
- Barat A, Sahoo PK. 2007. Karyotype analysis of *Channa punctatus* (Pisces) using restriction endonucleases. *Cytologia* 72: 471–473. <https://doi.org/10.1508/cytologia.72.471>.
- Barat A, Sahoo PK, Ponniah AG. 2002. Karyotype and Nucleolar Organizer Regions (NORs) in a few hill stream fishes. In: Ayyappan, S., Jena, J. K. and Joseph, M. M. (Eds.), *The Fifth Indian Fisheries Forum Proceedings*, AFSIB, Mangalore and AoA, Bhubaneswar, pp.111–114.
- Beardmore JA, Mair GC, Lewis RI. 2001. Monosex male production in finfish as exemplified by tilapia: applications, problems, and prospects. *Aquaculture* 197: 283–301. doi:10.1016/S0044-8486(01)00590-7.
- Desprez D, Mélard C, Hoareau MC, Belleme'ne Y, Bosc P, Baroiller JF. 2003. Inheritance of sex in two ZZ pseudofemale lines of tilapia, *Oreochromis aureus*. *Aquaculture* 218: 131–140. [https://doi.org/10.1016/S0044-8486\(02\)00572-0](https://doi.org/10.1016/S0044-8486(02)00572-0).

- Donsakul T, Magtoon W. 1995. Karyological studies of *Puntius brevis* (Bleeker, 1850) and *Cyclocheilichthys enoplos* in Thailand. Proceeding of 21st Congress on Science and Technology of Thailand, pp. 292–293.
- Donsakul T, Magtoon W. 2002. Karyotype of *Rasbora Caudimaculata*, *R. myersi*, *R. retrodorsalis*, *R. paviei* from Thailand. Academic conference seminar for research papers of Srinakharinwirot University (pp. 1–7). Bangkok.
- Donsakul T, Magtoon W, Rangsiruji W. 2005. Karyotype of five species of fish (Pla siew) in subfamily Rasboranae. 31st Congress on Science and Technology of Thailand. Suranaree University of Technology. Nakhon Ratchasima Province Thailand.
- Donsakul T, Rangsiruji A, Magtoon W. 2009. Karyotypes of five cyprinid fishes (Cyprinidae, Danioninae Danionini): *Rasbora agilis*, *R. dorsiocellata*, *R. rubrodorsalis*, *Boraras maculate* and *B. urophthalmoides* from Thailand. In Proceedings of the 47th Kasetsart University Annual Conference (pp. 320–327). Kasetsart University, Thailand
- Galetti Jr PM. 1998. Chromosome diversity in neotropical fish NOR studies. Italian Journal of Zoology. 65: 53–56. <https://doi.org/10.1080/11250009809386795>.
- Gold JR, Li YC, Shipley NS, Powers PK. 1990. Improved methods for working with fish chromosomes with a review of metaphase chromosome banding. Journal of Fish Biology 37: 563–575. <https://doi.org/10.1111/j.1095-8649.1990.tb05889.x>.
- Howell WM, Black DA. 1980. Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: A 1-step method. Experientia 36: 1014–1015. <https://doi.org/10.1007/BF01953855>.
- Khuda-Bukhsh AR. 1979. Karyology of two species of hillstream fishes, *Barilius bendelisis* and *Rasbora daniconius* (Fam: Cyprinidae). Current Sci. 48(17): 793–794.
- Manna GK, Khuda-Bukhsh, A.R. 1977. Karyomorphology of cyprinid fishes and cytological evaluation of the family. The Nucleus, 20: 119–127
- Petsut N, Panitvong N, Kulabong S, Petsut J, Nonpayom C. 2014. The first record of *Trigonostigma somphongsi* (Meinken, 1958), a critically endangered species, in its natural habitat of Thailand (Cypriniformes Cyprinidae). Biodiversity Journal 5(4): 471–474.
- Post A. 1965. Vergleichende Untersuchungen der Chromosomenzahlen bei Süßwasser-Teleosteen. In: Gyldenholm, A.O. and Scheel, J.J. Chromosome number of fishes I. Journal Fish Biology, 3: 47–93.
- Rooney DE. 2001. Human Cytogenetics: Constitutional analysis: A Practical approach. 3rd ed. London: Oxford University Press.
- Seetapan K, Moeikum T. 2004. Karyotypes of ten Cyprinid fishes (Family Cyprinidae). Journal Agricultural Extension, 22: 92–101
- Tan HH. 2020. *Trigonostigma truncata*, a new species of harlequin rasbora from Malay Peninsula (Teleostei: Danionidae). Raff. Bull. Zool. 68: 421–433. <https://doi.org/10.26107/RBZ-2020-0058>
- Turpin R, Lejeune J. 1965. Les Chromosomes Humains. Gauthier Pillars.
- Ueda T, Naoi H, Arai R. 2001. Flexibility on the karyotype evolution in bitterlings (Pisces, Cyprinidae). Genetica 111: 423–432. <https://doi.org/10.1023/a:1013703717626>
- Yeesaem N, Jantarat S, Yeesin P. 2019. Cytogenetic Characterization of *Rasbora einthovenii* in Sirindhorn Peat Swamp Forest, Narathiwat Province. Journal of Fisheries Technology Research, 13: 58–68.