

Source details

Nonlinear Functional Analysis and Applications

Scopus coverage years: from 2017 to 2023

Publisher: Kyungnam University Press

ISSN: 1229-1595 E-ISSN: 2466-0973

Subject area: Mathematics: Analysis Mathematics: Numerical Analysis Mathematics: Applied Mathematics
Mathematics: Control and Optimization

Source type: Journal

[View all documents >](#) [Set document alert](#) [Save to source list](#)

CiteScore 2022 1.6 ⓘ

SJR 2022 0.361 ⓘ

SNIP 2022 0.797 ⓘ

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology ⓘ

CiteScore 2022 counts the citations received in 2019-2022 to articles, reviews, conference papers, book chapters and data papers published in 2019-2022, and divides this by the number of publications published in 2019-2022. [Learn more >](#)

CiteScore 2022 ▾

1.6

=

356 Citations 2019 - 2022

228 Documents 2019 - 2022

Calculated on 05 May, 2023

CiteScoreTracker 2023 ⓘ

1.9

=

469 Citations to date

241 Documents to date

Last updated on 05 April, 2024 • Updated monthly

CiteScore rank 2022 ⓘ

Category	Rank	Percentile
Mathematics		
Analysis	#99/187	47th
Mathematics		
Numerical Analysis	#52/85	39th
Mathematics		
Applied Mathematics	#393/609	35th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#) ⓘ

About Scopus

- What is Scopus
- Content coverage
- Scopus blog
- Scopus API
- Privacy matters

Language

- 日本語版を表示する
- 查看简体中文版本
- 查看繁體中文版本
- Просмотр версии на русском языке

Customer Service

- Help
- Tutorials
- Contact us

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

All content on this site: Copyright © 2024 Elsevier B.V. ↗, its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the Creative Commons licensing terms apply. We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.



Nonlinear Functional Analys

Welcome. Please, turn your
ADBLOCK off!

It looks like you're using an ad blocker. That's
okay—who doesn't? But without advertising
income, we can't keep making this site awesome.

COUNTRY

South Korea

Universities and research
institutions in South Korea

Media Ranking in South Korea

SUBJECT ARE

Mathemati
Analysi:
Applied
Control
Numeri

H-INDEX

13

Thank You!

REFRESH

PUBLICATION TYPE

Journals

ISSN

12291595, 24660973

COVERAGE

2017-2023

INFORMATION

Homepage

How to publish in this journal

jongkyuk@kyungnam.ac.kr

SCOPE

The international mathematical journal NFAA will publish carefully selected original research papers on nonlinear functional analysis and applications, that is, ordinary differential equations, all kinds of partial differential equations, functional differential equations, integrodifferential equations, control theory, approximation theory, optimal control, optimization theory, numerical analysis, variational inequality, asymptotic behavior, fixed point theory, dynamic systems and complementarity problems. Papers for publication will be communicated and recommended by the members of the Editorial Board.

Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS ?

1

Thai Journal of Mathematics

THA

81%

similarity

2

Panamerican Mathematical Journal

USA

74%

similarity

3

Fixed Point Theory and Algorithms for Sciences and

GBR

72%

similarity



4

Carpathian Journal of Mathematics

ROU

71%

similarity




Welcome. Please, turn your
ADBLOCK off!








It looks like you're using an ad blocker. That's okay—who doesn't? But without advertising income, we can't keep making this site awesome.

Thank You!

ns per document



● Cites / Doc. (4 years)
● Cites / Doc. (3 years)
● Cites / Doc. (2 years)



Nonlinear Functional Analysis and Applications




Show this widget in your own website

Just copy the code below and paste within your html code:

`<a href="https://www.scimagojr.com/journalsearch.php?q=21100838755&tip=sid&clean=0"`

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool**.





certificate for acceptance SP- 22- 001-IWANA2022

1 ข้อความ

김종규 <jongkyuk@kyungnam.ac.kr>

14 เมษายน 2567 เวลา 00:02

ถึง: "nuttapol.pak@pcru.ac.th" <nuttapol.pak@pcru.ac.th>

Dear Prof. Nuttapol Pakkaranang (nuttapol.pak@pcru.ac.th)

NFAA-23-232

Title: Accelerated Strongly Convergent Extragradient Algorithms to solve Variational Inequalities and Fixed Point Problems in Real Hilbert Spaces

Authors: N. Wairojjana, N. Pholasa, C. Khunpanuk and N. Pakkaranang

Corresponding author: Nuttapol Pakkaranang

Dated: April 14, 2024

How are you?

Your paper was accepted to the journal Nonlinear Functional Analysis and Applications(NFAA).

(**Nonlinear Functional Analysis and Applications**)

(<http://nfaa.kyungnam.ac.kr/journal-nfaa/index.php/NFAA>)

Please make the NFAA-style file (La- tex-file like attachment sample file) of your paper and send it to me as soon as possible.

You should take more time, if needed, to check them carefully.

At any rate, please try to send the corrected NFAA-style file as quickly as possible.

It is best to return attached , by e-mail: (jongkyuk@kyungnam.ac.kr).

Please write down the 2020 Mathematics, Subject Classification codes, Keywords and phrases, and sort running title in your paper.

The list of references must be numbered in alphabetical order by the first author's surname.

Please use the indicated reference number in all future correspondence.

Should you have any questions, please feel free to contact me. Thank you for your attention

PS; The journal NFAA is cited in the SCOPUS.

<https://www.scopus.com/sourceid/21100838755?origin=sbrowse> .

In order to have publication in NFAA,

1. You have to pay the Article Processing Charge(APC), 350:00 US\$.

3. After receiving your style file, then I will send you the INVOICE for the processing Charge soon

Cordially yours

J. K. Kim

Editor-in-Chief of the Journals:

Nonlinear Functional Analysis and Applications(NFAA),

International Journal of Mathematical Sciences(IJMS)

Department of Mathematics Education

Kyungnam University,

Changwon, Gyeongnam, 631-701, Korea

Tel:82-(55)-249-2211 ,

Fax:82-(55)-243-8609

E-mail: jongkyuk@kyungnam.ac.kr

<http://nfaa.kyungnam.ac.kr/>

<http://www.serialspub.com>

https://www.researchgate.net/profile/Jong_Kim17/publications/

<http://scholar.google.com.pk/citations?user=k2Czf78AAAAJ>



Nonlinear Functional Analysis and Applications

Volume 29

Number 2

2024

Editor-in-Chief

Jong Kyu Kim, Kyungnam University, Korea

Print ISSN 1229-1595 / Online ISSN 2466-0973

Published Online: 28 May 2024



Nonlinear Functional Analysis and Applications (NFAA)

HOME	ABOUT	LOG IN	REGISTER	SEARCH	CURRENT	ARCHIVES	AIM AND
SCOPE	EDITORIAL BOARD						

Home > **Editorial Board**

Editorial Board

Editor-in-Chief

Jong Kyu Kim, Kyungnam University, Korea

E-mail: jongkyuk@kyungnam.ac.kr

Managing Editors

Kyung Soo Kim, Kyungnam University, Korea

E-mail: kksmj@kyungnam.ac.kr

Xiao Long Qin, UESTC, China

E-mail: ljhhqx1@aliyun.com

Honorary Editorial Board

Nasir U. Ahmed, University of Ottawa, Canada

Ravi P. Agarwal, Texas A&M University, USA

Ki Sik Ha, Pusan National University, Korea

Sehie Park, Seoul National University, Korea

H. M. Srivastava, University of Victoria, Canada

Ram U. Verma, Texas State University, USA

Editorial Board

George Anastassiou, University of Memphis, USA

Jurgen Appell, University of Wurzburg, Germany

Drumi Bainov, Medical University of Sofia, Bulgaria

Jean-Bernard Baillon, Paris 1 Pantheon-Sorbonne University, France

Krishnan Balachandran, Bharathiar University, India

Ronald E. Bruck, University of Southern California, USA

Pietro Cerone, Victoria University of Technology, Australia

Shih-sen Chang, Sichuan University, China

Sui Sun Cheng, Tsing Hua University, Taiwan

Yeol Je Cho, Gyeongsang National University, Korea

U Jin Choi, KAIST, Korea

Q-Heung Choi, Inha University, Korea

C. Corduneanu, University of Texas, USA

Sever Dragomir, Victoria University of Technology, Australia

Norimichi Hirano, Yokohama National University, Japan

Nan-Jing Huang, Sichuan University, China

Jong Soo Jung, Dong-A University, Korea

Athanassios G. Kartsatos, University of South Florida, USA

Do Sang Kim, Pukyong National University, Korea

Won Kyu Kim, Chungbuk National University, Korea

Poom Kumam, King Mongkut's University of Technology, Thailand

Anthony To-Ming Lau, University of Alberta, Canada

Byung Soo Lee, Kyungsung University, Korea

Gue Myung Lee, Pukyong National University, Korea

Yong-Hoon Lee, Pusan National University, Korea

Gang Li, Yangzhou University, China

Bor Luh Lin, University of Iowa, USA

Chia-Shiang Lin, Bishop's University, Canada

En-Bing Lin, Texas A&M International University, USA

Lai-Jiu Lin, National Changhua University of Education Changhua, Taiwan

USER

Username

Password

☐ Remember me

NOTIFICATIONS

- [View](#)
- [Subscribe / Unsubscribe](#)

JOURNAL CONTENT

Search

All

Browse

- [By Issue](#)
- [By Author](#)
- [By Title](#)

FONT SIZE

INFORMATION

- [For Readers](#)
- [For Authors](#)
- [For Librarians](#)

Xinzhi Liu, University of Waterloo, Canada
Giuseppe Marino, Universita della Calabria, Italy
Gheorghe Morosanu, Central European University, Hungary
Mohammad Sal Moslehian, Ferdowsi University, Iran
Shin-ichi Nakagiri, Kobe University, Japan
Young Man Nam, Kyungnam University, Korea
Muhammad Aslam Noor, COMSATS Institute of Information, Pakistan
Donal O'Regan, National University of Ireland, Ireland
Jong Yeoul Park, Pusan National University, Korea
Nicolae H. Pavel, Ohio University Athens, USA
Josip Pecaric, University of Zagreb, Croatia
Themistocles M. Rassias, National Technical University of Athens, Greece
Simeon Reich, The Technion-Israel Institute of Technology, Israel
Behzad Djafari Rouhani, Shahid Beheshti University, Iran
Brailey Sims, The University of Newcastle, Australia
Wataru Takahashi, Tokyo Institute of Technology, Japan
Alexander G. Ramm, Kansas State University, USA
Kok-Keong Tan, Dalhousie University, Canada
Jeong Sheok Ume, Changwon National University, Korea
Hong-Kun Xu, National Sun Yat-sen University, Taiwan
Masahiro Yamamoto, The University of Tokyo, Japan
George Xian-Zhi Yuan, The University of Queensland, Australia
Haiyun Zhou, Shijiazhuang Mechanical Engineering University, China

ISSN: 1229-1595 (Print), 2466-0973 (Online)

(51767) 7 Kyungnamdaehak-ro, Masanhappo-gu, Changwon-si, Gyeongsangnam-do, Republic of Korea



Nonlinear Functional Analysis and Applications (NFAA)

HOME	ABOUT	LOG IN	REGISTER	SEARCH	CURRENT	ARCHIVES	AIM AND SCOPE
SCOPE	EDITORIAL BOARD						

Home > Archives > **Volume 29, Number 2, June 2024**

Volume 29, Number 2, June 2024

Table of Contents

Articles

ACCELERATED STRONGLY CONVERGENT EXTRAGRADIENT ALGORITHMS TO SOLVE VARIATIONAL INEQUALITIES AND FIXED POINT PROBLEMS IN REAL HILBERT SPACES	PDF
<i>Nopparat Wairojjana, Nattawut Pholasa, Chainarong Khunpanuk, Nuttapol Pakkaranana</i>	307-332
THE GLOBAL EXISTENCE AND BEHAVIOR OF RADIAL SOLUTIONS OF A NONLINEAR p-LAPLACIAN TYPE EQUATION WITH SINGULAR COEFFICIENTS	PDF
<i>Hikmat El Baghourti, Arij Bouzelmate</i>	333-360
ON FIXED POINT THEOREMS SATISFYING COMPATIBILITY PROPERTY IN MODULAR G-METRIC SPACES	PDF
<i>Daniel Francis, Godwin Amechi Okeke, Ho Geun Hyun</i>	361-391
A MODIFIED KRASNOSELSKII-TYPE SUBGRADIENT EXTRAGRADIENT ALGORITHM WITH INERTIAL EFFECTS FOR SOLVING VARIATIONAL INEQUALITY PROBLEMS AND FIXED POINT PROBLEM	PDF
<i>Araya Kheawborisut, Wongvisarut Khuangsatung</i>	393-418
ON KIRCHHOFF TYPE EQUATIONS WITH SINGULAR NONLINEARITIES, SUB-CRITICAL AND CRITICAL EXPONENT	PDF
<i>Mohammed El Mokhtar Ould El Mokhtar, Saleh Fahad Aljurbua</i>	419-434
TAYLORS SERIES IN TERMS OF THE MODIFIED CONFORMABLE FRACTIONAL DERIVATIVE WITH APPLICATIONS	PDF
<i>Mohammed B. M. Altalla, B. Shanmukha, Ahmad El-Ajou, Mohammed N. A. Alkord</i>	435-450
L^r INEQUALITIES FOR POLYNOMIALS	PDF
<i>Reingachan N, Mayanglambam Singhajit Singh, Nirmal Kumar Singha, Khangembam Babina Devi, Barchand Chanam</i>	451-460
FIXED POINT THEOREMS OF EXTENSION AND MODIFIED EXTENSION α-F-CONTRACTION ON COMPLETE METRIC SPACE	PDF
<i>Saeed A. A. Al-Salehi, V. C Borkar</i>	461-475
MULTI-VALUED HICKS CONTRACTIONS IN b-MENGER SPACES	PDF
<i>Youssef Achtoun, Mohammed Sefian Lamarti, Ismail Tahiri</i>	477-485
FIXED POINT THEOREMS IN b-MENGER INNER PRODUCT SPACES	PDF
<i>Rachid Oubrahim</i>	487-499
INVESTIGATION OF A NEW COUPLED SYSTEM OF FRACTIONAL DIFFERENTIAL EQUATIONS IN FRAME OF HILFER-HADAMARD	PDF
<i>Ali Abd Alaziz Najem Al-Sudani, Ibrahim Abdurassool hammood Al-Nuh</i>	501-515
GENERAL MIXED HARMONIC VARIATIONAL INEQUALITIES	PDF
<i>Jong Kyu Kim, Avinash Lakshotra, Tirth Ram</i>	517-526
OPTIMAL STRATEGIES IN BIOECONOMIC DIFFERENTIAL GAMES: INSIGHTS FROM CHEBYSHEV TAU METHOD	PDF
<i>Shahd H. Alkharaz, Essam El-Siedy, Eliwa M. Roushdy, Muner M. Abou Hasan</i>	527-543
EXISTENCE RESULTS FOR BOUNDARY VALUE PROBLEMS OF VOLTERRA-FREDHOLM SYSTEM INVOLVING CAPUTO DERIVATIVE	PDF
<i>Shakir M. Atshan, Ahmed A. Hamoud</i>	545-558
CONTROL FUNCTION BASED COUPLED AND COMMON COUPLED FIXED POINT THEOREMS IN PARTIAL METRIC SPACES	PDF
<i>H. K. Nashine, G. V. V. Jagannadha Rao, W. H. Lim</i>	559-580
NEW BEST PROXIMITY POINT RESULTS FOR DIFFERENT TYPES OF NONSELF PROXIMAL CONTRACTIONS WITH AN APPLICATION	PDF
<i>Khairul Habib Alam, Yumnam Rohen, S. Surendra Singh, Kshetrimayum Mangjaobi Devi, L. Bishwakumar</i>	581-596
IMPACT OF FRACTIONAL CONFORMABLE DERIVATIVES ON A(H1N1) INFECTION MODEL	PDF
<i>Hind Ali Ahmad Eid, Mohammed N Alkord</i>	597-620

ISSN: 1229-1595 (Print), 2466-0973 (Online)

(51767) 7 Kyungnamdaehak-ro, Masanhappo-gu, Changwon-si, Gyeongsangnam-do, Republic of Korea

USER

Username

Password

☐ Remember me

[Log In](#)

NOTIFICATIONS

- [View](#)
- [Subscribe / Unsubscribe](#)

JOURNAL CONTENT

Search

All

[Search](#)

Browse

- [By Issue](#)
- [By Author](#)
- [By Title](#)

FONT SIZE

INFORMATION

- [For Readers](#)
- [For Authors](#)
- [For Librarians](#)

ACCELERATED STRONGLY CONVERGENT EXTRAGRADIENT ALGORITHMS TO SOLVE VARIATIONAL INEQUALITIES AND FIXED POINT PROBLEMS IN REAL HILBERT SPACES

Nopparat Wairojjana¹, Nattawut Pholasa²,
Chainarong Khunpanuk³ and Nuttapol Pakkaranang⁴

¹Applied Mathematics Program, Faculty of Science and Technology,
Valaya Alongkorn Rajabhat University under the Royal Patronage Pathum Thani Province,
Pathum Thani, 13180, Thailand
e-mail: nopparat@vru.ac.th

²School of Science, University of Phayao, Phayao 56000, Thailand
e-mail: nattawut_math@hotmail.com

³Mathematics and Computing Science Program, Faculty of Science and Technology,
Phetchabun Rajabhat University, Phetchabun 67000, Thailand
e-mail: improve2000ck@gmail.com

⁴Mathematics and Computing Science Program, Faculty of Science and Technology,
Phetchabun Rajabhat University, Phetchabun 67000, Thailand
e-mail: nuttapol.pak@pcru.ac.th

Abstract. Two inertial extragradient-type algorithms are introduced for solving convex pseudomonotone variational inequalities with fixed point problems, where the associated mapping for the fixed point is a ρ -demicontractive mapping. The algorithm employs variable step sizes that are updated at each iteration, based on certain previous iterates. One notable advantage of these algorithms is their ability to operate without prior knowledge of Lipschitz-type constants and without necessitating any line search procedures. The iterative sequence constructed demonstrates strong convergence to the common solution of the variational inequality and fixed point problem under standard assumptions. In-depth numerical applications are conducted to illustrate theoretical findings and to compare the proposed algorithms with existing approaches.

⁰Received December 22, 2022. Revised April 8, 2024. Accepted April 14, 2024.

⁰2020 Mathematics Subject Classification: 47J25, 47H09, 47H06, 47J05.

⁰Keywords: Variational inequality problem, fixed point problem, ρ -demicontractive operator, strong convergence theorems.

⁰Corresponding author: Nuttapol.Pakkaranang@pcru.ac.th.

1. INTRODUCTION

The investigation of common solution problems is driven by their potential applicability within mathematical models characterized by fixed point constraints, particularly in practical domains such as signal processing, network resource allocation, and image recovery. This aspect holds considerable significance across various fields, encompassing signal analysis, composite reduction, optimization methodologies, and image recovery challenges, as evidenced by relevant literature (e.g., Maing et al., [20, 21], Iiduka et al., [10], Qin et al., [27], An et al., [4]). This study scrutinizes two focal issues within this realm.

Let $\mathcal{K}$ denote a nonempty, closed and convex subset of a real Hilbert space $\mathcal{X}$ endowed with the inner product $\langle \cdot, \cdot \rangle$ and the corresponding induced norm $\| \cdot \|$. The study makes a significant contribution by exploring the convergence analysis of iterative algorithms designed to solve variational inequality problems and fixed point problems in real Hilbert spaces.

Considering an operator $\mathcal{N} : \mathcal{K} \rightarrow \mathcal{X}$, the variational inequality problem (VIP) (Stampacchia [30], 1964) is formulated as follows:

$$\text{Find } u^* \in \mathcal{K} \text{ such that } \langle \mathcal{N}(u^*), y - u^* \rangle \geq 0, \text{ for all } y \in \mathcal{K}. \quad (\text{VIP})$$

Let $VI(\mathcal{K}, \mathcal{N})$ denote the solution set corresponding to problem (VIP). Variational inequalities find applications across diverse domains, including but not limited to partial differential equations, optimization, engineering, applied mathematics, and economics (see, for instance, [1, 8, 11, 12, 15, 16, 19, 25, 31]). The variational inequality problem holds significant importance in the applied sciences. Numerous researchers have dedicated considerable effort not only to exploring the existence and stability of solutions but also to devising iterative algorithms for their resolution. Projection algorithms, in particular, play a pivotal role in ascertaining the numerical solutions to variational inequalities. [6, 7, 14, 17, 26, 29] along with others documented in works [2, 3, 9, 13, 23, 24, 35, 34].

The majority of algorithms employed for problem-solving rely on the projection technique computed within the feasible set $\mathcal{K}$. Among these methods, the extragradient algorithm, originating from the works of Korpelevich [17] and Antipin [5], stands out prominently. Consider $\mathcal{Q} : \mathcal{X} \rightarrow \mathcal{X}$ as a mapping. The associated fixed-point problem with $\mathcal{Q}$ is expressed as:

$$\mathcal{Q}(u^*) = u^*. \quad (\text{FP})$$

The solution set of the fixed point problem (FP) is denoted as $Fix(\mathcal{Q})$. A considerable portion of algorithms aimed at solving (FP) is derived from the foundational Mann iteration scheme. Specifically, commencing with $u_1 \in \mathcal{X}$, this scheme generates the sequence $\{u_{k+1}\}$ for every $k \geq 1$ according to the

recurrence relation:

$$u_{k+1} = \alpha_k u_k + (1 - \alpha_k) \mathcal{Q}(u_k). \quad (1.1)$$

For constructing weak convergence necessitates adherence to specific criteria by the variable sequence $\{\alpha_k\}$. The Halpern iteration introduces an alternative structured iterative algorithm, demonstrating heightened efficacy in achieving strong convergence within infinite-dimensional Hilbert spaces. The iterative sequence is delineated as follows:

$$u_{k+1} = \alpha_k u_1 + (1 - \alpha_k) \mathcal{Q}u_k, \quad (1.2)$$

where $u_1 \in \mathcal{X}$, and the sequence $\alpha_k \subset (0, 1)$ is non-summable and showcases gradual decline, conforming to the conditions:

$$\alpha_k \rightarrow 0 \quad \text{and} \quad \sum_{k=1}^{+\infty} \alpha_k = +\infty.$$

Additionally, the viscosity algorithm, as introduced by Moudafi [22], which merges the cost mapping $\mathcal{Q}$ with a contraction mapping iteratively, presents a generic variant of the Halpern iteration. Alongside the Halpern iteration, a generalized form known as the viscosity algorithm [22] exists, wherein the cost mapping $\mathcal{Q}$ is integrated with a contraction mapping within the iterative process. Lastly, the hybrid steepest descent approach, proposed in [33], represents another methodology offering substantial convergence benefits.

Tan et al. [32] introduced a new numerical algorithm called the extragradient viscosity algorithm. It's designed to solve variational inequalities that involve a fixed-point problem with a specific kind of mapping called a ρ -demicontractive mapping. This algorithm combines ideas from two other algorithms: the extragradient algorithm and the Mann-type technique. The authors showed that all these algorithms have strong convergence if the operator is both monotonic and satisfies the Lipschitz condition. These methods offer an advantage because they can be estimated numerically using optimization tools, as demonstrated in [32]. However, one drawback of these algorithms is that they heavily rely on viscosity and Mann-type techniques to achieve strong convergence. Strong convergence is crucial for iterative sequences, especially in situations with infinite dimensions. Only a few algorithms achieve strong convergence using inertial schemes. Using Mann and viscosity procedures can be challenging from an algorithmic perspective, potentially slowing down convergence and limiting practical usefulness. These algorithms also require more numerical and computational steps, making the system more complex.

This brings us to a fundamental question:

Can we make self-adjusting, strongly convergent inertial extragradient algorithms that don't need Mann and Viscosity-type methods to solve variational inequalities and fixed-point problems?

In response, we have developed two strong convergence extragradient-type algorithms. These are designed to solve monotone variational inequalities and the ρ -demicontractive fixed point problem in real Hilbert spaces. Our inspiration comes from the research discussed in [32]. Additionally, we have intentionally avoided using any hybrid techniques, like the Mann-type scheme or the viscosity scheme, to ensure the strong convergence of these algorithms.

The paper is organized into several sections. In Section 2, we present some fundamental findings. Section 3 introduces four distinct algorithms and verifies their convergence analysis. Lastly, Section 4 offers numerical data to showcase the practical application of the presented algorithms.

2. PRELIMINARIES

Let $\mathcal{K}$ denote a nonempty, closed and convex subset of a real Hilbert space $\mathcal{X}$. For any $u, y \in \mathcal{X}$, we establish the following properties:

- (i) $\|u + y\|^2 = \|u\|^2 + 2\langle u, y \rangle + \|y\|^2$;
- (ii) $\|u + y\|^2 \leq \|u\|^2 + 2\langle y, u + y \rangle$;
- (iii) $\|bu + (1 - b)y\|^2 = b\|u\|^2 + (1 - b)\|y\|^2 - b(1 - b)\|u - y\|^2$.

A metric projection $P_{\mathcal{K}}(u)$ of $u \in \mathcal{X}$ is defined as follows:

$$P_{\mathcal{K}}(u) = \arg \min\{\|u - y\| : y \in \mathcal{K}\}.$$

It is well established that $P_{\mathcal{K}}$ is nonexpansive and possesses the following significant properties:

- (1) $\langle u - P_{\mathcal{K}}(u), y - P_{\mathcal{K}}(u) \rangle \leq 0, \forall y \in \mathcal{K}$;
- (2) $\|P_{\mathcal{K}}(u) - P_{\mathcal{K}}(y)\|^2 \leq \langle P_{\mathcal{K}}(u) - P_{\mathcal{K}}(y), u - y \rangle, \forall y \in \mathcal{K}$.

Definition 2.1. Let $\mathcal{Q} : \mathcal{X} \rightarrow \mathcal{X}$ be a nonlinear mapping with $Fix(\mathcal{Q}) \neq \emptyset$. Then, $I - \mathcal{Q}$ is said to be demiclosed at zero if for any sequence $\{u_k\}$ in $\mathcal{X}$, the following statement holds:

$$u_k \rightharpoonup u \quad \text{and} \quad (I - \mathcal{Q})u_k \rightarrow 0 \quad \Rightarrow \quad u \in Fix(\mathcal{Q}),$$

where $Fix(\mathcal{Q}) = \{x \in \mathcal{X} : \mathcal{Q}x = x\}$ is the set of fixed points of $\mathcal{Q}$.

Definition 2.2. Let $\mathcal{N} : \mathcal{K} \rightarrow \mathcal{K}$ be an operator. It is said to be:

(1) *monotone* if

$$\langle \mathcal{N}(u_1) - \mathcal{N}(u_2), u_1 - u_2 \rangle \geq 0, \quad \forall u_1, u_2 \in \mathcal{K};$$

(2) *Lipschitz-continuous* with constant $L > 0$ such that

$$\|\mathcal{N}(u_1) - \mathcal{N}(u_2)\| \leq L\|u_1 - u_2\|, \quad \forall u_1, u_2 \in \mathcal{K};$$

(3) *sequentially weakly continuous* if a sequence $\{\mathcal{N}(u_k)\}$ converges weakly to $\mathcal{N}(u)$ for any sequence $\{u_k\}$ converging weakly to u .

Definition 2.3. Let $\mathcal{Q} : \mathcal{K} \rightarrow \mathcal{K}$ be a mapping such that $\text{Fix}(\mathcal{Q}) \neq \emptyset$. $\mathcal{Q}$ is said to be ρ -demicontractive if for any fixed number $0 \leq \rho < 1$, the following holds:

$$\|\mathcal{Q}(u_1) - u_2\|^2 \leq \|u_1 - u_2\|^2 + \rho\|(I - \mathcal{Q})(u_1)\|^2, \quad \forall u_2 \in \text{Fix}(\mathcal{Q}), u_1 \in \mathcal{X}.$$

Lemma 2.4. ([18]) Let $\mathcal{N} : \mathcal{X} \rightarrow \mathcal{X}$ be an operator that is L -Lipschitz continuous and monotone on $\mathcal{K}$. Consider $\mathcal{Q} = P_{\mathcal{K}}(I - \eta\mathcal{N})$, where $\eta > 0$. If $\{u_k\}$ is a sequence in $\mathcal{X}$ such that $u_k \rightarrow q$ and $u_k - \mathcal{N}(u_k) \rightarrow 0$, then $q \in VI(\mathcal{K}, \mathcal{N}) = \text{Fix}(\mathcal{Q})$.

Lemma 2.5. ([28]) Suppose that $\{c_k\} \subset [0, +\infty)$, $\{d_k\} \subset (0, 1)$ and $\{e_k\} \subset \mathbb{R}$ are sequences satisfying the following conditions:

$$c_{k+1} \leq (1 - d_k)c_k + d_k e_k, \quad \forall k \in \mathbb{N} \quad \text{and} \quad \sum_{k=1}^{+\infty} d_k = +\infty.$$

If $\limsup_{j \rightarrow +\infty} y_{k_j} \leq 0$ for any subsequence $\{c_{k_j}\}$ of $\{c_j\}$ such that

$$\liminf_{j \rightarrow +\infty} (c_{k_j+1} - c_{k_j}) \geq 0.$$

Then $\lim_{k \rightarrow +\infty} c_k = 0$.

3. MAIN RESULTS

In this section, we delve into the convergence analysis of two new inertial extragradient algorithms designed for solving fixed point and variational inequality problems. Initially, we scrutinize the proposed algorithms. To establish strong convergence, it is presupposed that the following conditions hold:

- ($\mathcal{N}1$) The common solution set, denoted by $\text{Fix}(\mathcal{Q}) \cap VI(\mathcal{K}, \mathcal{N})$, is nonempty.
- ($\mathcal{N}2$) The operator $\mathcal{N} : \mathcal{X} \rightarrow \mathcal{X}$ is monotone.
- ($\mathcal{N}3$) The operator $\mathcal{N} : \mathcal{X} \rightarrow \mathcal{X}$ is Lipschitz continuous.

- ($\mathcal{N}4$) The mapping $\mathcal{Q} : \mathcal{X} \rightarrow \mathcal{X}$ is ρ -demicontractive for $0 \leq \rho < 1$ and demiclosed at zero.
- ($\mathcal{N}5$) The operator $\mathcal{N} : \mathcal{X} \rightarrow \mathcal{X}$ is sequentially weakly continuous.

Algorithm 1 (Accelerated Tseng's Extragradient Algorithm With Monotone Variable Step Size Rule)

STEP 0: Choose $u_0, u_1 \in \mathcal{K}$, $\theta \in (0, 1)$, $\mu \in (0, 1)$, $\eta_1 > 0$, and a sequence $\{\varsigma_k\} \subset (0, 1)$ satisfying:

$$\lim_{k \rightarrow +\infty} \varsigma_k = 0 \quad \text{and} \quad \sum_{k=1}^{+\infty} \varsigma_k = +\infty.$$

STEP 1: Compute:

$$q_k = u_k + \theta_k(u_k - u_{k-1}) - \varsigma_k[u_k + \theta_k(u_k - u_{k-1})],$$

where θ_k is defined as follows:

$$0 \leq \theta_k \leq \hat{\theta}_k \quad \text{and} \quad \hat{\theta}_k = \begin{cases} \min \left\{ \frac{\theta}{2}, \frac{\chi_k}{\|u_k - u_{k-1}\|} \right\} & \text{if } u_k \neq u_{k-1}, \\ \frac{\theta}{2} & \text{otherwise.} \end{cases} \quad (3.1)$$

Additionally, a sequence $\chi_k = o(\varsigma_k)$ satisfies the condition $\lim_{k \rightarrow +\infty} \frac{\chi_k}{\varsigma_k} = 0$.

STEP 2: Compute:

$$y_k = P_{\mathcal{K}}(q_k - \eta_k \mathcal{N}(q_k)).$$

If $q_k = y_k$, then STOP; otherwise, proceed to **STEP 3**.

STEP 3: Compute:

$$p_k = y_k + \eta_k[\mathcal{N}(q_k) - \mathcal{N}(y_k)].$$

STEP 4: For any sequence $\alpha_k \subset (0, 1 - \rho)$, compute:

$$u_{k+1} = (1 - \alpha_k)p_k + \alpha_k \mathcal{Q}(p_k).$$

STEP 5: Compute:

$$\eta_{k+1} = \begin{cases} \min \left\{ \eta_k, \frac{\mu \|q_k - y_k\|}{\|\mathcal{N}(q_k) - \mathcal{N}(y_k)\|} \right\} & \text{if } \mathcal{N}(q_k) \neq \mathcal{N}(y_k), \\ \eta_k, & \text{otherwise.} \end{cases} \quad (3.2)$$

Update $k := k + 1$ and return to **STEP 1**.

Algorithm 2 (Accelerated Tseng's Extragradient Algorithm With Non-Monotone Variable Step Size Rule)

STEP 0: Choose initial points $u_0, u_1 \in \mathcal{K}$, parameters $\theta \in (0, 1)$, $\mu \in (0, 1)$, and $\eta_1 > 0$. Also, select a sequence $\{\beth_k\}$ such that $\sum_{k=1}^{+\infty} \beth_k < +\infty$ and a sequence $\{\varsigma_k\} \subset (0, 1 - \rho)$ satisfying:

$$\lim_{k \rightarrow +\infty} \varsigma_k = 0 \quad \text{and} \quad \sum_{k=1}^{+\infty} \varsigma_k = +\infty.$$

STEP 1: Compute

$$q_k = u_k + \theta_k(u_k - u_{k-1}) - \varsigma_k[u_k + \theta_k(u_k - u_{k-1})],$$

where θ_k is defined as follows:

$$0 \leq \theta_k \leq \hat{\theta}_k \quad \text{and} \quad \hat{\theta}_k = \begin{cases} \min \left\{ \frac{\theta}{2}, \frac{\chi_k}{\|u_k - u_{k-1}\|} \right\} & \text{if } u_k \neq u_{k-1}, \\ \frac{\theta}{2} & \text{otherwise,} \end{cases} \quad (3.3)$$

with $\chi_k = \mathcal{O}(\varsigma_k)$ satisfying $\lim_{k \rightarrow +\infty} \frac{\chi_k}{\varsigma_k} = 0$.

STEP 2: Compute

$$y_k = P_{\mathcal{K}}(q_k - \eta_k \mathcal{N}(q_k)).$$

If $q_k = y_k$, then STOP; otherwise, proceed to **STEP 3**.

STEP 3: Compute

$$p_k = y_k + \eta_k[\mathcal{N}(q_k) - \mathcal{N}(y_k)].$$

STEP 4: Choose a sequence $\alpha_k \subset (0, 1 - \rho)$. Compute

$$u_{k+1} = (1 - \alpha_k)p_k + \alpha_k \mathcal{Q}(p_k).$$

STEP 5: Compute

$$\begin{cases} \min \left\{ \eta_k + \beth_k, \frac{\mu \|q_k - y_k\|}{\|\mathcal{N}(q_k) - A(y_k)\|} \right\} & \text{if } \mathcal{N}(q_k) \neq \mathcal{N}(y_k), \\ \eta_k + \beth_k, & \text{otherwise,} \end{cases} \quad (3.4)$$

where $A(y_k)$ denotes a specific operator. Then, increment k and return to **STEP 1**.

Lemma 3.1. *Let $\{\eta_k\}$ be a sequence generated by the expression (3.2). Then, $\{\eta_k\}$ is monotonically decreasing and bounded by $\min \left\{ \frac{\mu}{L}, \eta_1 \right\} \leq \eta \leq \eta_1$.*

Proof. Given that the mapping $\mathcal{N}$ is Lipschitz continuous, we have:

$$\begin{aligned} \frac{\mu \|q_k - y_k\|}{\|\mathcal{N}(q_k) - \mathcal{N}(y_k)\|} &\geq \frac{\mu \|q_k - y_k\|}{L \|q_k - y_k\|} \\ &\geq \frac{\mu}{L}. \end{aligned} \quad (3.5)$$

This implies that the sequence $\{\eta_k\}$ is monotonically decreasing and bounded. $\square$

Lemma 3.2. *A sequence $\{\eta_k\}$ generated by the expression (3.4) is convergent to η and bounded by $\min\left\{\frac{\mu}{L}, \eta_1\right\} \leq \eta \leq \eta_1 + P$, where*

$$P = \sum_{k=1}^{+\infty} \beth_k.$$

Proof. Given that the mapping $\mathcal{N}$ is Lipschitz continuous, we have

$$\begin{aligned} \frac{\mu\|q_k - y_k\|}{\|\mathcal{N}(q_k) - \mathcal{N}(y_k)\|} &\geq \frac{\mu\|q_k - y_k\|}{L\|q_k - y_k\|} \\ &\geq \frac{\mu}{L}. \end{aligned} \quad (3.6)$$

By the definition of η_{k+1} , we have

$$\min\left\{\frac{\mu}{L}, \eta_1\right\} \leq \eta_k \leq \eta_1 + P.$$

Let

$$[\eta_{k+1} - \eta_k]^+ = \max\{0, \eta_{k+1} - \eta_k\}$$

and

$$[\eta_{k+1} - \eta_k]^- = \max\{0, -(\eta_{k+1} - \eta_k)\}.$$

Using the definition of $\{\eta_k\}$, we have

$$\sum_{k=1}^{+\infty} (\eta_{k+1} - \eta_k)^+ = \sum_{k=1}^{+\infty} \max\{0, \eta_{k+1} - \eta_k\} \leq P < +\infty. \quad (3.7)$$

This implies that the series $\sum_{k=1}^{+\infty} (\eta_{k+1} - \eta_k)^+$ converges. Next, we need to demonstrate the convergence of

$$\sum_{k=1}^{+\infty} (\eta_{k+1} - \eta_k)^-.$$

Suppose $\sum_{k=1}^{+\infty} (\eta_{k+1} - \eta_k)^- = +\infty$. Then we obtain

$$\eta_{k+1} - \eta_k = (\eta_{k+1} - \eta_k)^+ - (\eta_{k+1} - \eta_k)^-.$$

Hence,

$$\eta_{k+1} - \eta_1 = \sum_{k=0}^k (\eta_{k+1} - \eta_k) = \sum_{k=0}^k (\eta_{k+1} - \eta_k)^+ - \sum_{k=0}^k (\eta_{k+1} - \eta_k)^-. \quad (3.8)$$

Letting $k \rightarrow +\infty$ in (3.8), we get $\eta_k \rightarrow -\infty$ as $k \rightarrow +\infty$, which is a logical contradiction. Hence, due to the convergence of the series $\sum_{k=0}^k (\eta_{k+1} - \eta_k)^+$ and $\sum_{k=0}^k (\eta_{k+1} - \eta_k)^-$, taking $k \rightarrow +\infty$ in (3.8) yields $\lim_{k \rightarrow +\infty} \eta_k = \eta$. This completes the proof. $\square$

Lemma 3.3. *Let $\mathcal{N} : \mathcal{X} \rightarrow \mathcal{X}$ be an operator satisfying conditions $(\mathcal{N}1)$ – $(\mathcal{N}5)$. Suppose that $\{u_k\}$ is a sequence generated by Algorithms 1 and 2. For any $u^* \in VI(\mathcal{K}, \mathcal{N})$, we have*

$$\|p_k - u^*\|^2 \leq \|q_k - u^*\|^2 - \left(1 - \mu^2 \frac{\eta_k^2}{\eta_{k+1}^2}\right) \|q_k - y_k\|^2.$$

Proof. Consider the following expression

$$\begin{aligned} \|p_k - u^*\|^2 &= \|y_k + \eta_k[\mathcal{N}(u_k) - \mathcal{N}(y_k)] - u^*\|^2 \\ &= \|y_k - u^*\|^2 + \eta_k^2 \|\mathcal{N}(u_k) - \mathcal{N}(y_k)\|^2 \\ &\quad + 2\eta_k \langle y_k - u^*, \mathcal{N}(u_k) - \mathcal{N}(y_k) \rangle \\ &= \|y_k + u_k - u_k - u^*\|^2 + \eta_k^2 \|\mathcal{N}(u_k) - \mathcal{N}(y_k)\|^2 \\ &\quad + 2\eta_k \langle y_k - u^*, \mathcal{N}(u_k) - \mathcal{N}(y_k) \rangle \\ &= \|y_k - u_k\|^2 + \|u_k - u^*\|^2 + 2\langle y_k - u_k, u_k - u^* \rangle \\ &\quad + \eta_k^2 \|\mathcal{N}(u_k) - \mathcal{N}(y_k)\|^2 + 2\eta_k \langle y_k - u^*, \mathcal{N}(u_k) - \mathcal{N}(y_k) \rangle \\ &= \|u_k - u^*\|^2 + \|y_k - u_k\|^2 + 2\langle y_k - u_k, y_k - u^* \rangle \\ &\quad + 2\langle y_k - u_k, u_k - y_k \rangle + \eta_k^2 \|\mathcal{N}(u_k) - \mathcal{N}(y_k)\|^2 \\ &\quad + 2\eta_k \langle y_k - u^*, \mathcal{N}(u_k) - \mathcal{N}(y_k) \rangle. \end{aligned} \tag{3.9}$$

Additionally, we can express

$$\langle u_k - \eta_k \mathcal{N}(u_k) - y_k, y - y_k \rangle \leq 0, \quad \forall y \in \mathcal{K}. \tag{3.10}$$

For a given $u^* \in VI(\mathcal{K}, \mathcal{N})$, we can state

$$\langle u_k - y_k, u^* - y_k \rangle \leq \eta_k \langle \mathcal{N}(u_k), u^* - y_k \rangle. \tag{3.11}$$

By combining equations (3.9) and (3.11), we obtain

$$\begin{aligned}
\|p_k - u^*\|^2 &\leq \|u_k - u^*\|^2 + \|y_k - u_k\|^2 + 2\eta_k \langle \mathcal{N}(u_k), u^* - y_k \rangle \\
&\quad - 2\langle u_k - y_k, u_k - y_k \rangle + \eta_k^2 \|\mathcal{N}(u_k) - \mathcal{N}(y_k)\|^2 \\
&\quad - 2\eta_k \langle \mathcal{N}(u_k) - \mathcal{N}(y_k), u^* - y_k \rangle \\
&= \|u_k - u^*\|^2 - \|u_k - y_k\|^2 + \eta_k^2 \|\mathcal{N}(u_k) - \mathcal{N}(y_k)\|^2 \\
&\quad - 2\eta_k \langle \mathcal{N}(y_k), y_k - u^* \rangle. \tag{3.12}
\end{aligned}$$

Utilizing the concept of a mapping $\mathcal{N}$ on $\mathcal{K}$, we derive

$$\langle \mathcal{N}(u^*), y - u^* \rangle - \langle \mathcal{N}(y), y - u^* \rangle \leq 0, \quad \forall y \in \mathcal{K}.$$

Employing $u^* \in VI(\mathcal{K}, \mathcal{N})$, we obtain

$$\langle \mathcal{N}(y), y - u^* \rangle \geq 0, \quad \forall y \in \mathcal{K}.$$

Substituting $y = y_k \in \mathcal{K}$ yields

$$\langle \mathcal{N}(y_k), y_k - u^* \rangle \geq 0. \tag{3.13}$$

From equations (3.12) and (3.13), we deduce

$$\begin{aligned}
\|p_k - u^*\|^2 &\leq \|u_k - u^*\|^2 - \|u_k - y_k\|^2 + \mu^2 \frac{\eta_k^2}{\eta_{k+1}^2} \|u_k - y_k\|^2 \\
&= \|u_k - u^*\|^2 - \left(1 - \mu^2 \frac{\eta_k^2}{\eta_{k+1}^2}\right) \|u_k - y_k\|^2. \tag{3.14}
\end{aligned}$$

□

Theorem 3.4. *Let $\mathcal{N} : \mathcal{X} \rightarrow \mathcal{X}$ be an operator satisfying the conditions (N1)–(N5). Then, the sequence $\{u_k\}$ generated by Algorithms 1 and 2 converges strongly to $u^* \in VI(\mathcal{K}, \mathcal{N}) \cap \text{Fix}(\mathcal{Q})$, where $u^* = P_{VI(\mathcal{K}, \mathcal{N}) \cap \text{Fix}(\mathcal{Q})}(0)$.*

Proof. **Claim 1:** $\{u_k\}$ is a bounded sequence.

Let's consider that

$$u_{k+1} = (1 - \alpha_k)p_k + \alpha_k \mathcal{Q}(p_k).$$

By utilizing the definition of the sequence $\{u_{k+1}\}$, we have

$$\begin{aligned}
\|u_{k+1} - u^*\|^2 &= \|(1 - \alpha_k)p_k + \alpha_k \mathcal{Q}(p_k) - u^*\|^2 \\
&= \|p_k - u^*\|^2 + 2\alpha_k \langle p_k - u^*, \mathcal{Q}(p_k) - p_k \rangle + \alpha_k^2 \|\mathcal{Q}(p_k) - p_k\|^2 \\
&\leq \|p_k - u^*\|^2 + \alpha_k(\rho - 1) \|\mathcal{Q}(p_k) - p_k\|^2 + \alpha_k^2 \|\mathcal{Q}(p_k) - p_k\|^2 \\
&= \|p_k - u^*\|^2 - \alpha_k(1 - \rho - \alpha_k) \|\mathcal{Q}(p_k) - p_k\|^2. \tag{3.15}
\end{aligned}$$

By using the value of $\{q_k\}$, we obtain

$$\begin{aligned}
\|q_k - u^*\| &= \|u_k + \theta_k(u_k - u_{k-1}) - \varsigma_k u_k - \theta_k \varsigma_k(u_k - u_{k-1}) - u^*\| \\
&= \|(1 - \varsigma_k)(u_k - u^*) + (1 - \varsigma_k)\theta_k(u_k - u_{k-1}) - \varsigma_k u^*\| \\
&\leq (1 - \varsigma_k)\|u_k - u^*\| + (1 - \varsigma_k)\theta_k\|u_k - u_{k-1}\| + \varsigma_k\|u^*\| \\
&\leq (1 - \varsigma_k)\|u_k - u^*\| + \varsigma_k M_1,
\end{aligned} \tag{3.16}$$

for some fixed number M_1 , we have

$$(1 - \varsigma_k) \frac{\theta_k}{\varsigma_k} \|u_k - u_{k-1}\| + \|u^*\| \leq M_1.$$

By using $\eta_k \rightarrow \eta$ such that $\chi \in (0, 1 - \mu^2)$, we have

$$\lim_{k \rightarrow +\infty} \left(1 - \mu^2 \frac{\eta_k^2}{\eta_{k+1}^2} \right) = 1 - \mu^2 > \chi > 0.$$

Thus, there exists some fixed $k_0 \in \mathbb{N}$ such that

$$\left(1 - \mu^2 \frac{\eta_k^2}{\eta_{k+1}^2} \right) > \chi > 0, \quad \forall k \geq k_0. \tag{3.17}$$

By using Lemma 3.3, we can rewrite

$$\|p_k - u^*\|^2 \leq \|q_k - u^*\|^2, \quad \forall k \geq k_0. \tag{3.18}$$

From expressions (3.15), (3.16), and (3.18), we infer that

$$\|u_{k+1} - u^*\| \leq (1 - \varsigma_k)\|u_k - u^*\| + \varsigma_k M_1 - \alpha_k(1 - \rho - \alpha_k)\|\mathcal{Q}(p_k) - p_k\|^2. \tag{3.19}$$

Thus, for $\{\alpha_k\} \subset (0, 1 - \rho)$, we obtain

$$\begin{aligned}
\|u_{k+1} - u^*\| &\leq (1 - \varsigma_k)\|u_k - u^*\| + \varsigma_k M_1 \\
&\leq \max\{\|u_k - u^*\|, M_1\} \\
&\vdots \\
&\leq \max\{\|u_{k_0} - u^*\|, M_1\}.
\end{aligned} \tag{3.20}$$

Consequently, we may infer that the sequence $\{u_k\}$ is a bounded sequence.

Claim 2:

$$\begin{aligned}
&\left(1 - \mu^2 \frac{\eta_k^2}{\eta_{k+1}^2} \right) \|q_k - y_k\|^2 + \alpha_k(1 - \rho - \alpha_k) \|\mathcal{Q}(p_k) - p_k\|^2 \\
&\leq \|u_k - u^*\|^2 - \|u_{k+1} - u^*\|^2 + \varsigma_k M_2,
\end{aligned} \tag{3.21}$$

for some fixed $M_2 > 0$. Indeed, by using the definition of $\{u_{k+1}\}$, we have

$$\begin{aligned}
\|u_{k+1} - u^*\|^2 &= \|(1 - \alpha_k)p_k + \alpha_k \mathcal{Q}(p_k) - u^*\|^2 \\
&= \|p_k - u^*\|^2 + 2\alpha_k \langle p_k - u^*, \mathcal{Q}(p_k) - p_k \rangle + \alpha_k^2 \|\mathcal{Q}(p_k) - p_k\|^2 \\
&\leq \|p_k - u^*\|^2 + \alpha_k(\rho - 1) \|\mathcal{Q}(p_k) - p_k\|^2 + \alpha_k^2 \|\mathcal{Q}(p_k) - p_k\|^2 \\
&= \|p_k - u^*\|^2 - \alpha_k(1 - \rho - \alpha_k) \|\mathcal{Q}(p_k) - p_k\|^2.
\end{aligned} \tag{3.22}$$

By using Lemma 3.3, we obtain

$$\|p_k - u^*\|^2 \leq \|q_k - u^*\|^2 - \left(1 - \mu^2 \frac{\eta_k^2}{\eta_{k+1}^2}\right) \|q_k - y_k\|^2. \tag{3.23}$$

By using expression (3.16), we can obtain

$$\begin{aligned}
\|q_k - u^*\|^2 &\leq (1 - \varsigma_k)^2 \|u_k - u^*\|^2 + \varsigma_k^2 M_1^2 + 2M_1 \varsigma_k (1 - \varsigma_k) \|u_k - u^*\| \\
&\leq \|u_k - u^*\|^2 + \varsigma_k [\varsigma_k M_1^2 + 2M_1 (1 - \varsigma_k) \|u_k - u^*\|] \\
&\leq \|u_k - u^*\|^2 + \varsigma_k M_2,
\end{aligned} \tag{3.24}$$

where M_2 is some fixed constant > 0 . From expressions (3.22), (3.23) and (3.24), we obtain

$$\begin{aligned}
\|u_{k+1} - u^*\|^2 &\leq \|u_k - u^*\|^2 + \varsigma_k M_2 - \alpha_k(1 - \rho - \alpha_k) \|\mathcal{Q}(p_k) - p_k\|^2 \\
&\quad - \left(1 - \mu^2 \frac{\eta_k^2}{\eta_{k+1}^2}\right) \|q_k - y_k\|^2.
\end{aligned} \tag{3.25}$$

Claim 3:

Using the value of $\{q_k\}$, we can express as follows

$$\begin{aligned}
\|q_k - u^*\|^2 &= \|u_k + \theta_k(u_k - u_{k-1}) - \varsigma_k u_k - \theta_k \varsigma_k(u_k - u_{k-1}) - u^*\|^2 \\
&= \|(1 - \varsigma_k)(u_k - u^*) + (1 - \varsigma_k)\theta_k(u_k - u_{k-1}) - \varsigma_k u^*\|^2 \\
&\leq \|(1 - \varsigma_k)(u_k - u^*) + (1 - \varsigma_k)\theta_k(u_k - u_{k-1})\|^2 + 2\varsigma_k \langle -u^*, q_k - u^* \rangle \\
&= (1 - \varsigma_k)^2 \|u_k - u^*\|^2 + (1 - \varsigma_k)^2 \theta_k^2 \|u_k - u_{k-1}\|^2 \\
&\quad + 2\theta_k(1 - \varsigma_k)^2 \|u_k - u^*\| \|u_k - u_{k-1}\| \\
&\quad + 2\varsigma_k \langle -u^*, q_k - u_{k+1} \rangle + 2\varsigma_k \langle -u^*, u_{k+1} - u^* \rangle
\end{aligned}$$

$$\begin{aligned}
&\leq (1 - \varsigma_k) \|u_k - u^*\|^2 + \theta_k^2 \|u_k - u_{k-1}\|^2 \\
&\quad + 2\theta_k(1 - \varsigma_k) \|u_k - u^*\| \|u_k - u_{k-1}\| \\
&\quad + 2\varsigma_k \|u^*\| \|q_k - u_{k+1}\| + 2\varsigma_k \langle -u^*, u_{k+1} - u^* \rangle \\
&= (1 - \varsigma_k) \|u_k - u^*\|^2 + \varsigma_k \left[\theta_k \|u_k - u_{k-1}\| \frac{\theta_k}{\varsigma_k} \|u_k - u_{k-1}\| \right. \\
&\quad + 2(1 - \varsigma_k) \|u_k - u^*\| \frac{\theta_k}{\varsigma_k} \|u_k - u_{k-1}\| \\
&\quad \left. + 2\|u^*\| \|q_k - u_{k+1}\| + 2\langle u^*, u^* - u_{k+1} \rangle \right]. \tag{3.26}
\end{aligned}$$

Combining equations (3.18) and (3.26), we obtain

$$\begin{aligned}
\|u_{k+1} - u^*\|^2 &\leq (1 - \varsigma_k) \|u_k - u^*\|^2 + \varsigma_k \left[\theta_k \|u_k - u_{k-1}\| \frac{\theta_k}{\varsigma_k} \|u_k - u_{k-1}\| \right. \\
&\quad + 2(1 - \varsigma_k) \|u_k - u^*\| \frac{\theta_k}{\varsigma_k} \|u_k - u_{k-1}\| \\
&\quad \left. + 2\|u^*\| \|q_k - u_{k+1}\| + 2\langle u^*, u^* - u_{k+1} \rangle \right]. \tag{3.27}
\end{aligned}$$

Claim 4: The sequence $\|u_k - u^*\|^2$ converges to zero.

Let's define

$$c_k := \|u_k - u^*\|^2$$

and

$$\begin{aligned}
e_k &:= \theta_k \|u_k - u_{k-1}\| \frac{\theta_k}{\varsigma_k} \|u_k - u_{k-1}\| + 2(1 - \varsigma_k) \|u_k - u^*\| \frac{\theta_k}{\varsigma_k} \|u_k - u_{k-1}\| \\
&\quad + 2\|u^*\| \|q_k - u_{k+1}\| + 2\langle u^*, u^* - u_{k+1} \rangle.
\end{aligned}$$

Then, Claim 4 can be restated as

$$c_{k+1} \leq (1 - \varsigma_k) c_k + \varsigma_k e_k.$$

By Lemma 2.5, it's enough to prove that $\limsup_{j \rightarrow +\infty} e_{k_j} \leq 0$ for $\{c_{k_j}\}$ in $\{c_k\}$ such that

$$\liminf_{j \rightarrow +\infty} (c_{k_j+1} - c_{k_j}) \geq 0.$$

This is equivalent to stating

$$\limsup_{j \rightarrow +\infty} \langle u^*, u^* - u_{k_j+1} \rangle \leq 0$$

and

$$\limsup_{j \rightarrow +\infty} \|q_{k_j} - u_{k_j+1}\| \leq 0,$$

one from each subsequence $\{\|u_{k_j} - u^*\|\}$ of $\{\|u_k - u^*\|\}$ following

$$\liminf_{j \rightarrow +\infty} (\|u_{k_j+1} - u^*\| - \|u_{k_j} - u^*\|) \geq 0.$$

Suppose $\{\|u_{k_j} - u^*\|\}$ is such a subsequence of $\{\|u_k - u^*\|\}$ satisfying

$$\liminf_{j \rightarrow +\infty} (\|u_{k_j+1} - u^*\| - \|u_{k_j} - u^*\|) \geq 0.$$

Then, we have

$$\begin{aligned} & \liminf_{j \rightarrow +\infty} (\|u_{k_j+1} - u^*\|^2 - \|u_{k_j} - u^*\|^2) \\ &= \liminf_{j \rightarrow +\infty} (\|u_{k_j+1} - u^*\| - \|u_{k_j} - u^*\|)(\|u_{k_j+1} - u^*\| + \|u_{k_j} - u^*\|) \\ &\geq 0. \end{aligned} \tag{3.28}$$

As a result of Claim 2,

$$\begin{aligned} & \limsup_{j \rightarrow +\infty} \left[\left(1 - \frac{\mu^2 \eta_{k_j}^2}{\eta_{k_j+1}^2}\right) \|q_{k_j} - y_{k_j}\|^2 + \alpha_{k_j} (1 - \rho - \alpha_{k_j}) \|\mathcal{Q}(p_{k_j}) - p_{k_j}\|^2 \right] \\ &\leq \limsup_{j \rightarrow +\infty} [\|u_{k_j} - u^*\|^2 - \|u_{k_j+1} - u^*\|^2] + \limsup_{j \rightarrow +\infty} \varsigma_{k_j} K_2 \\ &= -\liminf_{j \rightarrow +\infty} [\|u_{k_j+1} - u^*\|^2 - \|u_{k_j} - u^*\|^2] \\ &\leq 0. \end{aligned} \tag{3.29}$$

The above expressions imply the following mathematical relationships

$$\lim_{j \rightarrow +\infty} \|q_{k_j} - y_{k_j}\| = 0, \quad \lim_{j \rightarrow +\infty} \|\mathcal{Q}(p_{k_j}) - p_{k_j}\| = 0. \tag{3.30}$$

Consequently, we have

$$\|p_{k_j} - y_{k_j}\| = \|y_{k_j} + \eta_{k_j} [\mathcal{N}(q_{k_j}) - \mathcal{N}(y_{k_j})] - y_{k_j}\| \leq \eta_{k_j} L \|q_{k_j} - y_{k_j}\|. \tag{3.31}$$

Thus, it follows that

$$\lim_{j \rightarrow +\infty} \|p_{k_j} - y_{k_j}\| = 0. \tag{3.32}$$

Hence, we can deduce

$$\lim_{j \rightarrow +\infty} \|p_{k_j} - q_{k_j}\| = 0. \tag{3.33}$$

Moreover, it's straightforward to derive

$$\lim_{j \rightarrow +\infty} \|q_{k_j} - u_{k_j}\| = 0. \tag{3.34}$$

Combining equations (3.33) and (3.34) leads to

$$\lim_{j \rightarrow +\infty} \|p_{k_j} - u_{k_j}\| = 0. \tag{3.35}$$

Given the equation $u_{k_j+1} = (1 - \alpha_{k_j})p_{k_j} + \alpha_{k_j}\mathcal{Q}(p_{k_j})$, we observe:

$$\lim_{j \rightarrow +\infty} \|u_{k_j+1} - p_{k_j}\| = \alpha_{k_j} \|\mathcal{Q}(p_{k_j}) - p_{k_j}\| \leq (1 - \rho) \|\mathcal{Q}(p_{k_j}) - p_{k_j}\|. \quad (3.36)$$

Thus, it follows that

$$\lim_{j \rightarrow +\infty} \|u_{k_j+1} - p_{k_j}\| = 0. \quad (3.37)$$

The expressions above imply that

$$\lim_{j \rightarrow +\infty} \|u_{k_j} - u_{k_j+1}\| \leq \lim_{j \rightarrow +\infty} \|u_{k_j} - p_{k_j}\| + \lim_{j \rightarrow +\infty} \|p_{k_j} - u_{k_j+1}\| = 0 \quad (3.38)$$

and

$$\lim_{j \rightarrow +\infty} \|q_{k_j} - u_{k_j+1}\| \leq \lim_{j \rightarrow +\infty} \|q_{k_j} - p_{k_j}\| + \lim_{j \rightarrow +\infty} \|p_{k_j} - u_{k_j+1}\| = 0. \quad (3.39)$$

Given that the sequence $\{u_{k_j}\}$ is bounded, without loss of generality, we assume that $\{u_{k_j}\}$ converges weakly to some $\hat{u} \in \mathcal{X}$.

Next, we need to prove that $\hat{u} \in VI(\mathcal{K}, \mathcal{N})$. Since $\{q_{k_j}\}$ converges weakly to $\hat{u}$ and because $\lim_{j \rightarrow +\infty} \|q_{k_j} - y_{k_j}\| = 0$, the sequence $\{y_{k_j}\}$ also converges weakly to $\hat{u}$. We then need to prove that $\hat{u} \in VI(\mathcal{K}, \mathcal{N})$. This implies

$$y_{k_j} = P_{\mathcal{X}}[q_{k_j} - \eta_{k_j}\mathcal{N}(q_{k_j})],$$

which is equivalent to

$$\langle q_{k_j} - \eta_{k_j}\mathcal{N}(q_{k_j}) - y_{k_j}, y - y_{k_j} \rangle \leq 0, \quad \forall y \in \mathcal{X}. \quad (3.40)$$

From the inequality above, we have

$$\langle q_{k_j} - y_{k_j}, y - y_{k_j} \rangle \leq \eta_{k_j} \langle \mathcal{N}(q_{k_j}), y - y_{k_j} \rangle, \quad \forall y \in \mathcal{X}. \quad (3.41)$$

Consequently, we obtain

$$\frac{1}{\eta_{k_j}} \langle q_{k_j} - y_{k_j}, y - y_{k_j} \rangle + \langle \mathcal{N}(q_{k_j}), y_{k_j} - q_{k_j} \rangle \leq \langle \mathcal{N}(q_{k_j}), y - q_{k_j} \rangle, \quad \forall y \in \mathcal{X}. \quad (3.42)$$

Given that $\min\{\frac{\mu}{L}, \eta_1\} \leq \eta \leq \eta_1$ and $\{q_{k_j}\}$ forms a bounded sequence, utilizing the limits $\lim_{j \rightarrow +\infty} \|q_{k_j} - y_{k_j}\| = 0$ and $j \rightarrow +\infty$ in (3.42), we deduce

$$\liminf_{j \rightarrow +\infty} \langle \mathcal{N}(q_{k_j}), y - q_{k_j} \rangle \geq 0, \quad \forall y \in \mathcal{X}. \quad (3.43)$$

Moreover, it implies

$$\begin{aligned} \langle \mathcal{N}(y_{k_j}), y - y_{k_j} \rangle &= \langle \mathcal{N}(y_{k_j}) - \mathcal{N}(q_{k_j}), y - q_{k_j} \rangle \\ &\quad + \langle \mathcal{N}(q_{k_j}), y - q_{k_j} \rangle + \langle \mathcal{N}(y_{k_j}), q_{k_j} - y_{k_j} \rangle. \end{aligned} \quad (3.44)$$

As $\lim_{j \rightarrow +\infty} \|q_{k_j} - y_{k_j}\| = 0$ and under the Lipschitz condition on the mapping $\mathcal{N}$, we obtain

$$\lim_{j \rightarrow +\infty} \|\mathcal{N}(q_{k_j}) - \mathcal{N}(y_{k_j})\| = 0, \quad (3.45)$$

which, coupled with (3.44) and (3.45), yields

$$\liminf_{j \rightarrow +\infty} \langle \mathcal{N}(y_{k_j}), y - y_{k_j} \rangle \geq 0, \quad \forall y \in \mathcal{X}. \quad (3.46)$$

For further proof, let's consider a positive sequence $\{\epsilon_j\}$ that converges to zero and decreases. For each $\{\epsilon_j\}$, there exists a least positive integer denoted by m_j such that

$$\langle \mathcal{N}(q_{k_i}), y - q_{k_i} \rangle + \epsilon_j > 0, \quad \forall i \geq m_j, \quad (3.47)$$

where the existence of m_j follows from expression (3.46). As $\{\epsilon_j\}$ is decreasing, the sequence m_j is evidently increasing. If there exists a natural number $N_0 \in \mathbb{N}$ such that $\mathcal{N}(u_{n_{m_k}}) \neq 0$ for all $n_{m_k} \geq N_0$, then we consider

$$\tilde{h}_{k_{m_n}} = \frac{\mathcal{N}(q_{k_{m_n}})}{\|\mathcal{N}(q_{k_{m_n}})\|^2}, \quad \forall k_{m_n} \geq N_0. \quad (3.48)$$

Using the provided value of $\tilde{h}_{k_{m_n}}$, we derive the following equation

$$\langle \mathcal{N}(q_{k_{m_n}}), \tilde{h}_{k_{m_n}} \rangle = 1, \quad \forall k_{m_n} \geq N_0. \quad (3.49)$$

Combining equations (3.47) and (3.49), we obtain

$$\langle \mathcal{N}(q_{k_{m_n}}), y + \epsilon_k \tilde{h}_{k_{m_n}} - q_{k_{m_n}} \rangle > 0. \quad (3.50)$$

Utilizing the definition of the pseudomonotone mapping $\mathcal{N}$, we express

$$\langle \mathcal{N}(y + \epsilon_k \tilde{h}_{k_{m_n}}), y + \epsilon_k \tilde{h}_{k_{m_n}} - q_{k_{m_n}} \rangle > 0. \quad (3.51)$$

For all $k_{m_n} \geq N_0$, we have

$$\begin{aligned} \langle \mathcal{N}(y), y - q_{k_{m_n}} \rangle &\geq \langle \mathcal{N}(y) - \mathcal{N}(y + \epsilon_k \tilde{h}_{k_{m_n}}), y + \epsilon_k \tilde{h}_{k_{m_n}} - q_{k_{m_n}} \rangle \\ &\quad - \epsilon_k \langle \mathcal{N}(y), \tilde{h}_{k_{m_n}} \rangle. \end{aligned} \quad (3.52)$$

Given that the sequence $\{q_{k_n}\}$ weakly converges to $\hat{u} \in \mathcal{X}$, it follows that $\{\mathcal{N}(q_{k_n})\}$ weakly converges to $\mathcal{N}(\hat{u})$. Suppose $\mathcal{N}(\hat{u}) \neq 0$, implying

$$\|\mathcal{N}(\hat{u})\| \leq \liminf_{n \rightarrow +\infty} \|\mathcal{N}(q_{k_n})\|. \quad (3.53)$$

Since $\{q_{k_{m_n}}\} \subset \{q_{k_n}\}$ and $\lim_{k \rightarrow +\infty} \epsilon_k = 0$, we have

$$0 \leq \lim_{n \rightarrow +\infty} \|\epsilon_k \tilde{h}_{k_{m_n}}\| = \lim_{n \rightarrow +\infty} \frac{\epsilon_k}{\|\mathcal{N}(q_{k_{m_n}})\|} \leq \frac{0}{\|\mathcal{N}(\hat{u})\|} = 0. \quad (3.54)$$

By letting $n \rightarrow +\infty$ in equation (3.52), we obtain

$$\langle \mathcal{N}(y), y - \hat{u} \rangle \geq 0, \quad \forall y \in \mathcal{X}. \quad (3.55)$$

Let $u \in \mathcal{X}$ be an arbitrary element and $0 < \psi \leq 1$. Consider the expression

$$\hat{u}_\psi = \psi u + (1 - \psi)\hat{u}.$$

Then, $\hat{u}_\psi \in \mathcal{X}$. From Equation (3.55), we have

$$\psi \langle \mathcal{N}(\hat{u}_\psi), u - \hat{u} \rangle \geq 0.$$

Thus,

$$\langle \mathcal{N}(\hat{u}_\psi), u - \hat{u} \rangle \geq 0.$$

As $\psi \rightarrow 0$, $\hat{u}_\psi \rightarrow \hat{u}$ along a line segment. By the continuity of the operator, $\mathcal{N}(\hat{u}_\psi)$ converges to $\mathcal{N}(\hat{u})$ as $\psi \rightarrow 0$. Hence, we have:

$$\langle \mathcal{N}(\hat{u}), u - \hat{u} \rangle \geq 0.$$

Therefore, $\hat{u}$ is a solution of problem (VIP). Given $u^* = P_{VI(\mathcal{K}, \mathcal{N}) \cap \text{Fix}(\mathcal{Q})}(0)$, we have:

$$\langle 0 - u^*, y - u^* \rangle \leq 0, \quad \forall y \in VI(\mathcal{K}, \mathcal{N}) \cap \text{Fix}(\mathcal{Q}).$$

From Equation (3.34), it follows that $\{q_{k_j}\}$ converges weakly to $\hat{u} \in \mathcal{X}$. Similarly, from Equation (3.35), $\{p_{k_j}\}$ converges weakly to $\hat{u} \in \mathcal{X}$. By the demiclosedness of $(I - \mathcal{Q})$, we conclude $\hat{u} \in \text{Fix}(\mathcal{Q})$. Therefore, $\hat{u} \in VI(\mathcal{K}, \mathcal{N}) \cap \text{Fix}(\mathcal{Q})$, leading to:

$$\lim_{j \rightarrow +\infty} \langle u^*, u^* - u_{k_j} \rangle = \langle u^*, u^* - \hat{u} \rangle \leq 0.$$

Furthermore, since $\lim_{j \rightarrow +\infty} \|u_{k_j+1} - u_{k_j}\| = 0$, we have

$$\limsup_{j \rightarrow +\infty} \langle u^*, u^* - u_{k_j+1} \rangle \leq \limsup_{j \rightarrow +\infty} \langle u^*, u^* - u_{k_j} \rangle + \limsup_{j \rightarrow +\infty} \langle u^*, u_{k_j} - u_{k_j+1} \rangle \leq 0.$$

Combining Claim 3 and Lemma 2.5, we observe that $u_k \rightarrow u^*$ as $k \rightarrow +\infty$. This completes the proof. $\square$

4. NUMERICAL ILLUSTRATIONS

This section investigates the algorithmic implications of the provided methodologies, as well as explores how variations in control parameters affect the numerical efficacy of the suggested algorithms.

Example 4.1. Consider a mapping $\mathcal{N} : \mathbb{R}^m \rightarrow \mathbb{R}^m$ defined as

$$\mathcal{N}(u) = Mu,$$

where $q = 0$. Here, $M = NN^T + B + D$, with $N = \text{rand}(m)$ and $K = \text{rand}(m)$ chosen randomly, and $B = 0.5K - 0.5K^T$, $D = \text{diag}(\text{rand}(m, 1))$. The feasible set $\mathcal{K}$ is defined as

$$\mathcal{K} = \{u \in \mathbb{R}^m : -10 \leq u_i \leq 10\}.$$

It is evident that the mapping $\mathcal{N}$ is monotone and Lipschitz continuous with constant $L = \|M\|$. Additionally, a function $\mathcal{Q} : \mathcal{X} \rightarrow \mathcal{X}$ is defined as

$$\mathcal{Q}(u) = \frac{1}{2}u.$$

The initial points for the tests are $u_0 = u_1 = (2, 2, \dots, 2)$. Different dimensions of the Hilbert space are considered to study its behavior in higher dimensions. The stopping condition for these experiments is $D_k = \|q_k - y_k\| \leq 10^{-10}$.

Figures 1 to 6 and Tables 1 to 2 depict empirical observations for Example 2. The following control criteria are enforced:

(1) Algorithm 1 (**Algo1**):

$$\begin{aligned} \eta_1 &= 0.43, \quad \theta = 0.56, \quad \mu = 0.64, \\ \chi_k &= \frac{10}{(1+k)^2}, \quad s_k = \frac{1}{(3k+5)}, \quad \alpha_k = \frac{2k}{(3k+2)}; \end{aligned}$$

(2) Algorithm 2 (**Algo2**):

$$\begin{aligned} \eta_1 &= 0.43, \quad \theta = 0.56, \quad \mu = 0.64, \\ \chi_k &= \frac{10}{(1+k)^2}, \quad s_k = \frac{1}{(3k+5)}, \quad \alpha_k = \frac{2k}{(3k+2)}; \end{aligned}$$

(3) Algorithm 1 in [32] (**Algo3.1**):

$$\begin{aligned} \tau_1 &= 0.43, \quad \theta = 0.56, \quad \mu = 0.64, \\ \chi_k &= \frac{10}{(1+k)^2}, \quad s_k = \frac{1}{(3k+5)}, \quad \alpha_k = \frac{2k}{(3k+2)}, \\ f(u) &= \frac{u}{3}; \end{aligned}$$

(4) Algorithm 2 in [32] (**Algo3.2**):

$$\begin{aligned} \tau_1 &= 0.43, \quad \theta = 0.56, \quad \mu = 0.64, \\ \chi_k &= \frac{10}{(1+k)^2}, \quad s_k = \frac{1}{(3k+5)}, \quad \alpha_k = \frac{k}{(3k+2)}, \\ f(u) &= \frac{u}{3}. \end{aligned}$$




FIGURE 1. Numerical comparison of Algorithm 1 and Algorithm 2 with Algorithm 1 in [32], Algorithm 2 in [32] when $m = 5$.




FIGURE 2. Numerical comparison of Algorithm 1 and Algorithm 2 with Algorithm 1 in [32], Algorithm 2 in [32] when $m = 5$.




FIGURE 3. Numerical comparison of Algorithm 1 and Algorithm 2 with Algorithm 1 in [32], Algorithm 2 in [32] when $m = 10$.




FIGURE 4. Numerical comparison of Algorithm 1 and Algorithm 2 with Algorithm 1 in [32], Algorithm 2 in [32] when $m = 10$.




FIGURE 5. Numerical comparison of Algorithm 1 and Algorithm 2 with Algorithm 1 in [32], Algorithm 2 in [32] when $m = 20$.




FIGURE 6. Numerical comparison of Algorithm 1 and Algorithm 2 with Algorithm 1 in [32], Algorithm 2 in [32] when $m = 20$.

TABLE 1. Numerical values for Figures 1–6.

The number of iterations				
m	Algo3.1	Algo3.2	Algo1	Algo2
5	54	46	27	21
10	59	49	42	23
20	61	51	44	25
50	67	56	45	34
100	73	61	48	36

TABLE 2. Numerical values for Figures 1–6.

Time required to complete				
m	Algo3.1	Algo3.2	Algo1	Algo2
5	0.41096080	0.33812320	0.20280990	0.16518980
10	0.64628270	0.57689310	0.49204840	0.29402830
20	0.48693460	0.36085080	0.30036720	0.26893690
50	0.57369690	0.40373730	0.37593220	0.29847481
100	0.67855063	0.53929280	0.41928400	0.30193762

Example 4.2. Let $\mathcal{K}$ denote the feasible set defined as follows:

$$\mathcal{K} := \{u \in L^2([0, 1]) : \|u\|_{L^2} \leq 1\}.$$

We define an operator $\mathcal{N} : \mathcal{K} \rightarrow \mathcal{X}$ by

$$\mathcal{N}(u)(t) = \max \{u(t), 0\} = \frac{u(t) + |u(t)|}{2}.$$

Here, $\mathcal{X} = L^2([0, 1])$ represents a real Hilbert space with its inner product and norm given by

$$\langle u, y \rangle = \int_0^1 u(t)y(t)dt, \quad \forall u, y \in \mathcal{X}$$

and

$$\|u\|_{L^2} = \sqrt{\int_0^1 |u(t)|^2 dt}.$$

It can be readily verified that $\mathcal{N}$ is monotone and 1-Lipschitz continuous. The projection onto $\mathcal{K}$ is straightforward

$$P_{\mathcal{K}}(u) = \begin{cases} \frac{u}{\|u\|_{L^2}} & \text{if } \|u\|_{L^2} > 1, \\ u & \text{if } \|u\|_{L^2} \leq 1. \end{cases}$$

A mapping $\mathcal{Q} : L^2([0, 1]) \rightarrow L^2([0, 1])$ takes the form

$$\mathcal{Q}(u)(t) = \int_0^1 tu(s)ds, \quad t \in [0, 1].$$

A simple analysis shows that $\mathcal{Q}$ is 0-demicontractive, with the solution being $u^*(t) = 0$. These trials commence with a halting requirement $D_k = \|q_k - y_k\|_{L^2} \leq 10^{-6}$. The tables 3 and 4 present numerical results pertaining to Example 4.2. The following conditions serve as control criteria:

(1) Algorithm 1 (**Algo1**):

$$\begin{aligned} \eta_1 &= 0.43, \quad \theta = 0.56, \quad \mu = 0.64, \\ \chi_k &= \frac{10}{(1+k)^2}, \quad \varsigma_k = \frac{1}{(3k+5)}, \quad \alpha_k = \frac{2k}{(3k+2)}; \end{aligned}$$

(2) Algorithm 2 (**Algo2**):

$$\begin{aligned} \eta_1 &= 0.43, \quad \theta = 0.56, \quad \mu = 0.64, \\ \chi_k &= \frac{10}{(1+k)^2}, \quad \varsigma_k = \frac{1}{(3k+5)}, \quad \alpha_k = \frac{2k}{(3k+2)}; \end{aligned}$$

(3) Algorithm 1 in [32] (**Algo3.1**):

$$\begin{aligned} \tau_1 &= 0.43, \quad \theta = 0.56, \quad \mu = 0.64, \\ \chi_k &= \frac{10}{(1+k)^2}, \quad \varsigma_k = \frac{1}{(3k+5)}, \quad \alpha_k = \frac{2k}{(3k+2)}, \\ f(u) &= \frac{u}{3}; \end{aligned}$$

(4) Algorithm 2 in [32] (**Algo3.2**):

$$\begin{aligned} \tau_1 &= 0.43, \quad \theta = 0.56, \quad \mu = 0.64, \\ \chi_k &= \frac{10}{(1+k)^2}, \quad \varsigma_k = \frac{1}{(3k+5)}, \quad \alpha_k = \frac{k}{(3k+2)}, \\ f(u) &= \frac{u}{3}. \end{aligned}$$

TABLE 3. Numerical values for Example 4.2.

Total number of iterations				
$u_0 = u_1$	Algo3.1	Algo3.2	Algo1	Algo2
t^3	76	65	45	32
$t \cos(t)$	87	68	49	33
$t \exp(t)$	92	74	57	41
$t2^t$	98	79	61	43

TABLE 4. Numerical values for Example 4.2.

Required CPU time				
$u_0 = u_1$	Algo3.1	Algo3.2	Algo1	Algo2
t^3	1.1745382	1.1275749	0.5867949	0.3525294
$t \cos(t)$	1.1976944	1.4869759	0.8563924	0.6273644
$t \exp(t)$	2.0575325	1.5486922	0.9949494	0.7264347
$t2^t$	2.0025344	1.69576969	1.2207376	1.03811294

5. CONCLUSION

The paper introduces two explicit extragradient-like algorithms designed to address an equilibrium problem within a real Hilbert space, featuring a pseudomonotone and Lipschitz-type bifunction, constrained by fixed points. A novel stepsize rule is proposed, independent of Lipschitz-type constants. Various experiments are conducted to demonstrate the numerical performance of the proposed algorithms and compare them with established approaches in the literature.

Acknowledgements: The first author would like to thank Faculty of Science and Technology and Research and Development Institute, Valaya Alongkorn Rajabhat University under the Royal Patronage Pathun Thani Province. The second author was supported by University of Phayao and Thailand Science Research and Innovation Fund (Fundamental Fund 2024). The fourth author would like to thank Professor Dr. Poom Kumam from King Mongkuts University of Technology Thonburi, Thailand for his advice and comments to improve the results of this paper. This research (Grant No. RGNS 65-168) was supported by Office of the Permanent Secretary, Ministry of Higher Education, Science, Research and Innovation (OPS MHESI), Thailand Science Research and Innovation (TSRI) and Phetchabun Rajabhat University.

REFERENCES

- [1] H.A. Abass, O.K. Narain and O.M. Onifade, *Inertial extrapolation method for solving systems of monotone variational inclusion and fixed point problems using Bregman distance approach*, Nonlinear Funct. Anal. Appl., **28**(2) (2023), 497–520.
- [2] K. Abuasbeh, A. Kanwal, R. Shafqat, B. Taufeeq, M.A. Almulla and M. Awadalla, *A method for solving time-fractional initial boundary value problems of variable order*, Symmetry, **15**(2) (2023), Article ID 519.
- [3] K. Abuasbeh and R. Shafqat, *Fractional brownian motion for a system of fuzzy fractional stochastic differential equation*, J. Math., **2022** (2022), Article ID 3559035.
- [4] N.T. An, N.M. Nam and X. Qin, *Solving k -center problems involving sets based on optimization techniques*, J. Global Optim., **76**(1) (2019), 189–209.
- [5] A.S. Antipin, *On a method for convex programs using a symmetrical modification of the Lagrange function*, Ekonomika i Matem. Metody, **12** (1976), 1164–1173.
- [6] L.C. Ceng, A. Petruşel, X. Qin and J.C. Yao, *A modified inertial subgradient extragradient method for solving pseudomonotone variational inequalities and common fixed point problems*, Fixed Point Theory, **21**(1) (2020), 93–108.
- [7] L.C. Ceng, J.C. Yao and Y. Shehu, *On mann implicit composite subgradient extragradient methods for general systems of variational inequalities with hierarchical variational inequality constraints*, J. Ineq. Appl., **2022** (2022), Article ID 78.
- [8] C.M. Elliott, *Variational and quasivariational inequalities: applications to free-boundary problems (Claudio Baiocchi and antônio Capelo)*, SIAM Rev., **29**(2) (1987), 314–315.
- [9] H.A. Hammad, H. Rehman and M.D. la Sen, *Advanced algorithms and common solutions to variational inequalities*, Symmetry, **12** (2020), Article ID 1198.
- [10] H. Iiduka and I. Yamada, *A subgradient-type method for the equilibrium problem over the fixed point set and its applications*, Optimization, **58**(2) (2009), 251–261.
- [11] G. Kassay, J. Kolumbán and Z. Páles, *On nash stationary points*, Publ. Math. Debrecen, **54**(3-4) (1999), 267–279.
- [12] G. Kassay, J. Kolumbán and Z. Páles, *Factorization of minty and stampacchia variational inequality systems*, Euro. J. Oper. Res., **143** (2002), 377–389.
- [13] A. Khan, R. Shafqat and A.U.K. Niazi, *Existence results of fuzzy delay impulsive fractional differential equation by fixed point theory approach*, J. Funct. Spaces, **2022** (2022), Article ID 4123949.
- [14] J.K. Kim, A.H. Dar and Salahuddin, *Existence theorems for the generalized relaxed pseudomonotone variational inequalities*, Nonlinear Funct. Anal. Appl., **25**(1) (2020), 25–34.
- [15] D. Kinderlehrer and G. Stampacchia, *An introduction to variational inequalities and their applications*, Classic in Applied Mathematics, Society for Industrial and Applied Mathematics, (SIAM), Philadelphia, PA, 2000.
- [16] I. Konnov, *Equilibrium models and variational inequalities*, **210**, Elsevier, 2007.
- [17] G. Korpelevich, *The extragradient method for finding saddle points and other problems*, Ékonom. i Mat. Metody, **12**(4) (1976), 747–756.
- [18] R. Kraikaew and S. Saejung, *Strong convergence of the Halpern subgradient extragradient method for solving variational inequalities in Hilbert spaces*, J. Optim. Theory Appl., **163**(2) (2014), 399–412.
- [19] G.M. Lee and J.H. Lee, *On Augmented Lagrangian Methods of Multipliers and Alternating Direction Methods of Multipliers for Matrix Optimization Problems*, Nonlinear Funct. Anal. Appl., **27** (4) (2022), 869–879.

- [20] P.E. Maingé, *Strong convergence of projected subgradient methods for nonsmooth and nonstrictly convex minimization*, Set-Valued Anal., **16** (2008), 899–912.
- [21] P.E. Maingé and A. Moudafi, *Coupling viscosity methods with the extragradient algorithm for solving equilibrium problems*, J. Nonlinear Convex Anal., **9**(2) (2008), 283–294.
- [22] A. Moudafi, *Viscosity approximation methods for fixed-points problems*, J. Math. Anal. Appl., **241** (2000), 46–55.
- [23] A. Moumen, R. Shafqat, Z. Hammouch, A.U.K. Niazi and M.B. Jeelani, *Stability results for fractional integral pantograph differential equations involving two caputo operators*, AIMS Math., **8**(3) (2023), 6009–6025.
- [24] K. Muangchoo, *A Viscosity type Projection Method for Solving Pseudomonotone Variational Inequalities*, Nonlinear Funct. Anal. Appl., **26**(2) (2021), 347–371.
- [25] A. Nagurney, *Network economics: a variational inequality approach*, Kluwer Academic Publishers Group, Dordrecht, 1993, <https://doi.org/10.1007/978-94-011-2178-1>.
- [26] M.A. Noor, *Some iterative methods for nonconvex variational inequalities*, Math. Comput. Modelling, **54** (11–12) (2011), 2955–2961.
- [27] X. Qin and N.T. An, *Smoothing algorithms for computing the projection onto a Minkowski sum of convex sets*, Comput. Optim. Appl., **74**(3) (2019), 821–850.
- [28] S. Saejung and P. Yotkaew, *Approximation of zeros of inverse strongly monotone operators in Banach spaces*, Nonlinear Anal., **75**(2) (2012), 742–750.
- [29] Y. Shehu, Q.L. Dong and D. Jiang, *Single projection method for pseudo-monotone variational inequality in Hilbert spaces*, Optimization, **68**(1) (2018), 385–409.
- [30] G. Stampacchia, *Formes bilinéaires coercitives sur les ensembles convexes*, C. R. Acad. Sci. Paris, **258** (1964), 4413–4416.
- [31] W. Takahashi, *Introduction to nonlinear and convex analysis*, Yokohama, 2009.
- [32] B. Tan, Z. Zhou and S. Li, *Viscosity-type inertial extragradient algorithms for solving variational inequality problems and fixed point problems*, J. Appl. Math. Comput., **68**(2) (2022), 1387–1411.
- [33] I. Yamada and N. Ogura, *Hybrid steepest descent method for variational inequality problem over the fixed point set of certain quasi-nonexpansive mappings*, Numer. Funct. Anal. Optim., **25**(7–8) (2004), 619–655.
- [34] P. Yordsorn, P. Kumam and H.U. Rehman, *Modified two-step extragradient method for solving the pseudomonotone equilibrium programming in a real Hilbert space*, Carpathian J. Math., **36**(2) (2020), 313–330.
- [35] P. Yordsorn, P. Kumam, H.U. Rehman and A.H. Ibrahim, *A weak convergence self-adaptive method for solving pseudomonotone equilibrium problems in a real Hilbert space*, Mathematics, **8**(7) (2020), Article ID 1165.