



Source details

Tropical Journal of Natural Product Research

Years currently covered by Scopus: from 2017 to 2024

Publisher: Faculty of Pharmacy, University of Benin

ISSN: 2616-0684 E-ISSN: 2616-0692

Subject area: [Pharmacology, Toxicology and Pharmaceutics: Pharmaceutical Science](#)

[Medicine: Complementary and Alternative Medicine](#)

[Pharmacology, Toxicology and Pharmaceutics: Pharmacology](#)

[Chemistry: Analytical Chemistry](#)

[View all](#)

Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

CiteScore 2023

1.1



SJR 2023

0.177



SNIP 2023

0.373



[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

CiteScore 2023

1.1 = $\frac{1,396 \text{ Citations } 2020 - 2023}{1,257 \text{ Documents } 2020 - 2023}$

Calculated on 05 May, 2024

CiteScoreTracker 2024

1.2 = $\frac{1,582 \text{ Citations to date}}{1,276 \text{ Documents to date}}$

Last updated on 08 July, 2024 • Updated monthly

CiteScore rank 2023

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics	#121/183	34th
Pharmaceutical Science		
Medicine		
Complementary	#79/105	25th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

[What is Scopus](#)
[Content coverage](#)
[Scopus blog](#)
[Scopus API](#)
[Privacy matters](#)

Language

[日本語版を表示する](#)
[查看简体中文版本](#)
[查看繁體中文版本](#)
[Просмотр версии на русском языке](#)

Customer Service

[Help](#)
[Tutorials](#)
[Contact us](#)

ELSEVIER

[Terms and conditions](#) [Privacy policy](#)

All content on this site: Copyright © 2024 Elsevier B.V. [↗](#), its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the Creative Commons licensing terms apply. We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies [↗](#).



Tropical Journal of Natural Product

Welcome. Please, turn your
ADBLOCK off!

It looks like you're using an ad blocker. That's
okay—who doesn't? But without advertising
income, we can't keep making this site awesome.

COUNTRY

Nigeria



Universities and research
institutions in Nigeria



Media Ranking in Nigeria

SUBJECT AREA AND C

Biochemistry, Gen
Biology

Biochemistry
Molecular Mex

Chemistry

Analytical Chemistry

Medicine

Complementary and Alternative
Medicine

Pharmacology, Toxicology and
Pharmaceutics

Drug Discovery

Pharmaceutical Science

Pharmacology

Thank You!

REFRESH

f Benin

H-INDEX

11

PUBLICATION TYPE

Journals

ISSN

26160684, 26160692

COVERAGE

2017-2023

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

[Contact](#)

SCOPE

The Tropical Journal of Natural Product Research is open access, peer-reviewed international journal aimed at making important contributions in the field of Natural Product Research, Pharmaceutical and Natural Sciences. The journal covers all aspects of Pharmaceutical research, chemistry and biochemistry of naturally occurring compounds, ethnomedicine, ethnopharmacology, pharmacognosy, biomedical research, biotechnology and related disciplines. The journal welcome submissions from a broad spectrum of scientific endeavour involving biological evaluation of natural substances of plant, microbial and animal origin against different disease targets, processes or therapeutic strategies that can lead to or assist in the prevention and management of chronic and infectious diseases, clinical therapeutics, Isolation and characterization of metabolites, structure elucidation, synthesis and experimental biosynthesis of natural Product as well as developments of methods in these areas. Research papers in the fields of chemistry-biology boundary, e.g. fermentation chemistry, plant tissue culture investigations etc. are also welcomed. Although the journal focuses mainly on original research articles, timely, concise and focused reviews on recent progress in active areas of natural Product is also encouraged.

Join the conversation about this journal

Quartiles



FIND SIMILAR JOURNALS ?

01

1
Pharmacognosy Journal

IND

76%
similarity

2
Journal of Traditional and
Complementary Medicine
TWN

76%
similarity

3
Pharmacologyonline
ITA


75%
similarity

4
Pharmacognosy Research
IND

73%
similarity

5
Oriental Pharmacy and
Experimental Medicine
USA

72%
similarity



Tropical Journal of Natural Product Research

Q3 Pharmaceutical Science

SJR 2023 0.18

best quartile


powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool**.



Metrics based on Scopus® data as of March 2024

Hesham Essam 5 months ago

please, I want to ask whether the journal is indexed in scopus till now 1-2-2024

reply

Yurnadi 4 months ago

The journal is indexed in scopus. Please check link below:

<https://www.scopus.com/record/display.uri?eid=2-s2.0-85186880420>

Melanie Ortiz 5 months ago

Dear Hesham, thank you very much for your comment. We suggest you consult the Scopus database directly. Keep in mind that the SJR is a static image (the update is made one time per year) of a database (Scopus) which is changing every day. The Scopus' update list can also be consulted here:

SCImago Team



Antioxidant and Anticancer Potentials and Metabolic Profiling of Benjakul, A Thai Herbal Preparation

Kunyarat Duenngai¹, Bundit Promraksa², Sutthiwan Janthamala³, Malinee Thanee⁴, Papitchaya Sirithawat³, Supaporn Wisungre¹, Sumalin Deechan¹, Nawarat Meechai¹, Phakamas Paratang¹, Anchalee Techasen^{3*}

¹Department of Thai Traditional Medicine, Faculty of Science and Technology, Phetchabun Rajabhat University, Phetchabun, 67000, Thailand

²Regional Medical Sciences Center 2 Phitsanulok, Department of Medical Sciences, Ministry of Public Health, Phitsanulok, 65000, Thailand

³Center for Medical Diagnostic Laboratories (CMDL), Faculty of Associated Medical Sciences, Khon Kaen University, Khon Kaen, 40002, Thailand

⁴Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen, 40002, Thailand

ARTICLE INFO

Article history:

Received 18 January 2024

Revised 10 April 2024

Accepted 11 April 2024

Published online 01 May 2024

ABSTRACT

Benjakul (BJK), a Thai folk remedy has been used in traditional medicine for the treatment of various illnesses. This study aimed to assess the antioxidant, and anticancer properties, and metabolic profile of Benjakul. The antioxidant activity of the ethanol extract of BJK and its component medicinal plants (*Piper retrofractum* Vahl., *Piper sarmentosum* Roxb., *Piper interruptum* Opiz., *Plumbago indica* L., and *Zingiber officinale* Rosc.) was assessed using the Ferric Reducing Antioxidant Power (FRAP) assay, and 2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical scavenging assay. The anticancer activity was assessed through cytotoxic effect against cholangiocarcinoma (CCA) (bile duct cancer) cell lines (KKU-213B and KKU-100) using the 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. The metabolic fingerprint and potential bioactive compounds in the ethanol extract of BJK and component medicinal plants was investigated using Nuclear Magnetic Resonance (NMR) spectroscopy. The FRAP and DPPH assays revealed that the ethanol extracts of BJK and its component medicinal plants possessed antioxidant activity with *Zingiber officinale* exhibiting the highest activity. The FRAP value and EC₅₀ for DPPH radical scavenging activity for *Zingiber officinale* were 94.860 ± 0.156 µg AAE/mg extract and 86.63 g/mL, respectively. The metabolic profiling identified vanillic acid and curcumin as the main bioactive metabolites in BJK and its component medicinal plants. The partial least squares (PLS) loading plot revealed that the methoxy group is associated with the inhibition of CCA cells, and phenolic hydroxyl protons as the potential pharmacophore of the bioactive molecules. Therefore, BJK could be considered as a potentially effective treatment for bile duct cancer.

Copyright: © 2024 Duenngai *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Keywords: Benjakul, Thai Traditional medicine, Antioxidant, Anticancer, Metabolic profile.

Introduction

Natural products have played an important role in the discovery of new drugs and their innovative mechanisms of action. Antioxidant effect is one of the most extensively studied properties exhibited by substances derived from natural products.¹ Many plants have been reported as sources of natural antioxidants, and the discovery and use of these naturally occurring antioxidants derived from plant products is beneficial to humankind.

Thai traditional medicine based on natural products especially medicinal plants plays an important role in the treatment of various diseases. The bioactive components and pharmacological effects of these plants are still insufficiently studied.¹

Phytochemicals with antioxidant properties derived from medicinal plants have been shown to extend the lifespan of animal models.² For example, proanthocyanidins and polyphenols extracted from blueberries, protect cells against oxidative stress.³

*Corresponding author. E mail: anchte@kku.ac.th
Tel: +66-43-202086

Citation: Duenngai K, Promraksa B, Janthamala S, Thanee M, Sirithawat P, Wisungre S, Deechan S, Meechai N, Paratang P, Techasen A. Antioxidant and Anticancer Potentials and Metabolic Profiling of Benjakul, A Thai Herbal Preparation. Trop J Nat Prod Res. 2024; 8(4):6877-6883

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria

A traditional Chinese multi-herbal extract was discovered to produce antioxidant and anti-aging effects.²

Benjakul (BJK), a Thai herbal remedy, is a common treatment for inflammation and cancer in Thai traditional medicine and it is included on the National Lists of Essential Medicines. BJK comprises five medicinal plants including *Piper retrofractum* Vahl. (Pr), *Piper sarmentosum* Roxb. (Ps), *Piper interruptum* Opiz. (Pi), *Plumbago indica* L. (Pn), and *Zingiber officinale* Rosc. (Zo). BJK has been reported to be used as anticancer remedies in Thai traditional medicine. It has been used as the initial medication to reduce pain and manage patients' symptoms prior to the administration of other treatments.^{4,5} Previous reports have indicated that the herbal remedy and its individual plant components exhibit cytotoxic effects against cancer cells⁶ anti-allergic⁷ and anti-inflammatory effects.⁷⁻¹⁰ In addition, isolated compounds from BJK have shown cytotoxic activity on breast cancer cells,¹¹ and large and small cell lung cancer cells.^{12,13}

It has been demonstrated that plant extracts and plant-derived compounds exhibit anticancer properties by activating the apoptotic pathway.¹⁴⁻¹⁷ Some specific plant components in the BJK formulation caused cell death by apoptotic signalling.¹⁸⁻²¹

At present, nuclear magnetic resonance (NMR) spectroscopy is increasingly being used in the fields of food science,²²⁻²⁴ medicine and health,²⁵⁻²⁹ and natural products.³⁰ Recently, NMR was used to analyze the Chinese herbal medicine, *Sophora flavescens*, and two novel chemical compounds were identified.³¹ In addition, a previous study used NMR-based metabolomics to identify changes in the metabolic profile of a local cultivar of *Andrographis paniculata* during its growth period.³²

The purpose of this study was to evaluate the antioxidant and anticancer activities of BJK and identify its bioactive constituents, which could serve as new therapeutic alternatives for the prevention and treatment of cancer.

Materials and Methods

Reagents and chemicals

Unless otherwise stated, the chemicals used in this study were purchased from Sigma-Aldrich (St. Louis, MO, USA). The F-12 Nutrient Mixture (HAM) and fetal bovine serum were purchased from Thermo Fisher Scientific (MA, United States).

Plant collection and identification

The BJK remedy is composed of 5 medicinal plant species, *Piper retrofractum* Vahl. (Pr), *Piper sarmentosum* Roxb. (Ps), *Piper interruptum* Opiz. (Pi), *Plumbago indica* L. (Pn), and *Zingiber officinale* Rosc. (Zo). Each medicinal plant was collected in July, 2022 from the northern region of Thailand. The Plant materials were authenticated by Ms. Dujhathai Anekchai, Pharm.D, at the Pharmaceutical Laboratory Service Centre, Faculty of Pharmaceutical Sciences, Prince of Songkla University, Songkhla, Thailand. The following herbarium no. SKP 146161803, SKP 146161907, SKP 146160902, SKP 148160901 and SKP 206261507 for *Piper retrofractum* Vahl., *Piper sarmentosum* Roxb., *Piper interruptum* Opiz., *Plumbago indica* L., and *Zingiber officinale* Rosc., respectively were issued.

Preparation of extracts

Each herbal plant was dried in an oven at 60°C, then ground into fine powder. The powdered plant (100 g) was extracted by maceration in 1 L of 95% ethanol at room temperature for 5 days. The ethanol extract was filtered using Whatman filter paper no. 1, and the extract was concentrated using a rotary evaporator (Buchi, Essen, Germany) at 50°C under reduced pressure. The concentrated ethanol extracts were stored at 4°C until used.³³

Evaluation of antioxidant activity

Ferric Reducing Antioxidant Power (FRAP) assay

The antioxidant capacity of the ethanol extracts was determined by FRAP assay following a slightly modified procedure.³⁴ The FRAP working solution was freshly prepared by mixing 20 mM ferric chloride (FeCl₃) in acetate buffer, 10 mM 2,4,6-Tri(2-pyridyl)-s-triazine (TPTZ) in acetate buffer and 0.25 mM sodium acetate buffer (pH 3.6) in a 1:1:10 ratio. Each extract was combined with 182 µL of FRAP solution, and the absorbance was measured at 593 nm using a microplate reader. Calibration curve was generated using ascorbic acid at concentrations ranging from 0 to 60 µg/mL. The antioxidant capacity of the extracts was estimated from the equation of the standard curve of ascorbic acid, and the results were expressed as micrograms of ascorbic acid equivalent per milligram of extract (µg AAE/mg extract).

2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical scavenging assay

The working reagent was prepared by dissolving DPPH in absolute ethanol at a concentration of 0.1 mM. Twenty microliters of the extract solution was mixed with 180 µL of DPPH solution. The mixture was incubated at room temperature in the dark for 30 min. The absorbance was measured using a spectrophotometer at a wavelength of 517 nm. The DPPH free radical scavenging capacity was determined and compared with standard ascorbic acid. The percentage of radical scavenging effect was calculated by the following equation:

$$\text{Percentage DPPH radical scavenging effect (\%)} = (\text{Ab} - \text{As})/\text{Ab} \times 100$$

Where; Ab is the absorbance of blank (DPPH without extract), and As is the absorbance of the reaction mixture (DPPH with the extract).

The EC₅₀ values (the concentration needed to achieve a 50% antioxidant effect) of the extracts was calculated from the plot of percentage radical scavenging activity versus concentration of the extracts.

Evaluation of anticancer activity

Cholangiocarcinoma cell lines

Cholangiocarcinoma (CCA) cell lines; KKKU-213B and KKKU-100 were purchased from the Japanese Collection of Research Bioresources (JCRB). The CCA cell lines were produced by Prof. Banchob Sripa of Khon Kaen University in Thailand, and both cell lines were deposited at the JCRB, Osaka, Japan, for thorough characterization. The cell lines were grown in Ham's F-12 medium supplemented with 10% fetal bovine serum and penicillin/streptomycin (50 U/mL and 50 µg/mL), and then placed in a humidified incubator at 37°C with 5% CO₂. A primary normal fibroblast cell line was purchased from American Type Culture Collection (ATCC). The cell line was cultured in Eagles's Minimum Essential Medium (EMEM) containing 10% fetal bovine serum and penicillin/streptomycin (50 U/mL and 50 µg/mL) and incubated in a humidified incubator of 5% CO₂ at 37°C.

Test for cytotoxic activity

The MTT colorimetric test was utilized to assess the cytotoxic effects of BJK. KKKU-213B and KKKU-100 cells were placed in a 96-well plate at a density of 2,000 cells per well. Cells were cultured for 24 h and then treated with various concentrations of the crude ethanol extracts for 48 h and 72 h at 37°C in a 5% CO₂ condition. The cells were then rinsed with phosphate buffered saline (PBS) and exposed to MTT reagent at 37°C for 2 h. The MTT reagent was removed, and DMSO was introduced to dissolve the formazan crystals, and the absorbance was measured at 570 nm using micro-plate reader.

The percentage cell viability was calculated using the following formula:

$$\text{Percentage cell viability (\%)} = \frac{\text{Absorbance of untreated cells} - \text{Absorbance of culture medium}}{\text{Absorbance of treated cells} - \text{Absorbance of culture medium}} \times 100$$

Nuclear magnetic resonance (NMR) acquisition for metabolic profiling

The ethanol extract (100 g) was dissolved in a 1 mL of CD₃OD (with 0.1% v/v tetramethylsilane, TMS). The mixture (600 µL) was placed in a 5-mm NMR tube for NMR analysis. The ¹H-NMR spectrum was obtained using a 400 MHz NMR instrument (Avance, Bruker BioSpin GmbH, Rheinstetten, Germany). The zg30 pulse sequence was applied with water signal suppression and 32 scans at 25°C. Following the capture, typical FID processing methods were performed using TopSpin (Bruker, Biospin, Italy), including Fourier transform, phase correction, and baseline correction. All the ¹H-NMR spectra were calibrated with reference to the internal standard TMS (δ = 0.00 ppm). Metabolite identification was performed by statistical total correlation spectroscopy (STOCSY) by comparing against metabolome databases such the human metabolome database (HMDB), biological magnetic resonance data bank (BMRB), and natural product magnetic resonance database (NP-MRD).^{35,36}

Statistical analysis

Statistical analysis was conducted using IBM SPSS V.23.0 software by SPSS Inc., Chicago, IL. Data were expressed as means ± standard deviation (SD) of three independent experiments. GraphPad Prism version 8.0 for Windows (GraphPad Software, San Diego, CA) was used to compute and construct a bar chart for cytotoxic activity and a dose-response curve for EC₅₀. One-way ANOVA was used to determine the statistical significant difference among means. P < 0.05 was considered statistically significant.

For NMR data, multivariate data analysis (MVDA) was applied to identify the bioactive metabolites. All spectra from 2.7 were aligned based on a hierarchical cluster-based peak alignment (CluPA) in R statistical software using the ASICS package,^{37,38} and normalized by probabilistic quotient normalization method.³⁹ Then, pareto scaling was applied using the Metabom8 package to all data prior to MVDA including unsupervised principal component analysis (PCA) and supervised partial least squares regression analysis (PLS). The fraction of the variance explained (R²) by a component, indicates how well the variation of a variable is explained. The goodness of fit prediction (Q²) was used to determine how well a variable could be predicted and the accuracy of prediction was estimated by cross validation. Quantitative

structure-activity relationships (QSAR) aim to maximize unexplained variation in dependent measures. Q^2 values equal to or greater than 0.4 were considered acceptable (Worley and Powers, 2015).

Results and Discussion

Antioxidant activity

Phyto-antioxidants, which are found in medicinal plants, can actually prevent oxidative stress. Both FRAP assay and DPPH radical scavenging assay were used to assess the antioxidant properties of BJK herbal remedy and its component medicinal plants extracts.

FRAP assay was used to determine the ability of antioxidants present in the herbal plants to reduce ferric ion (Fe^{3+}) to ferrous ion (Fe^{2+}) through the donation of an electron. The correlation coefficient (r) of the calibration curve, plotted using standard ascorbic acid concentrations versus absorbance was greater than 0.95. The results revealed that the ethanol extracts of BJK herbal remedy and its components medicinal plants had antioxidant activity ranging from 26.130 ± 0.962 to 94.860 ± 0.156 μ g AAE/mg extract, with the ethanol extract of *Z. officinale* exhibiting the highest activity with a FRAP value of 94.860 ± 0.156 μ g AAE/mg extract (Figure 1). However, in a previous study, the FRAP value of ginger juice, was found to be 348.70 μ gAAE/mg extract,⁴¹ a value significantly higher than what was obtained in our study. The variation in the FRAP value may be attributed to the differences in the extraction method used.

The DPPH scavenging activity of the extracts was expressed in terms of half maximal effective concentration (EC_{50}). The EC_{50} values of the extracts range from 86.63 to 237.40 μ g/mL compared to the standard ascorbic acid (15.10 μ g/mL). Lower EC_{50} value indicates higher DPPH radical scavenging activity, hence, a higher antioxidant potential. Overall, the ethanol extract of Zo had the lowest EC_{50} value (86.63 μ g/mL) followed by BJK, Pn, Ps, Pi and Pr (Figure 2A). At a concentration of 250 μ g/mL, Zo had the highest DPPH radical scavenging activity (Figure 2B). This indicates that the ethanol extract of Zo had the highest antioxidant activity compared to the other extracts.

Phyto-antioxidants, which are found in medicinal plants, can actually prevent oxidative stress, which has been implicated in cancer pathogenesis.⁴⁰ In the present study, DPPH radical scavenging and FRAP assays were used to assess the antioxidant properties of BJK remedy and its component medicinal plants extracts. *Z. officinale* (Zo) commonly called ginger exhibited the highest antioxidant activity in both FRAP and DPPH radical scavenging assays. This observation is similar to the findings from previous studies where the extract of *Z. officinale* was found to possess the highest antioxidant activity among other plants investigated.^{42,43}

Anticancer activity

CCA patients usually undergo chemotherapy because this cancer is often diagnosed at an advanced stage.⁴⁴ Nevertheless, CCA treatment proves difficult due to strong chemoresistance mechanisms,⁴⁵ which suggest a limited response to available chemotherapeutic agents. Therefore, this study has identified the herbal remedy Benjakul (BJK) as a new and potentially useful alternative treatment for CCA. The potential of BJK to inhibit the proliferation of KKKU-213B and KKKU-100 CCA cell lines was determined by treating the cells with the ethanol extracts of BJK and its herbal components for 48 and 72 h. At a concentration of 100 μ g/mL, each herbal extract exhibited strong anti-proliferative effect against CCA cell lines, with the most potent anti-proliferative effect induced by Zo, followed by Pn, Pr, BJK, Ps and Pi, in that order (Figure 3A-3D).

The present results demonstrated that the IC_{50} of BJK at 48 and 72 h was 93.61 μ g/mL and 76.13 μ g/mL for KKKU-213B, and 89.94 μ g/mL and 80.16 μ g/mL for KKKU-100 (Figure 4A and 4B). In addition, BJK was found to be non-toxic to normal fibroblast cells at the same concentration as shown in Figure 4C.

BJK exhibited dose-dependent cytotoxic effects on CCA cells while showing lower toxicity towards normal fibroblast cells. BJK has previously been utilized to treat many kinds of cancer to improve the quality of life for patients.¹³ According to previously published reports, the difference in the IC_{50} values of BJK for CCA compared to MCF-7

breast cancer and COR-L23 lung cancer cell lines is greater than twofold and threefold, respectively.^{11,13}

Metabolic profiling of BJK remedy and its component medicinal plants

All secondary metabolites in ethanol extracts of BJK and its component medicinal plants were examined concurrently using metabolic profiling. A Principal Component Analysis (PCA) transformed the high-dimensional NMR spectral data to a few latent variables, with the first principal component (PC1) accounting for the greatest variation and the second principal component (PC2) accounting for around 85% of the overall variance.




Figure 1: Antioxidant capacity of ethanol extracts of BJK and its component medicinal plants as determined by FRAP assay. Values are presented as mean $\pm$ SD of triplicate determination. BJK = Benjakul, Pr = *Piper retrofractum*, Ps = *Piper sarmentosum*, Pi = *Piper interruptum*, Pn = *Plumbago indica*, Zo = *Zingiber officinale*.




Figure 2: DPPH radical scavenging activity of BJK remedy and its medicinal plants extracts. (A): Half maximal effective concentration (EC_{50}) of the extracts compared to standard ascorbic acid. (B): Percentage scavenging activity of the extracts at 250 μ g/mL.

BJK = Benjakul, Pr = *Piper retrofractum*, Ps = *Piper sarmentosum*, Pi = *Piper interruptum*, Pn = *Plumbago indica*, Zo = *Zingiber officinale*.




Figure 3: Cytotoxic effects of BJK and its medicinal plant components on CCA cell lines. KKKU-213B (A & B) and KKKU-100 (C & D) were treated with ethanol extracts of BJK and its herbal components for 48 and 72 h.

The NMR spectra among the herbal extracts were distinguishable due to the differences in their phytochemical constituents (Figure 5). On the negative side of the plot, the score plot demonstrated that Pn was distinct from the other herbal extracts. The CCA cell inhibitory activity of Zo, Pr, and Pn was related to their metabolites. The PLS loading plot showed that the metabolites are positively correlated to the CCA cell inhibitory activity on both KKKU-213B and KKKU-100. The metabolites assignment of the peak at 3.51 ppm corresponded to methoxy (-OMe) group of vanillin and vanillic acid.⁴⁶ Hypoxia-inducible factor 1 (HIF-1) expression suppression has also been demonstrated as a mechanism by which vanillic acid inhibit the metastasis of colon cancer.⁴⁷ Vanillic acid has been shown to inhibit lung carcinogenesis by anti-inflammatory and free radical scavenging mechanisms in Swiss albino mice.⁴⁸ Vanillin extracted from millet was shown to inhibits the proliferation of HT-29 cells through G0/G1 cell cycle arrest and induction of apoptosis.⁴⁹

The bioactive metabolites in the crude herbal extracts, guided by CCA cell inhibition, were investigated using a partial least squares (PLS) regression analysis. The PLS score plot as shown in Figure 6 revealed a cluster of samples colored by CCA viability after being treated with each crude herbal extract. The cancer cell inhibitory activity was related to the metabolites of Zo, Pr, and Pn. The PLS loading plot showed that the methoxy (-OMe) group was correlated to the cancer cell inhibition of both KKKU-213B and KKKU-100. Methoxy and hydroxyl groups contribute to the antioxidant activity of phenolic acid through their ability to transfer proton and quench free radicals. Figure 7 shows the partial least squares discriminant analysis (PLS-DA) loading plot of the bioactive compounds. The doublet of doublets (dd) and doublets (d) peaks at 6.44 and 6.65 ppm, respectively, corresponds to the protons of the phenolic hydroxyl group of vanillic acid and curcumin, respectively. Silver nanoparticles of vanillic acid have been developed to improve drug delivery and enhance biological effect of vanillic acid,⁵⁰ but the metabolic alterations and health implications continue to pose a challenge. Hence, the biological mechanisms responsible for the anti-CCA effect of vanillic acid need to be studied in detail. This study is the first to show the effect of BJK on CCA cell lines. We believe that BJK herbal remedy could be potential therapy against bile duct cancer.

Conclusion

This study has evaluated the antioxidant, and anticancer activities, and examined the metabolic profile of BJK and its component medicinal plants. The ethanol extracts of BJK and its component medicinal plants possessed antioxidant and cytotoxic activities against cholangiocarcinoma (bile duct cancer) cell lines (KKU-213B and KKKU-100), and cytoprotective properties on normal fibroblast cells. Vanillic acid and curcumin were identified as the main bioactive compounds in

the plants. The protons of the phenolic hydroxyl group was identifies as the main pharmacophore of the potentially bioactive compounds (vanillic acid and curcumin). Therefore, BJK could be considered as a potentially effective treatment for bile duct cancer.

Conflict of Interest

The authors declare no conflict of interest.

Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.




Figure 4: Comparison of the cytotoxic potential of BJK against KKKU-213B (A), KKKU-100 (B) CCA cell lines and normal fibroblast cells (C) at the same concentration.




Figure 5: Unsupervised principal component analysis score plot. All phytochemical metabolites in BJK and its component medicinal plant extracts were examined using metabolic profiling. The NMR spectra were distinguishable among the different herbal extracts.




Figure 6: Partial least squares (PLS) regression score plot of crude herbal extract colored by the cell viability; (A): KCU-213B and (B): KCU-100. The PLS score plot revealed a cluster of samples colored by CCA viability after being treated with each crude herbal extract.




Figure 7: Partial least squares discriminant analysis (PLS-DA) loading plot of bioactive compounds; (A): KCU-213B and (B): KCU-100. The methoxyl and hydroxyl groups in phenolic acids can quench and transfer their proton to free radicals. The doublet of doublets (dd) and doublets (d) at 6.44 and 6.65 ppm, respectively, were the protons in the phenolic hydroxyl groups of vanillic acid and curcumin, respectively.

Acknowledgements

This work was supported by the Science Promotion Fund for research and innovation Fiscal Year 2021 innovative research and development plan (FRB640052/11).

We wish to thank the Institute of Research and Development, Phetchabun Rajabhat University. We would like to acknowledge Prof. Ross H Andrews of Imperial College London for editing the Manuscript via Publication Clinic KKU, Thailand.

References

- Yanda L, Tatsimo SJN, Tamokou JDD, Matsuet-Takongmo G, Meffo-Dongmo SC, Meli Lannang A, Sewald N. Antibacterial and antioxidant activities of isolated compounds from *Prosopis africana* leaves. *Int J Anal Chem*. 2022; 2022:4205823.
- Wetchakul P, Goon JA, Adekoya AE, Olatunji OJ, Ruangchuay S, Jaisamut P, Issuriya A, Kunworarath N, Limsuwan S, Chusri S. Traditional tonifying polyherbal infusion, Jatu-Phala-Tiga, exerts antioxidant activities and extends lifespan of *Caenorhabditis elegans*. *BMC Complement Alternat Med*. 2019; 19:1–11.
- Louis XL, Thandapilly SJ, Kalt W, Vinqvist-Tymchuk M, Aloud BM, Raj P, Yu L, Le H, Netticadan T. Blueberry polyphenols prevent cardiomyocyte death by preventing calpain activation and oxidative stress. *Food Funct*. 2014; 5(8):1785–1794.
- Itharat A, Singchangchai P, Ratanasuwan P. Wisdom of Southern Thai traditional doctors. Prince of Songkla University, Songkla. 1998. 126 p.
- Rachawat P, Pinsornsak P, Kanokkangsadal P, Itharat A. Clinical efficacy and safety of Benjakul remedy extract for treating primary osteoarthritis of knee compared with diclofenac: double blind, randomized controlled trial. *Evid-Based Complement Alternat Med*. 2017; 2017:9593580.
- Pawa KK. *In vitro* cytotoxic activity of Benjakul herbal preparation and its active compounds against human lung, cervical and liver cancer cells. *J Med Assoc Thai*. 2012; 95(1):S127–134.
- Makchuchit S, Rattarom R, Itharat A. The anti-allergic and anti-inflammatory effects of Benjakul extract (a Thai traditional medicine), its constituent plants and its some pure constituents using *in vitro* experiments. *Biomed Pharmacother*. 2017; 89:1018–1026.
- Burodom A and Itharat A. Inflammatory suppressive effect of Benjakul, a Thai traditional medicine on intestinal epithelial cell line. *J Med Plant Res*. 2013; 7(44):3286–3291.
- Thummawan C, Itharat A, Asasutjarit R. Anti-inflammatory activities of nanoemulsion containing Benjakul Remedy extract. *Thammasat Med J*. 2019; 19(2):359–370.
- Kuropakornpong P, Itharat A, Panthong S, Sireeratawong S, Ooraikul B. *In vitro* and *in vivo* anti-inflammatory activities of Benjakul: a potential medicinal product from Thai traditional medicine. *Evid-Based Complement Alternat Med*. 2020; 2020:9760948.
- Sakpakdeejaroen I and Itharat A. Cytotoxic compounds against breast adenocarcinoma cells (MCF-7) from Pikutbenjakul. *J Health Res*. 2009; 23(2):71–76.
- Rattarom R, Hansakul P, Sakpakdeejaroen I, Itharat A. Induction of apoptosis in human lung cancer NCI-H226 cells by ethanolic extract of Benjakul preparation and its isolated compound. *Planta Med*. 2013; 79(13):PN89.
- Itharat A, Rattarom R, Hansakul P, Sakpakdeejaroen I, Ooraikul B, Davies NM. The effects of Benjakul extract and its isolated compounds on cell cycle arrest and apoptosis in human non-small cell lung cancer cell line NCI-H226. *Res Pharm Sci*. 2021; 16(2):129.
- Pfeffer CM and Singh ATK. Apoptosis: a target for anticancer therapy. *Int J Mol Sci*. 2018; 19(2):448.
- Golshan A, Amini E, Emami SA, Asili J, Jalali Z, Sabouri-Rad S, Sanjar-Mousavi N, Tayarani-Najarian Z. Cytotoxic evaluation of different fractions of *Salvia chorassanica* Bunge on MCF-7 and DU 145 cell lines. *Res Pharm Sci*. 2016; 11(1):73.
- Homayoun M, Targhi RG, Soleimani M. Anti-proliferative and anti-apoptotic effects of grape seed extract on chemo-resistant OVCAR-3 ovarian cancer cells. *Res Pharm Sci*. 2020; 15(4):390.
- Ahmadi F, Mojarrab M, Ghazi-Khansari M, Hosseinzadeh L. A semipolar fraction of petroleum ether extract of *Artemisia aucheri* induces apoptosis and enhances the apoptotic response to doxorubicin in human neuroblastoma SK-NMC cell line. *Res Pharm Sci*. 2015; 10(4):335.
- Ahmad A, Banerjee S, Wang Z, Kong D, Sarkar FH. Plumbagin-induced apoptosis of human breast cancer cells is mediated by inactivation of NF- κ B and Bcl-2. *J Cell Biochem*. 2008; 105(6):1461–1471.
- Warin RF, Chen H, Soroka DN, Zhu Y, Sang S. Induction of lung cancer cell apoptosis through a p53 pathway by [6]-shogaol and its cysteine-conjugated metabolite M2. *J Agric Food Chem*. 2014; 62(6):1352–1362.
- Gomathinayagam R, Sowmyalakshmi S, Mardhatillah F, Kumar R, Akbarsha MA, Damodaran C. Anticancer mechanism of plumbagin, a natural compound, on non-small cell lung cancer cells. *Anticancer Res*. 2008; 28(2A):785–792.
- Chen CY, Liu TZ, Liu YW, Tseng WC, Liu RH, Lu FJ, Lin YS, Kuo SH, Chen CH. 6-shogaol jalkanone from ginger induces apoptotic cell death of human hepatoma p53 mutant *Mahlavu subline* via an oxidative stress-mediated caspase-dependent mechanism. *J Agric Food Chem*. 2007; 55(3):948–954.
- Rabbi F, Sharmin N, Khatun MCS, Hasan M, Muhit MA. Comparative Phytochemical and Biological Analyses of commercial Green Tea Products Marketed in Dhaka, Bangladesh. *Trop J Nat Prod Res*. 2023; 7(12):5470–5478.
- Gougeon L, Da Costa G, Le Mao I, Ma W, Teissedre PL, Guyon F, Richard T. Wine analysis and authenticity using 1 H-NMR metabolomics data: Application to Chinese wines. *Food Anal Methods*. 2018; 11:3425–3434.
- Ravaglia LM, Freitas D dos S, Ricci TG, Nazario CED, Alcantara GB. Sodium quantitation in soft drinks: A rapid methodology by qNMR. *Magn Reson Chem*. 2020; 58(2):186–190.
- Herlyanti K, Fakhrudin N, Murwanti R, Wahyono A. Identification of Cubebin from Cubeb (*Piper cubeba*) as An Antiplatelet Agent. *Trop J Nat Prod Res*. 2023; 7(7):3392–3397.
- Jaurila H, Koivukangas V, Koskela M, Gäddnäs F, Myllymaa S, Kullaa A, Salo T, Ala-Kokko TI. 1H NMR based metabolomics in human sepsis and healthy serum. *Metabol*. 2020; 10(2):70.
- Petrás M, Kalenska D, Samos M, Bolek T, Sarlinova M, Racay P, Halasova E, Strbak O, Stasko J, Musak L. NMR plasma metabolomics study of patients overcoming acute myocardial infarction: in the first 12 h after onset of chest pain with statistical discrimination towards metabolomic biomarkers. *Physiol Res*. 2020; 69(5):823.
- Igbo EU, Igoli OJ, Onyiriuka OS, Ogukwe EC, Ayuk AA, Gray IA. Isolation and Characterization of Pyropheophorbide-a from *Moringa oleifera* Lam. *Trop J Nat Prod Res*. 2019; 3(10):314 - 318.
- Tamvapee P, Samosorn S, Watanapokasin R. Identification of Active Fatty Acid Mixture Extracted from Rice Bran Oil Using NMR Spectroscopy. *Trop J Nat Prod Res*. 2023; 7(4):2756–2759.
- Wang ZF, You YL, Li FF, Kong WR, Wang SQ. Research progress of NMR in natural product quantification. *Molecules*. 2021; 26(20):6308.

31. Lin Y, Li JJ, He L, Li QR, Long QD, Zhang X, Zeng Z. A new modified pterocarpan glycoside from *Sophora flavescens*. Nat Prod Res. 2023; 37(20):3374-3379.
32. Tajidin NE, Shaari K, Maulidiani M, Salleh NS, Ketaren BR, Mohamad M. Metabolite profiling of *Andrographis paniculata* (Burm. f.) (Nees. young and mature leaves at different harvest ages using ¹H NMR-based metabolomics approach. Sci Rep. 2019; 9(1):16766.
33. Promraksa B, Phetcharaburanin J, Namwat N, Techasen A, Boonsiri P, Loilome W. Evaluation of anticancer potential of Thai medicinal herb extracts against cholangiocarcinoma cell lines. PLoS One. 2019; 14(5):e0216721.
34. Benzie IFF and Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. Anal Biochem. 1996; 239(1):70-76.
35. Cloarec O, Dumas ME, Craig A, Barton RH, Trygg J, Hudson J, Blancher C, Gauguier D, Lindon JC, Holmes E. Statistical total correlation spectroscopy: an exploratory approach for latent biomarker identification from metabolic ¹H NMR data sets. Anal Chem. 2005; 77(5):1282-1289.
36. Wishart DS, Guo A, Oler E, Wang F, Anjum A, Peters H, Dizon R, Sayeeda Z, Tian S, Lee BL. HMDB 5.0: the human metabolome database for 2022. Nucl Acids Res. 2022; 50(D1):D622-631.
37. Lefort G, Liaubet L, Canlet C, Tardivel P, Pèrè MC, Quesnel H, Paris A, Iannuccelli N, Vialaneix N, Servien R. ASICS: an R package for a whole analysis workflow of 1D ¹H NMR spectra. Bioinformatics. 2019; 35(21):4356-4363.
38. Vu TN, Valkenburg D, Smets K, Verwaest KA, Dommissie R, Lemiere F, Verschoren A, Goethals B, Laukens K. An integrated workflow for robust alignment and simplified quantitative analysis of NMR spectrometry data. BMC Bioinformatics. 2011; 12(1):1-14.
39. Dieterle F, Ross A, Schlotterbeck G, Senn H. Probabilistic quotient normalization as robust method to account for dilution of complex biological mixtures. Application in ¹H NMR metabolomics. Anal Chem. 2006; 78(13):4281-4290.
40. Ma L, Zhang M, Zhao R, Wang D, Ma Y, Ai L. Plant natural products: promising resources for cancer chemoprevention. Molecules. 2021; 26(4):933.
41. Bekkouch O, Dalli M, Harnafi M, Touiss I, Mokhtari I, Assri S El, Harnafi H, Choukri M, Ko SJ, Kim B. Ginger (*Zingiber officinale* Roscoe), lemon (*Citrus limon* L.) juices as preventive agents from chronic liver damage induced by CCl₄: A Biochemical and Histological Study. Antioxidants. 2022; 11(2):390.
42. Khalaf NA, Shakya AK, Al-Othman A, El-Agbar Z, Farah H. Antioxidant activity of some common plants. Turk J Biol. 2008; 32(1):51-55.
43. Ghasemzadeh A, Jaafar HZE, Rahmat A. Antioxidant activities, total phenolics and flavonoids content in two varieties of Malaysia young ginger (*Zingiber officinale* Roscoe). Molecules. 2010; 15(6):4324-4333.
44. Marin JJG, Lozano E, Herraez E, Asensio M, Di Giacomo S, Romero MR, Briz O, Serrano MA, Efferth T, Macias RIR. Chemoresistance and chemosensitization in cholangiocarcinoma. Biochim Biophys Acta Mol Basis Dis. 2018; 1864(4 Pt B):1444-1453.
45. Wu HJ and Chu PY. Role of cancer stem cells in cholangiocarcinoma and therapeutic implications. Int J Mol Sci. 2019; 20(17):4154.
46. Bogdan M, Floare CG, Pirnau A. ¹H NMR investigation of self-association of vanillin in aqueous solution. In: J Physics: Conf Ser. IOP Publishing; 2009. 012002 p.
47. Gong J, Zhou S, Yang S. Vanillic acid suppresses HIF-1 α expression via inhibition of mTOR/p70S6K/4E-BP1 and Raf/MEK/ERK pathways in human colon cancer HCT116 cells. Int J Mol Sci. 2019; 20(3):465.
48. Velli SK and Thiruvengadam D. Vanillic acid inhibits lung carcinogenesis by modulates glycoprotein abnormalities, membrane-bound enzymes, and inflammatory markers. Int J Pharm Pharm Sci. 2020; 12(3):83-88.
49. Ramadoss DP and Sivalingam N. Vanillin extracted from Proso and Barnyard millets induce apoptotic cell death in HT-29 human colon cancer cell line. Nutr Cancer. 2020; 72(8):1422-1437.
50. Alamri ES, El Rabey HA, Alzahrani OR, Almutairi FM, Attia ES, Bayomy HM, Albalwi RA, Rezk SM. Enhancement of the Protective Activity of Vanillic Acid against Tetrachloro-Carbon (CCl₄) Hepatotoxicity in Male Rats by the Synthesis of Silver Nanoparticles (AgNPs). Molecules. 2022; 27(23):8308.