

Already have a manuscript?

Use our Manuscript Matcher to find the best relevant journals!

[Find a Match](#)

Filters

[Clear All](#)

Web of Science Coverage

Core Collection

- ☐ Science Citation Index Expanded (SCIE)
- ☐ Social Sciences Citation Index (SSCI)
- ☐ Arts & Humanities Citation Index (AHCI)
- ☐ Emerging Sources Citation Index (ESCI)

Current Contents

- ☐ Agriculture, Biology & Environmental Sciences
- ☐ Arts & Humanities
- ☐ Business Collection
- ☐ Clinical Medicine
- ☐ Electronics & Telecommunications Collection

Refine Your Search Results

Caryologia

[Search](#)

Sort By: Title (A-Z)

Search Results

Found 1 results (Page 1)

[Share These Results](#)

Exact Match Found

CARYOLOGIA

[OPEN ACCESS](#)

Publisher: FIRENZE UNIV PRESS , JOURNALS DIVISION, BORGO ALBIZI, 28, FIRENZE, ITALY, 50122

ISSN / eISSN: 0008-7114 / 2165-5391

Web of Science Core Collection: **Not indexed in Web of Science Core Collection**

Additional Web of Science Indexes: Biological Abstracts | BIOSIS Previews | Zoological Record

[Share This Journal](#)

[View profile page](#)

* Requires free login.

1 - 1 of 1

Items per page:

10

Editorial Disclaimer: As an independent organization, Clarivate does not become involved in and is not responsible for the editorial management of any journal or the business practices of any publisher. Publishers are accountable for their journal performance and compliance with ethical publishing standards. The views and opinions expressed in any journal are those of the author(s) and do not necessarily reflect the views or opinions of Clarivate. Clarivate remains neutral in relation to territorial disputes, and allows journals, publishers, institutes and authors to specify their address and affiliation details including territory.

Criteria for selection of newly submitted titles and re-evaluation of existing titles in the Web of Science are determined by the Web of Science Editors in their sole discretion. If a publisher's editorial policy or business practices negatively impact the quality of a journal, or its role in the surrounding literature of the subject, the Web of Science Editors may decline to include the journal in any Clarivate product or service. The Web of Science Editors, in their sole discretion, may remove titles from coverage at any point if the titles fail to maintain our standard of quality, do not comply with ethical standards, or otherwise do not meet the criteria determined by the Web of Science Editors. If a journal is deselected or removed from coverage, the journal will cease to be indexed in the Web of Science from a date determined by the Web of Science Editors in their sole discretion – articles published after that date will not be indexed. The Web of Science Editors' decision on all matters relating to journal coverage will be final.

[Clarivate.™ Accelerating innovation.](#)





Caryologia

COUNTRY United Kingdom Universities and research institutions in United Kingdom Media Ranking in United Kingdom	SUBJECT AREA AND CATEGORY Agricultural and Biological Sciences Agricultural and Biological Sciences (miscellaneous) Biochemistry, Genetics and Molecular Biology Genetics	PUBLISHER Firenze University Press Universita degli Studi di Firenze in Scimago Institutions Rankings	H-INDEX 33
PUBLICATION TYPE Journals	ISSN 00087114, 21655391	COVERAGE 1949-2023	INFORMATION Homepage How to publish in this journal caryo@unifi.it

SCOPE

Caryologia is devoted to the publication of original papers, and occasionally of reviews, about plant, animal and human karyological, cytological, cytogenetic, embryological and ultrastructural studies. Articles about the structure, the organization and the biological events relating to DNA and chromatin organization in eukaryotic cells are considered. Caryologia has a strong tradition in plant and animal cytosystematics and in cytotoxicology. Bioinformatics articles may be considered, but only if they have an emphasis on the relationship between the nucleus and cytoplasm and/or the structural organization of the eukaryotic cell.

Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS

1

Cytologia

JPN

36%

similarity

2

Genetika

SRB

29%

similarity

3

Turkish Journal of Botany

TUR

23%

similarity

4

Genetic Resources and Crop Evolution

NLD

23%

similarity













5

Revisita Botanica

CHE

https://www.scimagojr.com/journalsearch.php?q=18430&tip=sid&clean=0

1/3



Metrics based on Scopus® data as of March 2024

Loading comments...

ตอบรับการตีพิมพ์ วันที่ 29 กันยายน 2567

1 จาก 1,208

[Caryo] Editor Decision

ทดลองจดหมาย 



**Alessio Papini**
ถึง ชัน, Surachest, Kan, Alongklod 

อา. 29 ก.ย. 20:09 (13 ชั่วโมงที่ผ่านมา)    

Sumalee Phimphan, Surachest Alumsumang, Kan Khoomsab, Itsara Tangsuwan, Alongklod Tanomtong:

We have reached a decision regarding your submission to Caryologia, "Karyomorphology and microsatellites characterization of *Limnonectes gyldenstolpei* : First report from Thailand".

Our decision is to: Accept Submission

[Caryologia, International Journal of Cytology, Cytosystematics and Cytogenetics](#) | E-ISSN: 2165-5391 |
Principal contact: Alessio Papini alpapini@unifi.it | Support contact: Alessandro Pierno alessandro.pierno@unifi.it
| Published by [Firenze University Press](#) | [University of Florence](#) | [About the Journal](#) | [Editorial Team](#)



CURRENT ISSUE

JUST ACCEPTED

ALL ISSUES

ABOUT


SUBMISSIONS

POLICIES

Q SEARCH

HOME / ARCHIVES / Vol. 77 No. 3 (2024)

Vol. 77 No. 3 (2024)



PUBLISHED: 2025-03-25

FULL ISSUE

- PDF

ARTICLES

The genome of the southern short-horned tree dragon *Acanthosaura meridiona* Trivalairat, Sumontha, Kunya & Chaingkul, 2022 (Squamata, Draconinae) was analyzed using classical and molecular techniques to identify and study its chromosomal and repetitive elements

DOI: <https://doi.org/10.36253/caryologia-2961>

Praween Supanuam; Sittisak Jantararat, Thaintip Kraiprom, Somsak Buathip, Sarun Jumrusthanasan, Sarawut Kaewsri, Nattasuda Donbundit, Phichaya Buasriyot, Weera Thongnetr, Sumalee Phimphan, Alongklod Tanomtong
3-10



Amelioration strategy of saline stress in wheat with salicylic acid: a review

DOI: <https://doi.org/10.36253/caryologia-2812>

Syeda Afia Fairaj, Uttam Kumar Ghosh, Md. Moshiul Islam, Khurshida Jahan, Anamika, Sazada Siddiqui, Mohammed O. Alshaharani, Ayesha Siddiqua, Habab Merghani Yassin
11-25



Genomic in situ hybridization (GISH) and performance analysis in intergeneric hybrids from five consecutive generations of *Erianthus* x *Saccharum*

DOI: <https://doi.org/10.36253/caryologia-2727>

V.P. Sobhakumari, K. Mohanraj
27-36



Giemsa-based chromosome staining and comparative fluorescent banding pattern in five valuable Indian plant species

DOI: <https://doi.org/10.36253/caryologia-3007>

Timir Baran Jha, Mihir Halder, Biplab Kumar Bhowmick

37-45



Karyomorphology and microsatellites characterization of *Limnonectes gyldenstolpei*: first report from Thailand

DOI: <https://doi.org/10.36253/caryologia-2868>

Sumalee Phimphan, Surachest Aiumsumang, Kan Khoomsab, Itsara Tangsuwan, Alongklod Tanomtong

47-52



Chromosome, ploidy analysis, and flow cytometric genome size estimation of *Datura stramonium* and *D. innoxia* medicinal plant

DOI: <https://doi.org/10.36253/caryologia-2768>

Zahra Morovati, Ghasem Karimzadeh, Mohamad Reza Nagahavi, Sajad Rashidi Monfared

53-61



Report of genomic doubling in *Cyamopsis tetragonoloba* (L.) Taub. (Fabaceae): salient features and effects

DOI: <https://doi.org/10.36253/caryologia-2478>

Shefali Singh, Girjesh Kumar

63-71



[MAKE A SUBMISSION](#)

INFORMATION

[For Readers](#)[For Authors](#)[For Librarians](#)

Caryologia: International Journal of Cytology, Cytosystematics and Cytogenetics | E-ISSN: **2165-5391** |

contact: alessio.papini@unifi.it

Published by [Firenze University Press](#) | [University of Florence](#) | Via Cittadella, 7 - 50144 Firenze (FI) - Italy | contact:

info@fupress.com |

[Privacy policy](#) | [Terms of use](#) | [Legal notice](#) | [Accessibility Statement](#) | [Login/Registration](#) |

[Announcements](#) | [Fupress Journals](#) | [Open Access Policy of the University of Florence](#) |

The *Caryologia* site and its metadata are licensed under a [Creative Commons Attribution 4.0 International License \(CC-BY- 4.0\)](#)



Platform &
workflow by
OJS / PKP

Editorial Team

Editor in Chief

[Alessio Papini](#) - *Department of Biology*
University of Florence
Via La Pira, 4 – 0121 Firenze, Italy

Associate Editors

Alfonso Carabez-Trejo - *Mexico City, Mexico*
Katsuhiko Kondo - *Hagishi-Hiroshima, Japan*
Canio G. Vosa - *Pisa, Italy*

Subject Editors

Mycology

Renato Benesperi - *University of Florence, Italy*

Human and Animal Cytogenetics

Michael Schmid - *University of Würzburg, Germany*

Plant Cytogenetics

Lorenzo Peruzzi – *University of Pisa, Italy*

Cytotoxicity

Cleiton Pereira de Souza - *São Paulo, Brazil*

Plant Karyology and Phylogeny

Andrea Coppi – *University of Florence, Italy*

Histology and Cell Biology

Alessio Papini – *University of Florence, Italy*

Nanoparticles

Sara Falsini - University of Florence, Italy

Zoology

Mauro Mandrioli - University of Modena and Reggio Emilia, Italy

Genotoxicity

Nicoleta Anca Sutan - University of Pitesti, Romania

Editorial Assistant

Sara Falsini - University of Florence, Italy

Editorial Advisory Board

- Andrew Bajer - *University of Oregon, Eugene, USA*
 Erika Bellini - *University of Pisa, Pisa, Italy*
 Andrea Bennici - *University of Florence, Florence, Italy*
 Graziella Berta - *Amedeo Avogadro University of Eastern Piedmont, Alessandria, Italy*
 Elisabetta Bianchi - *University of Florence, Florence, Italy*
 Davide Bizzaro - *Polytechnic University of Marche, Ancona, Italy*
 Elena Bonciu - *University of Craiova, Romania*
 Antonio Brito Da Cunha - *University of Sao Paulo, Sao Paulo, Brazil*
 Ernesto Capanna - *University "La Sapienza", Rome, Italy*
 Duccio Cavalieri - *Edmund Mach Foundation, San Michele all'Adige, Italy*
 Roberto Cremonini - *University of Pisa, Pisa, Italy*
 Mauro Cresti - *University of Siena, Siena, Italy*
 Giovanni Cristofolini - *University of Bologna, Bologna, Italy*
 Paolo Crosti - *University of Milano-Bicocca, Milan, Italy*
 Giovanni Delfino - *University of Florence, Firenze, Italy*
 Saverio D'Emerico - *University of Bari, Bari, Italy*
 Fabio Garbari - *University of Pisa, Pisa, Italy*
 Antonio Giovino - *Council for Agricultural Research and Economics, Bagheria, Italy*
 Claudia Giuliani - *University of studies of Milan, Milan, Italy*
 Marcelo Guerra - *Federal University of Pernambuco, Recife, Brazil*
 Waheeb K. Heneen - *The Swedish University of Agricultural Sciences, Svalöf, Sweden*
 Alessio Iannucci - *University of Florence, Florence, Italy*
 Leopoldo Iannuzzi - *CNR-ISPAA, Napoli, Italy*
 Janusz Limon - *Medical University of Gdansk, Gdansk, Poland*
 Jiangtao Liu - *University of Lanzhou, Lanzhou, China*
 Nils Mandahl - *Lund University, Lund, Sweden*
 Mauro Mandrioli - *University of Modena and Reggio Emilia, Modena, Italy*
 Gian Carlo Manicardi - *University of Modena and Reggio Emilia, Modena, Italy*
 Palmer Marchi - *University "La Sapienza", Rome, Italy*
 Fábio Porto-Foresti - *Universidade Estadual Paulista, São Paulo, Brazil*
 Monica Ruffini Castiglione - *University of Pisa, Pisa, Italy*
 Anca Sutan - *University of Pitesti, Romania*
 Luigi Sanità di Toppi - *University of Pisa, Pisa, Italy*
 Claus Steinlein - *University of Würzburg, Würzburg, Germany*
 Joan Vallès - *University of Barcelona, Barcelona, Catalonia, Spain*

[MAKE A SUBMISSION](#)

INFORMATION

[For Readers](#)[For Authors](#)[For Librarians](#)

Caryologia: International Journal of Cytology, Cytosystematics and Cytogenetics | E-ISSN: **2165-5391** |

contact: alessio.papini@unifi.it

Published by [Firenze University Press](#) | [University of Florence](#) | Via Cittadella, 7 - 50144 Firenze (FI) - Italy | contact:

info@fupress.com |

[Privacy policy](#) | [Terms of use](#) | [Legal notice](#) | [Accessibility Statement](#) | [Login/Registration](#) |

[Announcements](#) | [Fupress Journals](#) | [Open Access Policy of the University of Florence](#) |

The *Caryologia* site and its metadata are licensed under a [Creative Commons Attribution 4.0 International License \(CC-BY- 4.0\)](#)



Platform &
workflow by
OJS / PKP



Citation: Phimphan, S., Aiumsumang, S., Khoomsab, K., Tangsuwan, I. & Tanomtong, A. (2024). Karyomorphology and microsatellites characterization of *Limnonectes gyldenstolpei*: first report from Thailand. *Caryologia* 77(3): 47-52. doi: 10.36253/caryologia-2868

Received: Jul 14, 2024

Accepted: Sep 30, 2024

Published: March 25, 2025

© 2024 Author(s). This is an open access, peer-reviewed article published by Firenze University Press (<https://www.fupress.com>) and distributed, except where otherwise noted, under the terms of the CC BY 4.0 License for content and CC0 1.0 Universal for metadata.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Competing Interests: The Author(s) declare(s) no conflict of interest.

Karyomorphology and microsatellites characterization of *Limnonectes gyldenstolpei*: first report from Thailand

SUMALEE PHIMPHAN^{1,*}, SURACHEAT AIUMSUMANG¹, KAN KHOOMSAB², ITSARA TANGSUWAN³, ALONGKLOD TANOMTONG⁴

¹ Biology Program, Faculty of Science and Technology, Phetchabun Rajabhat University, Phetchabun 67000, Thailand

² Education Science Program, Faculty of Science and Technology, Phetchabun Rajabhat University, Phetchabun 67000, Thailand

³ Natural Resources and Environmental Management Program, Faculty of Science and Technology, Phetchabun Rajabhat University, Phetchabun 67000, Thailand

⁴ Department of Biology, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand

*Corresponding author. E-mail: Sumalee.phi@pcru.ac.th, joodoof@gmail.com

Abstract. In the present investigation, karyotype and microsatellites pattern in the chromosome of Gyldenstolpe's Frog (*Limnonectes gyldenstolpei*) have been analyzed. The aspect of chromosome numbers, morphology, nucleolus organizer region (NOR) locations and microsatellites pattern [d(CA)₁₅, d(CGG)₁₀, d(GC)₁₅, d(TA)₁₅]. We provided the karyotype and idiogram of this species by conventional staining, Ag-NOR banding and Fluorescence *in situ* hybridization techniques. For the study, five male and female samples collected from northern Thailand, were used. The metaphase chromosome preparations were prepared from the bone marrows by the standard protocol. The result shows that *L. gyldenstolpei* had the diploid chromosome number (2n) was 26 and the fundamental number (NF) were 56 in both males and females. The karyotype is composed of 4 large metacentric, 4 large submetacentric, 2 medium metacentric, 14 small metacentric and 2 small submetacentric chromosomes. The NORs bearing chromosome were in close to the telomere region on chromosome pair 1. In addition, the microsatellite d(CGG)₁₀ and (GC)₁₅ hybridization results confirmed the NOR region. The *in situ* localization pattern of d(CA)₁₅ microsatellites was positive on all telomere chromosome, while microsatellites d(TA)₁₅ have no signal on chromosome. Here we provide a classical and some molecular genetics information for *L. gyldenstolpei* useful as a species specific marker.

Keywords: *Limnonectes gyldenstolpei*, karyotype, chromosome, microsatellites.

INTRODUCTION

Limnonectes gyldenstolpei is a species of frog in the Dicroglossidae family. It has been recorded throughout much of Thailand, northeastern Lao,

southwestern Cambodia, and central Vietnam. It has recently also been recorded from the Phong Nha-Kẻ Bàng National Park in central Vietnam (Luu et al., 2013). The members of the genus *Limnonectes* have a broad distribution in Asia from eastern and southern China, eastwards to Japan, throughout Indochina and southwards to Malaysia, Indonesia, Philippines, and New Guinea (Frost, 2016). *Limnonectes* is one of the most diverse groups amphibians with 69 currently species recognized and 15 of which have been described in the last ten years (Frost, 2016).

The gross chromosome numbers of 1,000 amphibian species were reported by Kuramoto (1990). But only 837 of the 3,521 anuran species have been analyzed chromosomally (King, 1991). The amphibian fauna in Thailand comprises of 176 species in 8 families and 3 orders (Khonsue and Thirakhupt, 2001). In the genus *Limnonectes* reported 11 species (Niyomwan, et al., 2019). The list shows chromosome number variation occurs in most of the seven families of anuran amphibians classified with 33 genera. The typical karyotype of the family Dicroglossidae is diploid chromosome number ($2n$)=22, 24 and 26. For the genus *Limnonectes*, there were some cytogenetic studies reported the diploid number was $2n$ =22-26, NF =44-52, including *L. kuhlii* and *L. blythii* (Supaprom, 2003), *L. pileatus* (Supaprom, 2003; Supaprom and Baimai, 2004), *L. gruniens* and *L. modestus* (Nasaruddin, 2009), *L. blythii* (Donsakul and Rangsiruji, 2005; Phimphan et al., 2020) and *L. taylori* (Phimphan and Aiumsumang, 2019). All previous knowledge demonstrated that there are several patterns of chromosomes (number, type, size).

This is the first report describing the molecular cytogenetic and karyotype study of chromosome size, standardized idiogram, karyotype formula and meiotic cell division of the *L. gyldenstolpei* species. The molecular data, microsatellite probes are used to detect if there is some specific hybridization pattern in *L. gyldenstolpei* has not been studied yet. The results obtain can be fulfilled to the basic knowledge. In addition, our knowledge advances cytogenetic information for further study on taxonomy relationship. Moreover, we provide useful basic information for the conservation and chromosome evolution study of this frog.

MATERIAL AND METHODS

Field surveys were conducted in rainy season from northern (16.42°N 101.16°E), Thailand. Five males and five females of *L. gyldenstolpei* were mature obtained during. The frogs were transferred to the laboratory and were kept under standard conditions for 3 days before the

experimentation. Experiments were performed in accordance with ethical protocols (Ref No. U1-04498-2559). The chromosomes were prepared *in vivo* with slight adaptations as follows (Sangpakdee et al., 2016). The colchicine was injected into the frogs' abdominal cavity. Then, the frogs were left in a box for eight hours and then killed. The bone marrow was collected by cutting the head and the end of femurs and tibias, and then a syringe was used to inject 0.075 M KCl into the marrow to drive out the bone marrow tissue or cells into the plate. We gently cut the tissue to pieces as small as possible. We transferred 8 mL of cell sediments to a centrifuge tube and incubated it for 30 min at 37 °C. After centrifugation at 1500 rpm for 8 min, the KCl was discarded. Cells were fixed in fresh cool fixative up to 8 mL by gradually adding it before being centrifuged again at 1500 rpm for 8 min. The fixation was repeated until the supernatant was clear, usually three times. Finally, the pellet was mixed with 1 mL fixative (depending on the amount of cell). The mixture was dropped onto a clean and cold slide by a micropipette, and then the air-dry technique was applied.

Conventional staining was done using 10% Giemsa's solution for 10 min (Phimphan and Aiumsumang, 2019). Ag-NOR banding was performed (Howell and Black, 1980) by applying two drops of 2% gelatin on the slides, followed with four drops of 50% silver nitrate. The slides were then covered with a cover slip and incubated at 60°C for 5 min or until the slide changed brownish. After that the slides were dipped in distilled water to remove the cover glass and air-dried on the slide. The microsatellites (CA)₁₅, (CGG)₁₀, (GC)₁₅, and (TA)₁₅ were synthesized according to (Kubat et al., 2008; Supiwong et al., 2014). These sequences were directly labeled with Cy₃ at the 5'terminus during synthesis by Sigma (St. Louis, MO, USA).

Chromosome counting was performed on mitotic metaphase cells under a light microscope. Twenty clearly observable and well-spread chromosomes of each male and female were selected and photographed. The length of the short arm chromosome (Ls) and the length of the long arm chromosome (Ll) were measured, and the length of the total arm chromosome (LT, $LT = Ls + Ll$) calculated. The relative length (RL), the centromeric index (CI) and standard deviation (SD) of RL and CI were estimated. The CI ($q/p+q$) between 0.50–0.59, 0.60–0.69, 0.70–0.89 and 0.90–0.99 were described as metacentric, submetacentric, acrocentric and telocentric chromosomes, respectively (Levan, 1964). The fundamental number (number of chromosome arm, NF) was obtained by assigning a value of two to metacentric, submetacentric and acrocentric chromosomes and one to telocentric chromosome. All parameters were used in karyotyping and idiogramming.

RESULTS AND DISCUSSION

The results showed *L. gyldenstolpei* had diploid chromosome number of $2n=26$ and fundamental number (NF)=52, the karyotype comprised four large metacentric, four large submetacentric, two medium metacentric, 14 small metacentric and two small submetacentric chromosomes. The karyotype formula of *L. gyldenstolpei* is $2n(26)=L^m_4+L^{sm}_4+M^m_2+S^m_{14}+S^{sm}_2$ in both males and female, while sex chromosomes were cytologically indistinguishable (Fig. 1A). The average lengths of each chromosome including short and long arm length, total length, relative length, and centromeric index were calculated and presented in Table 1. The previous relevant literatures have been reported that the numbers of diploid chromosome and fundamental number in *Limnonectes* studied herein are $2n=22-26$ and $NF=44-52$ including, *L. kuhlii* and *L. blythii* (Supaprom, 2003), *L. pileatus* (Supaprom, 2003; Supaprom and Baimai, 2004), *L. gruniens* and *L. modestus* (Nasaruddin, 2009), *L. blythii* (Donsakul and Rangsiruji, 2005; Phimphan et al., 2020) and *L. taylori* (Phimphan and Aiumsumang, 2019). (Table 2). Comparison to closely related species, *L. gyldenstolpei* had diploid chromosome number similar to *L. gruniens* and *L. modestus* ($2n=24$), but is higher than that in *L. taylori* ($2n=22$) and lower than *L. kuhlii* and *L. pileatus* ($2n=24$). This result was the first report on *L. gyldenstolpei*. These characteristics are consistent with the theory that reorganization from the original karyotype resulted from Robertsonian fissions, fusions, or pericentric inversions (Gorman 1973; King 1978). Our results confirmed $2n$ for *L. gyldenstolpei* species but with differences in the diploid chromosome number. This incongruence reflects probably the number of the *Limnonectes* chromosomes, especially those of polyploids.

After Ag-NOR staining, these regions produce numerous gene expressions and contain more non-histone protein than others regions on the chromosome. Accordingly, the dark band (NOR-positive) is induced by the reduction of organic silver by these proteins that change from silver to dark (Sharma et al., 2002). If these regions were active during the interphase prior to mitosis, they can be detected by silver nitrate staining (Howell and Black 1980). The NOR could be detected to near telomeric region on long arm chromosome pairs 1 (Fig. 1B). We found one pair of Ag-NOR sites in all of the samples examined. However, the results were similar to the previous report on *L. kuhlii* and *L. blythii* (Supaprom, 2003), *L. pileatus* (Supaprom, 2003; Supaprom and Baimai, 2004), *L. gruniens* and *L. modestus* (Nasaruddin, 2009), *L. blythii* (Donsakul and Rangsiruji, 2005; Phimphan et al., 2020) and *L. taylori* (Phimphan and Aium-




Figure 1. Metaphase chromosome plates and karyotypes of *Limnonectes gyldenstolpei*, $2n=26$ by conventional staining [A.], Ag-NOR banding [B.], $d(CA)_{15}$ [C.], $d(CGG)_{10}$ [D.], $d(GC)_{15}$ [E.] and $d(TA)_{15}$ microsatellite probe [F.]. Note scale bars indicate 10 micrometers.

sumang, 2019). The most striking variation is seen in the morphology of the secondary constrictions. Generally, one major nucleolar organizer region is present per genome (n), which may vary in its position between species. However, closely related and often morphologically very similar species share the same type and location of their nucleolar organizer regions, which can therefore provide an effective taxonomic.

Here the first molecular cytogenetic study in metaphase chromosomes stained by FISH. The *in situ* hybridized localization of microsatellites $d(CA)_{15}$, $d(CGG)_{10}$, $d(GC)_{15}$, and $d(TA)_{15}$. Microsatellites, also known as simple sequence repeats, consist of very short motifs (1-6 nucleotides in length) repeated in tandem arrays. Generally, they are located in the heterochromatic regions (telomeres, centromeres and in the sex chromosomes) of genomes, where a significant fraction of repetitive DNA is expected to be localized (Supiwong et al., 2013). The

Table 1. Mean length of short arm chromosome (Ls), length of long arm chromosome (Ll), length of total chromosomes (LT), relative length (RL), centromeric index (CI) and standart deviation (SD) from 20 metaphases of male and female *Limnonectes gyldenstolpei*, $2n$ (diploid)=26.

Chromosome pairs	Ls	Ll	LT	CI±SD	RL±SD	Chromosome size	Chromosome type
1*	6.162	7.628	13.790	0.552±0.012	0.161±0.005	Large	metacentric
2	4.166	6.849	11.015	0.622±0.014	0.128±0.004	Large	submetacentric
3	3.469	5.991	9.460	0.633±0.019	0.110±0.004	Large	submetacentric
4	3.786	5.168	8.954	0.580±0.022	0.105±0.002	Large	metacentric
5	3.700	4.503	8.203	0.550±0.018	0.096±0.003	Medium	metacentric
6	2.179	2.966	5.145	0.577±0.014	0.060±0.002	Small	metacentric
7	1.866	2.892	4.758	0.606±0.032	0.056±0.002	Small	submetacentric
8	1.876	2.799	4.675	0.597±0.037	0.054±0.002	Small	metacentric
9	1.766	2.590	4.356	0.595±0.017	0.051±0.002	Small	metacentric
10	1.799	2.475	4.274	0.577±0.021	0.050±0.002	Small	metacentric
11	1.609	2.387	3.996	0.596±0.015	0.047±0.001	Small	metacentric
12	1.505	2.260	3.765	0.598±0.018	0.044±0.001	Small	metacentric
13	1.486	1.967	3.452	0.572±0.024	0.040±0.002	Small	metacentric

* NORs bearing chromosomes (satellite chromosome).

Table 2. Review of cytogenetic publications of family Dicroglossidae (genus *Limnonectes*).

Species	$2n$	Karyotype formula	NF	NORs	FISH	Reference
<i>L. gruniens</i>	24	24m	48	-	-	Nasaruddin et al. 2009
<i>L. modestus</i>	24	20m+4t	44	-	-	Nasaruddin et al. 2009
<i>L. kuhlii</i>	26	8m+14sm	52	2	-	Supaprom 2003
<i>L. pileatus</i>	26	16m+10sm	52	2	-	Supaprom 2003
	26	16m+10sm	52	2	-	Supaprom and Baimai 2004
<i>L. taylori</i>	22	16m+6sm	44	2	-	Phimphan and Aiumsumang 2019
	24	10m+12sm+2a	48	-	-	Donsakul and Rangsiruji 2005
<i>L. blythii</i>	24	20m+4sm	48	2	+	Phimphan et al. 2021a
<i>L. gyldenstolpei</i>	26	20m+6sm	52	2	+	Present study

$2n$ diploid chromosome number, NF=fundamental number (number of chromosome arms), *m* metacentric, *sm* submetacentric, *a* acrocentric, *t* telocentric chromosome, NORs Ag-NOR banding, FISH Fluorescence in situ hybridization, + positive and - not available.

result of *L. gyldenstolpei* analyzed was being abundantly distributed on all telomere chromosomes such as, the accumulation of (CA)₁₅ in long arm chromosomal pair 1 (Fig 1C), while (CGG)₁₀ and (GC)₁₅ detected subtelomeric region on long arm chromosomal pair 1 (Fig. 1D, 1E) and (TA)₁₅ sequences are not present in the all chromosome (Fig. 1F). However, an intriguing feature exclusive for *L. gyldenstolpei* was the strong accumulation of all microsatellites at the regions of specific chromosomal pair, indicating that these microsatellites may be used as chromosomal markers in this frog species. In the frog genomes, microsatellites are usually abundant in the telomeric and centromeric regions, Otherwise, the dinu-

cleotides (CA)₁₅, (GC)₁₅ and (CGG)₁₀ accumulated exclusively in telomeric and subcentromeric chromosomal regions, corroborating findings from other frog groups studied to date (Phimphan, et al. 2021a; 2021b). These molecular cytogenetics data could also be a substantial prerequisite for future frog genome projects. This study discovered that the cytogenetic maps of *L. gyldenstolpei* allowed us to map out the steps involved in this species' chromosomal rearrangement. This is the first report on the Fluorescence *in situ* hybridization (FISH) study of this species in Thailand.

The present study on the meiotic cell division of *L. gyldenstolpei* found that during interphase, nucleolus




Figure 2. Meiosis cell division of the *Limnonectes gyldenstolpei*, $2n=26$ on interphase (A.), leptotene (B.), pachytene (C.), diakinesis (D.), metaphase I (E.) and metaphase II (F.) by conventional staining technique. Scale bars indicate 10 micrometers.

could be clearly seen, while chromatin was absent. In prophase, metaphase I (meiosis I) the homologous chromosomes showed synapsis, which can be defined as the 13 bivalent and 13 haploid chromosomes at metaphase II as diploid species. It is confirmed for this species had $2n=26$ in similar to previous reports. The largest metacentric chromosome pair 1 is the largest bivalent. We found that *L. gyldenstolpei* had the distinct character of the observable leptotene (initiation of chromosome shrinking), pachytene (completion of chromosome synapsis) and diakinesis (terminalization) according to Patawang (Patawang et al., 2013) (Fig. 2). In conclusion, this study provides the first chromosome, molecular cytological details and Ag-NOR marker for *L. gyldenstolpei* from Thailand. The results support the karyotype of genus *Limnonectes* are conserved among several other species. However, the chromosomal morphology may be slightly different depending on populations of *L. gyldenstolpei* present in different countries. Our results added new knowledge that can be used for karyological comparative analyses in *Limnonectes* species, on the basis of classical and banding approach within this taxon.

ACKNOWLEDGMENTS

This study was supported by the National Research Council of Thailand under the Phetchabun Rajabhat University (Grant No. TSRI2567/64).

REFERENCES

- Donsakul T, Rangsiruji A. 2005. Liver karyotypes in *Limnonectes blythii*, *Rana erythraea*, *Rana leptoglossa*, *Occidozyga martensii* and *Glyphoglossus molossus* (Amphibia, Anura). Proceedings of 43rd Kasetsart University Annual Conference, Thammasart University, Bangkok.
- Frost DR. 2016. Amphibian Species of the World: an online reference. Version 6.0. American Museum of Natural History, New York, USA.
- Gorman GC. 1973. The chromosomes of the Reptilia, a cytotaxonomic interpretation. Cytotaxonomy and vertebrate evolution. New York: Academic Press, p. 349–424.
- Howell WM, Black DA. 1980. Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: a 1-step method. *Experientia*. 36: 1014–5.
- King M. 1978. A new chromosome form of *Hemidactylus frenatus* (Dumeril and Bibron). *Herpetologica* 34: 216–218.
- Khonsue W, Thirakhupt K. 2001. A Checklist of the Amphibians in Thailand. *Trop Nat Hist*. 1: 69–82.
- Kubat Z, Hobza R, Vyskot B, Kejnovsky E. 2008. Microsatellite accumulation in the Y chromosome of *Silene latifolia*. *Genome*. 51: 350–356.
- Kuramoto M. 1980. Karyotype of several frogs from Korea, Taiwan and Philippines. *Experientia*. 36: 826–828.
- Levan A, Fredga K, Sandberg AA. 1964. Nomenclature for centromeric position on chromosomes. *Hereditas*. 52: 201–220.
- Luu, VQ, Truong QN, Cuong TP, Kien ND, Thanh NV, Sladjana M, Michael B, Thomas Z, 2013. No end in sight? Further new records of amphibians and reptiles from Phong Nha – Ke Bang National Park, Quang Binh Province, Vietnam”. *Biodiversity*. 4(2): 285–300.
- Nasaruddin, Suriana, Dwi Arinto Adi, Salamansyah. 2009. The Karyotype of Seven Species of Amphibians (Anuran Order) from South-east Sulawesi. *Veteriner*. 10: 77–76.
- Niyomwan P, Priwan S, Pawangkanan P. 2019. Amphibians of Thailand. Bangkok. 487 pp.

- Patawang I, Tanomtong A, Phimphan S, Chuaynkern Y, Chuaynkern C, Phaengphairee P, Khrueanet W, and Nithikulworawong N. 2013. The identification of sex-chromosomes and karyological analysis of rice frog, *Fejervarya limnocharis* (Anura, Ranidae) from Northeast Thailand. *Cytologia*. 79:141–150.
- Phimphan S, Aiumsumang S. 2019. Chromosomal characteristics of Taolor's stream frog (*Limnonectes taylori*) (Amphibia, Anura) from Thailand. *The Nucleus*. 64(1): 129–133.
- Phimphan S, Aiumsumang S, Tanomtong A, Jantararat S. 2021a. Karyomorphological delineation and linear differentiation of microsatellite patterns, and meiosis in giant Asian river frog (*Limnonectes blythii*) from Thailand. *The Nucleus* 64(2): 249–254.
- Phimphan S, Aiumsumang S, Tanomtong, A. 2021b. Characterization of Chromosome and Repetitive elements in the Genome of Frog *Rana nigrovittata* (Anuran, Ranidae) from Thailand: Revealed by conventional, Ag-NOR banding, C-banding and FISH techniques. *Cytology and Genetics*. 55(6): 583–589.
- Sangpakdee W, Phimphan S, Tengjaroenkul B, Pinthong K, Neeratanaphan L, Tanomtong, A. 2016. Cytogenetic Study of Three Microhylid Species (Anura, Microhylidae) from Thailand. *Cytologia*. 82: 67–74.
- Sharma OP, Tripathi NK, Sharma KK. 2002. A review of chromosome banding in fishes. In: Sobti, RC (ed.). Some aspects of chromosome structure and functions. Narosa Publishing House, New Delhi.
- Supaprom T. 2003. Cytogenetics of Amphibians in Thailand. Ph.D. Dissertation, Mahidol university.
- Supaprom T, Baimai V. 2004. Karyotypes of ten species of ranid frogs (Anura: Ranidae) from Thailand. *Amphibia-Reptilia*. 25:104–111.
- Supiwong W, Liehr T, Cioffi MB, Chaveerach A, Kosyakova A, Pinthong K, Tanee T, Tanomtong A. 2013. Karyotype and cytogenetic mapping of 9 classes of repetitive DNAs in the genome of the naked catfish *Mystus bocourti* (Siluriformes, Bagridae). *Molecular Cytogenetics*. 6:51
- Supiwong W, Liehr L, Cioffi MB, Chaveerach A, Kosyakova N, Pinthong K, Tanee T, Tanomtong, A. 2014. Chromosomal evolution in naked catfishes (Bagridae, Siluriformes): A comparative chromosome mapping study. *Zool. Anz*. 253: 316–320.