



atom together <atom2414@gmail.com>

Fill final version to template and Copyright assignment form Science & Technology Asia

Science & Technology Asia <sciencetechnologyasia@gmail.com>

28 มีนาคม 2562 13:53

ถึง: atom together <atom2414@gmail.com>

Dear Dr. Wilaiporn Pongpian,

Your article title: **Analysis of Chemical Properties and Gamma - Oryzanol Content in Luem Pua Rice Bran Oil (Oryza sativa L.)** has been accepted in Science & Technology Asia Vol.24 No.2 (April-June 2019)

Please, return to us

1. fill up Copyright assignment form.


Sincerely
Chatchada

Coordinator
Science & Technology Asia
Research Administration Division
Thammasat University (Rangsit Campus)
Klong Nueng, Klong Luang, Pathum Thani 12120 Thailand
0-2564-4440-79 Ext. 1810
Coordinator
Science & Technology Asia
Research Administration Division
Thammasat University (Rangsit Campus)
Klong Nueng, Klong Luang, Pathum Thani 12120 Thailand
0-2564-4440-79 Ext. 1810

ในวันที่ อ. 26 มี.ค. 2019 เวลา 11:38 atom together <atom2414@gmail.com> เขียนว่า:
[ข้อความที่เกี่ยวข้องถูกซ่อนไว้]

 STA Copyright Transfer Form.pdf
44K

สำเนาถูกต้อง



SCIENCE & TECHNOLOGY ASIA

A Publication of Thammasat University, Thailand

(Formerly Thammasat International Journal of Science and Technology)

ISSN 2586-900

ISSN 2586-902

Vol. 24 No. 2 April – Jun

สีเทา
สีเทา
(ทาสีเทา สีเทา สีเทา)

2019

Editorial Team

Editors

Charturong Tantibundhit	Thammasat University	Thailand
Kesara Na-Bangchang	Thammasat University	Thailand
Wutiphol Sintunavarat	Thammasat University	Thailand

Associate Editors

Poom Kumam	King Mongkut's University of Technology Thonburi	Thailand
Wararit Panichkitkosolkul	Thammasat University	Thailand

Editorial Board Members

Andrew Owen	University of Liverpool	United Kingdom
Ariana Pitea	University Politehnica of Bucharest	Romania
Bashir Ahmad	King Abdulaziz University	Saudi Arabia
Carol Sibley	University of Washington	United States of America
Cecilia C. Maramba-Lazarte	UP College of Medicine	Philippines
Dhananjay Gopal	S.V. National Institute of Technology	India
Franz Pernkopf	Graz University of Technology	Austria
Gao CHEN	Chinese Academy of Sciences	China
Hassen Aydi	Imam Abdulrahman Bin Faisal University	Arabia
Janusz Brzdek	Pedagogical University	Poland
Jessada Tariboon	King Mongkut's University of Technology North Bangkok	Thailand
Juan Martinez-Moreno	University of Jaen	Spain
Johann Schinnert	University of Vienna	Austria
L. Jeyaseelan	Christian Medical College Vellore	India

Mitsuru Ikeda	Japan Advanced Institute of Science and Technology	Japan
Nguyen Van Dung	Ton Duc Thang University	Vietnam
Paul Haddad	University of Tasmania	Australia
Peerasak Intarapaiboon	Thammasat University	Thammasat University
Penwisa Pisitsak	Thammasat University	Thailand
Phadungsak Ratanadecho	Thammasat University	Thailand
Saifon Chaturantabut	Thammasat University	Thammasat University
Sarute Ummartyotin	Thammasat University	Thailand
Sotiris K. Ntouyas	Ntouyas, University of Ioannina	Greece
Suthep Suantai	Chiang Mai University	Thailand
Wai Shiu Fred WONG	National University of Singapore	Singapore
Xianghai CAI Kunming	Chinese Academy of Sciences	China
Yeol-je Cho	Gyeongsang National University	Korea

Home ThaiJo

THAIJO (<https://www.tci-thaijo.org/>)

Indexed In



ASEAN
CITATION
INDEX

(<https://asean-cites.org/index.php?r=contents%2Findex&id=9>)



(https://www.kmutt.ac.th/jif/public_html/list%20journal.php)



Elektronische
Zeitschriftenbibliothek

([https://ezb.uni-regensburg.de/searchres.phtml?](https://ezb.uni-regensburg.de/searchres.phtml?bibid=AAAAA&colors=7&lang=de&jq_type1=QS&jq_term1=Thammasat+International+Journal)

[bibid=AAAAA&colors=7&lang=de&jq_type1=QS&jq_term1=Thammasat+International+Journal](https://ezb.uni-regensburg.de/searchres.phtml?bibid=AAAAA&colors=7&lang=de&jq_type1=QS&jq_term1=Thammasat+International+Journal)



([https://scholar.google.co.th/scholar?](https://scholar.google.co.th/scholar?as_q=&as_epq=&as_oq=&as_eq=&as_occt=any&as_sauthors=&as_publication=Science+%26)

[as_q=&as_epq=&as_oq=&as_eq=&as_occt=any&as_sauthors=&as_publication=Science+%26](https://scholar.google.co.th/scholar?as_q=&as_epq=&as_oq=&as_eq=&as_occt=any&as_sauthors=&as_publication=Science+%26)

Information

For Readers (<https://www.tci-thaijo.org/index.php/SciTechAsia/information/readers>)

For Authors (<https://www.tci-thaijo.org/index.php/SciTechAsia/information/authors>)

For Librarians (<https://www.tci-thaijo.org/index.php/SciTechAsia/information/librarians>)

Current Issue

 (<https://www.tci-thaijo.org/index.php/SciTechAsia/gateway/plugin/WebFeedGatewayPlugin/atom>)

 (<https://www.tci-thaijo.org/index.php/SciTechAsia/gateway/plugin/WebFeedGatewayPlugin/rss2>)

สวัสดีค่ะ
สวัสดีค่ะ
(สวัสดีค่ะ)



(<https://www.tci-thaijo.org/index.php/SciTechAsia/gateway/plugin/WebFeedGatewayPlugin/rss>)

Research Administration Division, Office of the Rector Building
Thammasat University, Rangsit Campus
99 Khlong Nueng, Khlong Luang, Pathum Thani 12120, Thailand
Tel: +66-2-564 4443 Ext.1810
Email: sciencetechnologyasia@gmail.com

Home (<https://www.tci-thaijo.org/index.php/SciTechAsia/index>)

/ Archives (<https://www.tci-thaijo.org/index.php/SciTechAsia/issue/archive>)

/ Vol.24 No.2 (April - June 2019)



(<https://www.tci-thaijo.org/index.php/SciTechAsia/issue/view/14211>)

Published: 2019-05-23

Analysis of Chemical Properties and Gamma - Oryzanol Content in Luen Pua Rice Bran Oil (*Oryza sativa* L.) (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190599>)

Wilaiporn Pongpian, Sasikarn Panpraneecharoen

1-5

Wilaiporn Pongpian
Sasikarn Panpraneecharoen
(Thammasat International Journal of Science and Technology)

PDF (662KB) (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190599/1333>)

Effect of Ammonium Sulfate on the Porous Creating of Regenerated Cellulose Hydrogel from Palm Oil Trunk (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190603>)

Nantharat Phruksaphithak, Anuwat Sukthong, Warintorn Muanpannarai, Atima Theptong

6-14

PDF (1MB) (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190603/1333>)

Erector Spinae Muscle Activity during Lifted People with Two Persons in Lifting Phase of Manual Human Handling (MHH) (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190616>)

Chanya Jiemjai, Santhanee Khruakhorn, Patcharee Kooncumchoo

15-19

PDF (514KB) (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190616/133>)

Non-Scanning Acquisition Technique for Extracting Small Depth Difference on the Area of Interest (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190621>)

Suchada Rianmora, Maroay Phlernjai

20-33

PDF (1MB) (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190621/1333>)

Using Adaptive Integral Gain for Overshoot Reduction in PID Control Systems (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190623>)

Pinit Ngamsom

34-44

PDF (813KB) (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190623/133>)

Comparatively Rapid Screening Tests for Diagnosis of Hepatitis B Virus Infection Using Loop-Mediated Isothermal Amplification (LAMP) Paired with Lateral Flow Dipstick (LFD), Gold Nanoparticles (AuNPs) and Real-Time Turbidimetry (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190624>)

Suphitcha Augkarawaritsawong, Surangrat Srisurapanon, Sirirat Wachiralurpan, Supatra Areeki Kosum Chansiri

45-57

PDF (1MB) (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190624/1333>)

Determination of Collagen Content and Antioxidant Activity in *Sesbania grandiflora* (L.) Extracts (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190631>)

Sathaporn Satsue, Wilailak Suksai, Kanaporn Tansriprapasiri, Kritsana Pachuen, Napasorn Kummeekorn, Watcharin Phuttasanbundid, Sunisa Sanlad

PDF (777KB) (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190631/133>)

Intron Length Polymorphism in Candidate Genes for Secondary Growth its Application in Diversity Assessment of Amazonian Accessions of *Hevea brasiliensis* (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190630>)

Gunlayarat Bhusudsawang, Kittipat Ukoskit

66-75

PDF (862KB) (<https://www.tci-thaijo.org/index.php/SciTechAsia/article/view/190630/133>)

Home ThaiJo

THAIJO (<https://www.tci-thaijo.org/>)

Indexed In



ASEAN
CITATION
INDEX

(<https://asean-cites.org/index.php?r=contents%2Findex&id=9>)



(https://www.kmutt.ac.th/jif/public_html/list%20journal.php)

ศูนย์ดัชนีการอ้างอิงวารสารไทย
(ศูนย์ดัชนีการอ้างอิงวารสารไทย)



Elektronische
Zeitschriftenbibliothek

([https://ezb.uni-regensburg.de/searchres.phtml?](https://ezb.uni-regensburg.de/searchres.phtml?bibid=AAAAA&colors=7&lang=de&jq_type1=QS&jq_term1=Thammasat+International+Journal)

[bibid=AAAAA&colors=7&lang=de&jq_type1=QS&jq_term1=Thammasat+International+Journal](https://ezb.uni-regensburg.de/searchres.phtml?bibid=AAAAA&colors=7&lang=de&jq_type1=QS&jq_term1=Thammasat+International+Journal)



([https://scholar.google.co.th/scholar?](https://scholar.google.co.th/scholar?as_q=&as_epq=&as_oq=&as_eq=&as_occt=any&as_sauthors=&as_publication=Science+%26)

[as_q=&as_epq=&as_oq=&as_eq=&as_occt=any&as_sauthors=&as_publication=Science+%26](https://scholar.google.co.th/scholar?as_q=&as_epq=&as_oq=&as_eq=&as_occt=any&as_sauthors=&as_publication=Science+%26)

Information

For Readers (<https://www.tci-thaijo.org/index.php/SciTechAsia/information/readers>)

For Authors (<https://www.tci-thaijo.org/index.php/SciTechAsia/information/authors>)

For Librarians (<https://www.tci-thaijo.org/index.php/SciTechAsia/information/librarians>)


Current Issue

 (<https://www.tci-thaijo.org/index.php/SciTechAsia/gateway/plugin/WebFeedGatewayPlugin/atom>)

 (<https://www.tci-thaijo.org/index.php/SciTechAsia/gateway/plugin/WebFeedGatewayPlugin/rss2>)

 (<https://www.tci-thaijo.org/index.php/SciTechAsia/gateway/plugin/WebFeedGatewayPlugin/rss>)

Research Administration Division, Office of the Rector Building
Thammasat University, Rangsit Campus
99 Khlong Nueng, Khlong Luang, Pathum Thani 12120, Thailand
Tel: +66-2-564 4443 Ext.1810
Email: sciencetechnologyasia@gmail.com



Analysis of Chemical Properties and Gamma - Oryzanol Content in Luem Pua Rice Bran Oil (*Oryza sativa* L.)

Wilaiporn Pongpian* and Sasikarn Panpraneecharoen

Chemistry Program, Faculty of Science and Technology, Phetchabun Rajabhat University,
Phetchabun 67000, Thailand.

Received 26 February 2019; Received in revised form 13 February 2019
Accepted 28 February 2019; Available online 23 May 2019

ABSTRACT

The study investigated the soxhlet extraction of Luem Pua rice bran oil (*Oryza sativa* L.) using 12 hours of extraction time, with comparison of product with hexane and ethanol as solvent. The chemical properties and gamma-oryzanol content in the rice bran oil were investigated. The results showed that the highest yield of rice bran oil was obtained from extraction using 70:30 %v/v ethanol:water solvent with percent yield of 21.89 ± 0.35 g (mean $\pm$ SD)/100g of dry rice bran) and maximum amount of gamma-oryzanol of 0.4647 mg/L of rice bran oil. The chemical properties and gamma-oryzanol content of Luem Pua rice bran oil indicates acceptable quality of oil when stored for a long time causing foul odour due to lipid peroxidation.

Keywords: Rice bran oil; Luem pua rice; Gamma-oryzanol; Peroxide value; Acid value

1. Introduction

Rice bran oil is extracted from the hard outer brown layer of rice husk in the rice bran milling process. Rice bran oil is an edible oil that is popular in Asian countries such as Japan, India, Korea, China, and Indonesia as a cooking oil [1]. It has been found that rice bran oil is suitable for cooking due to its high smoke point and mild flavor. Rice bran oil has consistency of

several phytochemical compounds such as tocotrienols, tocopherols, phytosterols, polyphenols, squalene, and gamma-oryzanol that are reported to have beneficial effects to human health [2]. In recent years, research has been done to develop various methods for extracting rice bran oil from rice bran. Gamma-oryzanol is a potent antioxidant component present only in rice bran oil that has been reported to have positive biological

*Corresponding author: atom2414@gmail.com

effects, such as decreasing the level of lowdensity lipoprotein (LDL), lowering cholesterol level, reducing blood pressure, and preventing colorectal cancer, and can be improved with pharmaceutical dosage forms, cosmetic formulations, or food products [3]. The gamma-oryzanol value demonstrated is dependent on species of rice, cultivation area and the method of extraction [4]. Massarolo et al. evaluated the effect of the solid state cultivation time of rice bran by *Rhizopus oryzae* on gamma-oryzanol recovery and its antioxidant properties. They found that the solid state cultivation leads to an increase in the gamma-oryzanol recovery followed by improvement in the functional properties of rice bran [5]. Leum Pua is native Thai sticky rice that contains antioxidants higher than white rice and other colored rice. Pornputtapitak et al. found that the lipid components in Leum Pua rice bran oil consisted of oleic acid, linoleic acid, palmitic acid and gamma-oryzanol [6]. Moreover, Srisuwan et al. suggested that Luem Pua rice might be developed as a nutraceutical for improving mood and brain function, especially learning and memory [7].

Thus, the aim of this research was to compare the three solvent extraction of Luem Pua rice bran from Phetchabun province. The chemical properties and gamma-oryzanol of extracted Luem Pua rice bran oil were investigated.

2. Materials and Methods

2.1 Material

Leum Pua rice bran (*Oryza sativa* L.), provided by a local rice mill from Amphoe Khao Kho, Phetchabun Province, Thailand, was used as a raw material for the oil extraction and gamma oryzanol analysis. Gamma-Olyzanol standard was purchased from Sigma-Aldrich (Japan). Hexane and ethanol were purchased from Labscan (Thailand).

2.2 Oil extraction of Leum Pua rice bran

The 25 g on a dry weight basis of Leum Pua rice bran was extracted with 200 ml of different solvents, hexane, ethanol and 70:30 %v/v ethanol:water by using Soxhlet apparatus for 12 hr; after that the solvent was evaporated under vacuum. Then, the yield of extracted oil was calculated by using Eq 1. The dried material was sampled in triplicate.

$$\% \text{ yield} = \frac{\text{weight of extracted oil (g)}}{\text{weight of dried seed (g)}} \times 100 \dots (1)$$

2.3 Characterization of Leum Pua rice bran oil

Acid value (AV), free fatty acid (FFA) content, and peroxide value (PV) were determined following AOCS official methods (AOCS, 2004). FFA was calculated as oleic acid and expressed as percentage of the total lipids. Gamma-oryzanol content in rice bran oil was determined by using a spectrophotometer (Specord, USA) that was slightly modified following the method of Thanonkaew et al. [8]. Consecutively, rice bran oil samples were diluted in iso-propanol. The content of gamma-oryzanol was determined by measurement of UV absorption at 320 nm and was quantified against the standard curve. The calibration curve was obtained with gamma-oryzanol standard in a concentration range of 1-5 mg/L which was adjusted to 10 ml in iso-propanol.

The data were presented as the mean values and standard deviation (SD).

3. Results and Discussion

3.1 Oil content

Previous research reported the commercial rice bran contains 15–20% of oil and also an endogenous lipase which degrades the oil and produces free fatty acids [9]. In this study, we used the hexane, ethanol and 70:30 %v/v ethanol:water as the extraction solvents. The results are shown in Table 1.

Table 1. Oil content of Leum Pua rice bran extracted oil.

Solvents	Extracted oil (g)	%Yield (g/100 g of dry basis)
Hexane	3.81±0.02	15.25±0.08
Ethanol	5.23±0.28	20.94±1.12
70:30 %v/v ethanol:water	5.47±0.09	21.89±0.35

Table 1 shows that the extracted oil content ranged from 15.25±0.08 to 21.89±0.35% by weight. Extraction using 70:30 %v/v ethanol:water yielded the highest oil content. Azrina et al. studied the rice bran oil extraction using 3:2 chloroform:methanol mixture obtaining result of 16.4% of yield [10]. The heating may disrupt the cellular wall, which results in greater porosity when the cell membrane ruptures, and improves the efficiency of the pressing extraction of oil from materials. In another research, the extraction yield of cold-pressed rice bran oil was 3.29-5.53% of yield [8].

3.2 Chemical properties of Leum Pua rice bran oil

Acid value (AV), free fatty acid (FFA) and peroxide value (PV) were the parameters used for determination of chemical quality of the extracted rice bran oil. Generally, the content of AV, FFA and PV was positively related to the activity of enzyme lipase. Acid value can be used for purity check of oil whereby decomposition reactions may have already progressed. The free fatty acids were lower than 5% in rice bran oil that was used for edible production. The high free fatty acid content of 10-20% was used in industrial grade of productions which could be present at considerate amounts in crude oils [11].

Table 2 shows the acid value, free fatty acid and peroxide value in this study. The results show that the lowest value of the chemical properties was determined by

using the 70:30 %v/v ethanol:water. Using hexane and ethanol as solvent extraction resulted in a slightly different value of the chemical properties. Furthermore, the 70:30 %v/v ethanol:water should be used as a solvent because it is less toxic to the environment than other solvents.

Raja Rajan and Gopala Krishina reported the refined rice bran oil had the free fatty acid in range 0.4-4.8% and that result was supported in this study [9]. The peroxide value indicated the quantity of peroxides in the oil which were important intermediates of oxidative reactions. The temperature of thermal extraction causes decomposition and free radicals formation. In table 2, the highest peroxide values are shown when the solvent extraction occurs at high temperature. Thanonkaew et al. reported the 11.72-12.13 meq Eqv O₂/kg oil of the peroxide value when using cold-pressed rice bran oil [8].

Table 2. The Chemical properties of Leum Pua rice bran extracted oil.

Solvents	Chemical properties of Leum Pua rice bran extracted oil		
	Acid value (mg KOH/g oil)	Free fatty acid (%)	Peroxide value (meq Eqv O ₂ /kg oil)
Hexane	4.97±0.68	2.49±0.34	21.50±0.50
Ethanol	4.85±0.31	2.42±0.16	19.83±0.58
70:30 %v/v ethanol:water	4.55±0.37	2.27±0.19	18.50±0.50

3.3 Gamma - oryzanol content of Leum Pua rice bran oil

Gamma-oryzanol can be used in nutraceutical, pharmaceutical and cosmoceutical productions [12]. In this research, the 5 ppm of gamma-oryzanol standard solution was used in scanning at the maximum wavelength (λ_{max}) which showed at 320 nm. The result found that the 0.4647 mg/L of the gamma-oryzanol content of extracted oil using 70:30 %v/v ethanol:water was the highest value shown in Table 3.

Table 3. Gamma - oryzanol content of Leum Pua rice bran oil.

Solvents	Gamma -oryzanol content (mg/L)
Hexane	0.0451±0.01
Ethanol	0.3872±0.04
Ethanol:Water (70:30)	0.4647±0.03

Thanonkaew et al. reported the 2.03 ±0.05 g/100g oil of the gamma-oryzanol when using cold-pressed rice bran oil [8]. Srikaeo found the concentration of gamma-oryzanol in RBO ranges from 115 to 780 ppm [13]. However, gamma-oryzanol content in rice bran oil is dependent on the temperature and method of processing [14].

4. Conclusion

The oil of Leum Pua rice bran was extracted by soxhlet extraction using hexane, ethanol and 70:30 %v/v ethanol:water as the solvents for 12 hr. The results found that the highest oil content 21.89±0.35 g/100g of dry basis was extracted by ethanol to water ratio 70:30. Acid value, free fatty acid and peroxide value of extracted oil were the lowest in oil extracted using ethanol to water ratio 70:30 as a solvent at 4.55±0.37 mgKOH/goil, 2.27±0.19% and 18.50±0.50 meq Eqv O₂/kg oil, respectively. The highest of the gamma-oryzanol content was 0.4647 mg/L of the extracted oil using ethanol to water ratio 70:30. Furthermore, the chemical properties could be indicative of the oil quality, which is the major cause of odour due to lipid peroxidation.

Acknowledgements

The authors acknowledge Miss Pornpimon Channai for assistance of this research. We also would like to thank the Faculty of Science and Technology, Phetchabun Rajabhat University for partial support of this research.

References

- [1] Pali V. Rice bran oil- Unique gift of nature:A review. *Agri. Reviews*. 2013;34(4):288-294.
- [2] Wongwaiwech D, Weerawatanakorn M, Tharatha S and Ho CT. Comparative study on amount of nutraceuticals in by-products from solvent and cold pressing methods of rice bran oil processing. *Journal of food and drug analysis*. 2019;27:71-82.
- [3] Jayadeep A and Malleshi NG. Nutrients, composition of tocotrienols, tocopherols, and c-oryzanol, and antioxidant activity in brown rice before and after biotransformation. *Journal of Food*. 2011;9(1):82–87.
- [4] Sakunpak A, Suksaeree J, Monton C, Pathompak P and Kraisintu K. Quantitative analysis of γ -oryzanol content in cold pressed rice bran oil by TLC-image analysis method. *Asian Pacific Journal of Tropical Biomedicine*. 2014;4(2):119-123.
- [5] Massarolo KC, Denardi de Souza T, Collazzo CC, Furlong EB and Almeida de Souza Soares L. The impact of *Rhizopus oryzae* cultivation on rice bran: Gamma-oryzanol recovery and its antioxidant properties. *Food Chemistry*. 2017;228:43-49.
- [6] Pornputtapitak W, Pantakitcharoenkul J, Panpakdee R, Teeranachaideekul V and Sinchaipanid N. Development of γ -Oryzanol Rich Extract from Leum Pua Glutinous Rice Bran Loaded Nanostructured Lipid Carriers for Topical Delivery. *Journal of Oleo Science*. 2018;67(2):125-133
- [7] Srisuwan S, Arkaravichien T, Mahatheeranont S, Puangsombat P, Seekhaw P, Na Lampang Noenplab A and Sattayasai J. Effects of Aqueous Extract of Unpolished Dark Purple Glutinous Rice, Var Luem Pua, on ROS in SK-N-SH Cells and Scopolamine-induced Memory Deficit in Mice. *Tropical Journal of Pharmaceutical Research* September. 2015;14(9):1635-1641

- [8] Thanonkaew A, Wongyai S, McClements DJ and Decker EA. Effect of stabilization of rice bran by domestic heating on mechanical extraction yield, quality, and antioxidant properties of cold-pressed rice bran oil (*Oryza sativa* L.). LWT-Food Science and Technology. 2012; 48:231-236.
- [9] Raja Rajan RG and Gopala Krishna AG. Refining of high free fatty acid rice bran oil its quality characteristics. Journal of food lipid. 2009;16(4):589-604.
- [10] Azrina A, Maznah I and Azizah AH. Extraction and Determination of Oryzanol in Rice Bran of Mixed Herbarium UKMB; AZ 6807:MR 185, AZ 6808: MR 211, AZ6809: MR 29. ASEAN Food Journal. 2008;15(1):89-96.
- [11] Gopala Krishna AG. Nutritional components of rice bran oil in relation to processing. Lipid Technology. 2002:80-84.
- [12] Patel M and Naik SN. Gamma-oryzanol from rice bran oil – A review. Journal of Scientific & Industrial Research. 2004;63:569-578.
- [13] Srikaeo K. Chapter 35 - Organic Rice Bran Oils in Health. Wheat and Rice in Disease Prevention and Health. 2014: 453-465.
- [14] Srisaipet A and Nuddagul M. Influence of temperature on gamma-oryzanol stability of edible rice bran oil during heating. International Journal of Chemical Engineering and Applications. 2014;5:303-306.